



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁶ :

C07K 14/415, C12N 15/29, C12Q 1/68

A1

(11) International Publication Number:

WO 95/28423

(43) International Publication Date:

26 October 1995 (26.10.95)

(21) International Application Number: PCT/US95/04589

(22) International Filing Date: 13 April 1995 (13.04.95)

(30) Priority Data:

08/227,360	13 April 1994 (13.04.94)	US
08/310,912	22 September 1994 (22.09.94)	US

(71) Applicants: THE GENERAL HOSPITAL CORPORATION [US/US]; 55 Fruit Street, Boston, MA 02114 (US). THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 22nd floor, 300 Lakeside Drive, Oakland, CA 94612-3550 (US). THE UNITED STATES OF AMERICA, represented by THE SECRETARY OF AGRICULTURE [US/US]; United States Department of Agriculture, 14th & Independence Avenue, SW, Washington, DC 20250 (US). COMMONWEALTH SCIENTIFIC AND INDUSTRIAL RESEARCH ORGANISATION [AU/AU]; 407 Royal Parade, Parkville, VIC 3052 (AU).

(72) Inventors: AUSUBEL, Frederick, M.; 271 Lake Avenue, Newton, MA 02161 (US). STASKAWICZ, Brian, J.; 18945 Marciel Court, Castro Valley, CA 94546 (US). BENT, Andrew, F.; 308 West Iowa Street, Urbana, IL 61801 (US). DAHLBECK, Douglas; 18851 Lenross Court, Castro Valley, CA 94546 (US). KATAGIRI, Fumiaki; Seven Hardan Road, Somerville, MA 02144 (US). KUNKEL, Barbara, N.; 6042 Pershing Avenue, St. Louis, MO 63112 (US). MINDRINOS, Michael, N.; 400 Brookline Avenue, Boston, MA 02215 (US). YU, Guo-Liang; 13524 Straw Bale Lane, Darnestown, MD 20878 (US). BAKER, Barbara; 804 Vista Heights Road, Richmond, CA 94805 (US). ELLIS, Jeffrey; 10 Gibbs Place, Weetangera 2614 (AU). SALMERON, John; 1308 Blackberry Lane, Hillsborough, NC 27278 (US).

(74) Agent: CLARK, Paul, T.; Fish & Richardson, 225 Franklin Street, Boston, MA 02110-2804 (US).

(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TT, UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).

Published

With international search report.

(54) Title: RPS GENE FAMILY, PRIMERS, PROBES, AND DETECTION METHODS

(57) Abstract

Disclosed is substantially pure DNA encoding an *Arabidopsis thaliana* Rps2 polypeptide; substantially pure Rps2 polypeptide; and methods of using such DNA to express the Rps2 polypeptide in plant cells and whole plants to provide, in transgenic plants, disease resistance to pathogens. Also disclosed are conserved regions characteristic of the RPS family and primers and probes for the identification and isolation of additional RPS disease-resistance genes.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

RPS GENE FAMILY, PRIMERS, PROBES, AND DETECTION METHODSStatement as to Federally Sponsored Research

5 This invention was made in part with Government funding and the Government therefore has certain rights in the invention.

Background of the Invention

10 The invention relates to recombinant plant nucleic acids and polypeptides and uses thereof to confer disease resistance to pathogens in transgenic plants.

Plants employ a variety of defensive strategies to combat pathogens. One defense response, the so-called hypersensitive response (HR), involves rapid localized
15 necrosis of infected tissue. In several host-pathogen interactions, genetic analysis has revealed a gene-for-gene correspondence between a particular avirulence (*avr*) gene in an avirulent pathogen that elicits an HR in a host possessing a particular resistance gene.

20 Summary of the Invention

In general, the invention features substantially pure DNA (for example, genomic DNA, cDNA, or synthetic DNA) encoding an Rps polypeptide as defined below. In related aspects, the invention also features a vector, a
25 cell (e.g., a plant cell), and a transgenic plant or seed thereof which includes such a substantially pure DNA encoding an Rps polypeptide.

In preferred embodiments, an RPS gene is the RPS2 gene of a plant of the genus *Arabidopsis*. In various
30 preferred embodiments, the cell is a transformed plant cell derived from a cell of a transgenic plant. In related aspects, the invention features a transgenic plant containing a transgene which encodes an Rps

- 2 -

polypeptide that is expressed in plant tissue susceptible to infection by pathogens expressing the *avrRpt2* avirulence gene or pathogens expressing an avirulence signal similarly recognized by an *Rps* polypeptide.

5 In a second aspect, the invention features a substantially pure DNA which includes a promoter capable of expressing the *RPS2* gene in plant tissue susceptible to infection by bacterial pathogens expressing the *avrRpt2* avirulence gene.

10 In preferred embodiments, the promoter is the promoter native to an *RPS* gene. Additionally, transcriptional and translational regulatory regions are preferably native to an *RPS* gene.

The transgenic plants of the invention are
15 preferably plants which are susceptible to infection by a pathogen expressing an avirulence gene, preferably the *avrRpt2* avirulence gene. In preferred embodiments the transgenic plant is from the group of plants consisting of but not limited to *Arabidopsis*, tomato, soybean, bean,
20 maize, wheat and rice.

In another aspect, the invention features a method of providing resistance in a plant to a pathogen which involves: (a) producing a transgenic plant cell having a transgene encoding an *Rps2* polypeptide wherein the
25 transgene is integrated into the genome of the transgenic plant and is positioned for expression in the plant cell; and (b) growing a transgenic plant from the transgenic plant cell wherein the *RPS2* transgene is expressed in the transgenic plant.

30 In another aspect, the invention features a method of detecting a resistance gene in a plant cell involving: (a) contacting the *RPS2* gene or a portion thereof greater than 9 nucleic acids, preferably greater than 18 nucleic acids in length with a preparation of genomic DNA from
35 the plant cell under hybridization conditions providing

- 3 -

detection of DNA sequences having about 50% or greater sequence identity to the DNA sequence of Fig. 2 encoding the Rps2 polypeptide.

In another aspect, the invention features a method of producing an Rps2 polypeptide which involves: (a) providing a cell transformed with DNA encoding an Rps2 polypeptide positioned for expression in the cell; (b) culturing the transformed cell under conditions for expressing the DNA; and (c) isolating the Rps2 polypeptide.

In another aspect, the invention features substantially pure Rps2 polypeptide. Preferably, the polypeptide includes a greater than 50 amino acid sequence substantially identical to a greater than 50 amino acid sequence shown in Fig. 2, open reading frame "a". Most preferably, the polypeptide is the *Arabidopsis thaliana* Rps2 polypeptide.

In another aspect, the invention features a method of providing resistance in a transgenic plant to infection by pathogens which do not carry the *avrRpt2* avirulence gene wherein the method includes: (a) producing a transgenic plant cell having transgenes encoding an Rps2 polypeptide as well as a transgene encoding the *avrRpt2* gene product wherein the transgenes are integrated into the genome of the transgenic plant; are positioned for expression in the plant cell; and the *avrRpt2* transgene and, if desired, the *RPS2* gene, are under the control of regulatory sequences suitable for controlled expression of the gene(s); and (b) growing a transgenic plant from the transgenic plant cell wherein the *RPS2* and *avrRpt2* transgenes are expressed in the transgenic plant.

In another aspect, the invention features a method of providing resistance in a transgenic plant to infection by pathogens in the absence of avirulence gene

- 4 -

expression in the pathogen wherein the method involves:

- (a) producing a transgenic plant cell having integrated in the genome a transgene containing the *RPS2* gene under the control of a promoter providing constitutive
5 expression of the *RPS2* gene; and (b) growing a transgenic plant from the transgenic plant cell wherein the *RPS2* transgene is expressed constitutively in the transgenic plant.

- In another aspect, the invention features a method
10 of providing controllable resistance in a transgenic plant to infection by pathogens in the absence of avirulence gene expression in the pathogen wherein the method involves: (a) producing a transgenic plant cell having integrated in the genome a transgene containing
15 the *RPS2* gene under the control of a promoter providing controllable expression of the *RPS2* gene; and (b) growing a transgenic plant from the transgenic plant cell wherein the *RPS2* transgene is controllably expressed in the transgenic plant. In preferred embodiments, the *RPS2*
20 gene is expressed using a tissue-specific or cell type-specific promoter, or by a promoter that is activated by the introduction of an external signal or agent, such as a chemical signal or agent.

- In other aspects, the invention features a
25 substantially pure oligonucleotide including one or a combination of the sequences:

5' GGNATGGGNGGNNTNGGNAARACNAC 3', [SEQ ID NO: 158]
wherein N is A, T, G, or C; and R is A or G;

- 5' NARNGGNARNCC 3', [SEQ ID NO: 169] wherein N is
30 A, T, G or C; and R is A or G;

5'NCGNGWNGTNAKDAWNCGNGA 3', [SEQ ID NO: 159]
wherein N is A, T, G or C; W is A or T; D is A, G, or T;
and K is G or T;

- 5 -

5' GGWNTBGGWAARACHAC 3', [SEQ ID NO: 160] wherein N is A, T, G or C; R is G or A; B is C, G, or T; H is A, C, or T; and W is A or T;

5' TYGAYGAYRTBKRBRA 3', [SEQ ID NO: 163] wherein R is G or A; B is C, G, or T; D is A, G, or T; Y is T or C; and K is G or T;

5' TYCCAVAYRTCRTCNA 3', [SEQ ID NO: 164] wherein N is A, T, G or C; R is G or A; V is G or C or A; and Y is T or C;

10 5' GGWYTBCCWYTBGCHYT 3', [SEQ ID NO: 170] wherein B is C, G, or T; H is A, C, or T; W is A or T; and Y is T or C;

5' ARDGCVARWGGVARNCC 3', [SEQ ID NO: 171] wherein N is A, T, G or C; R is G or A; W is A or T; D is A, G, or T; and V is G, C, or A; and

15 5' ARRTTRTCRTADSWRAWYTT 3', [SEQ ID NO: 174] wherein R is G or A; W is A or T; D is A, G, or T; S is G or C; and Y is C or T.

In other aspects, the invention features a recombinant plant gene including one or a combination of the DNA sequences:

5' GGNATGGGNGGNNTNGGNAARACNAC 3', [SEQ ID NO: 162] wherein N is A, T, G or C; and R is A or G;

5' NARNGGNARNCC 3', [SEQ ID NO: 169] wherein N is A, T, G or C; and R is A or G;

25 5' NCGNGWNGTNAKDAWNCNGA 3', [SEQ ID NO: 167] wherein N is A, T, G or C; W is A or T; D is A, G or T; and K is G or T.

In another aspect, the invention features a substantially pure plant polypeptide including one or a combination of the amino acid sequences:

Gly Xaa₁ Xaa₂ Gly Xaa₃ Gly Lys Thr Thr Xaa₄ Xaa₅, [SEQ ID NO: 191] wherein Xaa₁ is Met or Pro; Xaa₂ is Gly or Pro; Xaa₃ is Ile, Leu, or Val; Xaa₄ is Ile, Leu, or Thr; and Xaa₅ is Ala or Met;

- 6 -

Xaa₁ Xaa₂ Xaa₃ Leu Xaa₄ Xaa₅ Xaa₆ Asp Asp Xaa₇
Xaa₈, [SEQ ID NO; 192]

wherein Xaa₁ is Phe or Lys; Xaa₂ is Arg or Lys; Xaa₃ is
Ile, Val, or Phe; Xaa₄ is Ile, Leu, or Val; Xaa₅ is Ile or
5 Leu; Xaa₆ is Ile or Val; Xaa₇ is Ile, Leu, or Val; and
Xaa₈ is Asp or Trp;

Xaa₁ Xaa₂ Xaa₃ Xaa₄ Xaa₅ Thr Xaa₆ Arg, [SEQ ID NO:
193]

wherein Xaa₁ is Ser or Cys; Xaa₂ is Arg or Lys; Xaa₃ is
10 Phe, Ile, or Val; Xaa₄ is Ile, or Met; Xaa₅ is Ile, Leu,
or Phe; Xaa₆ is Ser, Cys, or Thr;

Gly Leu Pro Leu Xaa₁ Xaa₂ Xaa₃ Xaa₄, [SEQ ID NO.:
194]

wherein Xaa₁ is Thr, Ala, or Ser; Xaa₂ is Leu or Val; Xaa₃
15 is Ile, Val, or Lys; and Xaa₄ is Val or Thr; and

Xaa₁ Xaa₂ Ser Tyr Xaa₃ Xaa₄ Leu, [SEQ ID NO: 195]
wherein Xaa₁ is Lys or Gly; Xaa₂ is Ile or Phe; Xaa₃ is
Asp or Lys; and Xaa₄ is Ala, Gly, or Asn.

In another aspect, the invention features a method
20 of isolating a disease-resistance gene or fragment
thereof from a plant cell, involving: (a) providing a
sample of plant cell DNA; (b) providing a pair of
oligonucleotides having sequence homology to a conserved
region of an RPS disease-resistance gene; (c) combining
25 the pair of oligonucleotides with the plant cell DNA
sample under conditions suitable for polymerase chain
reaction-mediated DNA amplification; and (d) isolating
the amplified disease-resistance gene or fragment
thereof.

- 7 -

In preferred embodiments, the amplification is carried out using a reverse-transcription polymerase chain reaction, for example, the RACE method

In another aspect, the invention features a method
5 of identifying a plant disease-resistance gene in a plant cell, involving: (a) providing a preparation of plant cell DNA (for example, from the plant genome); (b) providing a detectably-labelled DNA sequence (for example, prepared by the methods of the invention) having
10 homology to a conserved region of an RPS gene; (c) contacting the preparation of plant cell DNA with the detectably-labelled DNA sequence under hybridization conditions providing detection of genes having 50% or greater sequence identity; and (d) identifying a disease-
15 resistance gene by its association with the detectable label.

In another aspect, the invention features a method of isolating a disease-resistance gene from a recombinant plant cell library, involving: (a) providing a
20 recombinant plant cell library; (b) contacting the recombinant plant cell library with a detectably-labelled gene fragment produced according to the PCR method of the invention under hybridization conditions providing detection of genes having 50% or greater sequence
25 identity; and (c) isolating a member of a disease-resistance gene by its association with the detectable label.

In another aspect, the invention features a method of isolating a disease-resistance gene from a recombinant
30 plant cell library, involving: (a) providing a recombinant plant cell library; (b) contacting the recombinant plant cell library with a detectably-labelled RPS oligonucleotide of the invention under hybridization conditions providing detection of genes having 50% or
35 greater sequence identity; and (c) isolating a disease-

- 8 -

resistance gene by its association with the detectable label.

In another aspect, the invention features a recombinant plant polypeptide capable of conferring
5 disease-resistance wherein the plant polypeptide includes a P-loop domain or nucleotide binding site domain. Preferably, the polypeptide further includes a leucine-rich repeating domain.

In another aspect, the invention features a
10 recombinant plant polypeptide capable of conferring disease-resistance wherein the plant polypeptide contains a leucine-rich repeating domain.

In another aspect, the invention features a plant disease-resistance gene isolated according to the method
15 involving: (a) providing a sample of plant cell DNA; (b) providing a pair of oligonucleotides having sequence homology to a conserved region of an RPS disease-resistance gene; (c) combining the pair of
oligonucleotides with the plant cell DNA sample under
20 conditions suitable for polymerase chain reaction-mediated DNA amplification; and (d) isolating the amplified disease-resistance gene or fragment thereof.

In another aspect, the invention features a plant disease-resistance gene isolated according to the method
25 involving: (a) providing a preparation of plant cell DNA; (b) providing a detectably-labelled DNA sequence having homology to a conserved region of an RPS gene; (c) contacting the preparation of plant cell DNA with the detectably-labelled DNA sequence under hybridization
30 conditions providing detection of genes having 50% or greater sequence identity; and (d) identifying a disease-resistance gene by its association with the detectable label.

In another aspect, the invention features a plant
35 disease-resistance gene according to the method

- 9 -

involving: (a) providing a recombinant plant cell library; (b) contacting the recombinant plant cell library with a detectably-labelled RPS gene fragment produced according to the method of the invention under
5 hybridization conditions providing detection of genes having 50% or greater sequence identity; and (c) isolating a disease-resistance gene by its association with the detectable label.

In another aspect, the invention features a method
10 of identifying a plant disease-resistance gene involving: (a) providing a plant tissue sample; (b) introducing by biolistic transformation into the plant tissue sample a candidate plant disease-resistance gene; (c) expressing the candidate plant disease-resistance gene within the
15 plant tissue sample; and (d) determining whether the plant tissue sample exhibits a disease-resistance response, whereby a response identifies a plant disease-resistance gene.

Preferably, the plant tissue sample is either
20 leaf, root, flower, fruit, or stem tissue; the candidate plant disease-resistance gene is obtained from a cDNA expression library; and the disease-resistance response is the hypersensitive response.

In another aspect, the invention features a plant
25 disease-resistance gene isolated according to the method involving: (a) providing a plant tissue sample; (b) introducing by biolistic transformation into the plant tissue sample a candidate plant disease-resistance gene; (c) expressing the candidate plant disease-resistance
30 gene within the plant tissue sample; and (d) determining whether the plant tissue sample exhibits a disease-resistance response, whereby a response identifies a plant disease-resistance gene.

In another aspect, the invention features a
35 purified antibody which binds specifically to an rps

- 10 -

family protein. Such an antibody may be used in any standard immunodetection method for the identification of an RPS polypeptide.

In another aspect, the invention features a DNA
5 sequence substantially identical to the DNA sequence shown in Figure 12.

In another aspect, the invention features a substantially pure polypeptide having a sequence substantially identical to a Prf amino acid sequence
10 shown in Figure 5 (A or B).

By "disease resistance gene" is meant a gene encoding a polypeptide capable of triggering the plant defense response in a plant cell or plant tissue. An RPS gene is a disease resistance gene having about 50% or
15 greater sequence identity to the RPS2 sequence of Fig. 2 or a portion thereof. The gene, RPS2, is a disease resistance gene encoding the Rps2 disease resistance polypeptide from *Arabidopsis thaliana*.

By "polypeptide" is meant any chain of amino
20 acids, regardless of length or post-translational modification (e.g., glycosylation or phosphorylation).

By "substantially identical" is meant a polypeptide or nucleic acid exhibiting at least 50%, preferably 85%, more preferably 90%, and most preferably
25 95% homology to a reference amino acid or nucleic acid sequence. For polypeptides, the length of comparison sequences will generally be at least 16 amino acids, preferably at least 20 amino acids, more preferably at least 25 amino acids, and most preferably 35 amino acids.
30 For nucleic acids, the length of comparison sequences will generally be at least 50 nucleotides, preferably at least 60 nucleotides, more preferably at least 75 nucleotides, and most preferably 110 nucleotides.

Sequence identity is typically measured using
35 sequence analysis software (e.g., Sequence Analysis

- 11 -

Software Package of the Genetics Computer Group,
University of Wisconsin Biotechnology Center, 1710
University Avenue, Madison, WI 53705). Such software
matches similar sequences by assigning degrees of
5 homology to various substitutions, deletions,
substitutions, and other modifications. Conservative
substitutions typically include substitutions within the
following groups: glycine alanine; valine, isoleucine,
leucine; aspartic acid, glutamic acid, asparagine,
10 glutamine; serine, threonine; lysine, arginine; and
phenylalanine, tyrosine.

By a "substantially pure polypeptide" is meant an
Rps2 polypeptide which has been separated from components
which naturally accompany it. Typically, the polypeptide
15 is substantially pure when it is at least 60%, by weight,
free from the proteins and naturally-occurring organic
molecules with which it is naturally associated.
Preferably, the preparation is at least 75%, more
preferably at least 90%, and most preferably at least
20 99%, by weight, Rps2 polypeptide. A substantially pure
Rps2 polypeptide may be obtained, for example, by
extraction from a natural source (e.g., a plant cell); by
expression of a recombinant nucleic acid encoding an Rps2
polypeptide; or by chemically synthesizing the protein.
25 Purity can be measured by any appropriate method, e.g.,
those described in column chromatography, polyacrylamide
gel electrophoresis, or by HPLC analysis.

A protein is substantially free of naturally
associated components when it is separated from those
30 contaminants which accompany it in its natural state.
Thus, a protein which is chemically synthesized or
produced in a cellular system different from the cell
from which it naturally originates will be substantially
free from its naturally associated components.
35 Accordingly, substantially pure polypeptides include

- 12 -

those derived from eukaryotic organisms but synthesized in *E. coli* or other prokaryotes.

By "substantially pure DNA" is meant DNA that is free of the genes which, in the naturally-occurring
5 genome of the organism from which the DNA of the invention is derived, flank the gene. The term therefore includes, for example, a recombinant DNA which is incorporated into a vector; into an autonomously replicating plasmid or virus; or into the genomic DNA of
10 a prokaryote or eukaryote; or which exists as a separate molecule (e.g., a cDNA or a genomic or cDNA fragment produced by PCR or restriction endonuclease digestion) independent of other sequences. It also includes a recombinant DNA which is part of a hybrid gene encoding
15 additional polypeptide sequence.

By "transformed cell" is meant a cell into which (or into an ancestor of which) has been introduced, by means of recombinant DNA techniques, a DNA molecule encoding (as used herein) an Rps2 polypeptide.

20 By "positioned for expression" is meant that the DNA molecule is positioned adjacent to a DNA sequence which directs transcription and translation of the sequence (i.e., facilitates the production of, e.g., an Rps2 polypeptide, a recombinant protein or a RNA
25 molecule).

By "reporter gene" is meant a gene whose expression may be assayed; such genes include, without limitation, β -glucuronidase (GUS), luciferase, chloramphenicol transacetylase (CAT), and β -
30 galactosidase.

By "promoter" is meant minimal sequence sufficient to direct transcription. Also included in the invention are those promoter elements which are sufficient to render promoter-dependent gene expression controllable
35 for cell-type specific, tissue-specific or inducible by

- 13 -

external signals or agents; such elements may be located in the 5' or 3' regions of the native gene.

By "operably linked" is meant that a gene and a regulatory sequence(s) are connected in such a way as to
5 permit gene expression when the appropriate molecules (e.g., transcriptional activator proteins) are bound to the regulatory sequence(s).

By "plant cell" is meant any self-propagating cell bounded by a semi-permeable membrane and containing a
10 plastid. Such a cell also requires a cell wall if further propagation is desired. Plant cell, as used herein includes, without limitation, algae, cyanobacteria, seeds suspension cultures, embryos, meristematic regions, callus tissue, leaves, roots, shoots, gametophytes,
15 sporophytes, pollen, and microspores.

By "transgene" is meant any piece of DNA which is inserted by artifice into a cell, and becomes part of the genome of the organism which develops from that cell. Such a transgene may include a gene which is partly or
20 entirely heterologous (i.e., foreign) to the transgenic organism, or may represent a gene homologous to an endogenous gene of the organism.

By "transgenic" is meant any cell which includes a DNA sequence which is inserted by artifice into a cell
25 and becomes part of the genome of the organism which develops from that cell. As used herein, the transgenic organisms are generally transgenic plants and the DNA (transgene) is inserted by artifice into the nuclear or plastidic genome.

30 By "pathogen" is meant an organism whose infection into the cells of viable plant tissue elicits a disease response in the plant tissue.

By an "RPS disease-resistance gene" is meant any member of the family of plant genes characterized by
35 their ability to trigger a plant defense response and

- 14 -

having at least 20%, preferably 30%, and most preferably 50% amino acid sequence identity to one of the conserved regions of one of the RPS members described herein (i.e., either the RPS2, L6, N, or Prf genes). Representative
5 members of the RPS gene family include, without limitation, the rps2 gene of *Arabidopsis*, the L6 gene of flax, the Prf gene of tomato, and the N gene of tobacco.

By "conserved region" is meant any stretch of six or more contiguous amino acids exhibiting at least 30%,
10 preferably 50%, and most preferably 70% amino acid sequence identity between two or more of the RPS family members, RPS2, L6, N, or Prf. Examples of preferred conserved regions are shown (as boxed or designated sequences) in Figures 5 A and B, 6, 7, and 8 and include,
15 without limitation, nucleotide binding site domains, leucine-rich repeats, leucine zipper domains, and P-loop domains.

By "detectably-labelled" is meant any means for marking and identifying the presence of a molecule, e.g.,
20 an oligonucleotide probe or primer, a gene or fragment thereof, or a cDNA molecule. Methods for detectably-labelling a molecule are well known in the art and include, without limitation, radioactive labelling (e.g., with an isotope such as ³²P or ³⁵S) and nonradioactive
25 labelling (e.g., chemiluminescent labelling, e.g., fluorescein labelling).

By "biolistic transformation" is meant any method for introducing foreign molecules into a cell using velocity driven microprojectiles such as tungsten or gold
30 particles. Such velocity-driven methods originate from pressure bursts which include, but are not limited to, helium-driven, air-driven, and gunpowder-driven techniques. Biolistic transformation may be applied to the transformation or transfection of a wide variety of
35 cell types and intact tissues including, without

- 15 -

limitation, intracellular organelles (e.g., chloroplasts and mitochondria), bacteria, yeast, fungi, algae, pollen, animal tissue, plant tissue (e.g., leaf, seedling, embryo, epidermis, flower, meristem, and root), pollen, and cultured cells.

By "purified antibody" is meant antibody which is at least 60%, by weight, free from proteins and naturally-occurring organic molecules with which it is naturally associated. Preferably, the preparation is at least 75%, more preferably 90%, and most preferably at least 99%, by weight, antibody, e.g., an rps2-specific antibody. A purified rps antibody may be obtained, for example, by affinity chromatography using recombinantly-produced rps protein or conserved motif peptides and standard techniques.

By "specifically binds" is meant an antibody which recognizes and binds an rps protein but which does not substantially recognize and bind other molecules in a sample, e.g., a biological sample, which naturally includes rps protein.

Other features and advantages of the invention will be apparent from the following description of the preferred embodiments thereof, and from the claims.

Detailed Description

The drawings will first be described.

Drawings

Figs. 1A - 1F are a schematic summary of the physical and RFLP analysis that led to the cloning of the RPS2 locus.

Fig. 1A is a diagram showing the alignment of the genetic and the RFLP maps of the relevant portion of *Arabidopsis thaliana* chromosome IV adapted from the map published by Lister and Dean (1993) Plant J. 4:745-750.

- 16 -

The RFLP marker L11F11 represents the left arm of the YUP11F11 YAC clone.

Fig. 1B is a diagram showing the alignment of relevant YACs around the *RPS2* locus. YAC constructs designated YUP16G5, YUP18G9 and YUP11F11 were provided by J. Ecker, University of Pennsylvania. YAC constructs designated EW3H7, EW11D4, EW11E4, and EW9C3 were provided by E. Ward, Ciba-Geigy, Inc.

Fig. 1C is a diagram showing the alignment of cosmid clones around the *RPS2* locus. Cosmid clones with the designation H are derivatives of the EW3H7 YAC clone whereas those with the designation E are derivatives of the EW11E4 YAC clone. Vertical arrows indicate the relative positions of RFLP markers between the ecotypes La-er and the *rps2-101N* plant. The RFLP markers were identified by screening a Southern blot containing more than 50 different restriction enzyme digests using either the entire part or pieces of the corresponding cosmid clones as probes. The cosmid clones described in Fig. 1C were provided by J. Giraudat, C.N.R.S., Gif-sur-Yvette, France.

Figs. 1D and 1E are maps of *EcoRI* restriction endonuclease sites in the cosmids E4-4 and E4-6, respectively. The recombination break points surrounding the *RPS2* locus are located within the 4.5 and 7.5 kb *EcoRI* restriction endonuclease fragments.

Fig. 1F is a diagram showing the approximate location of genes which encode the RNA transcripts which have been identified by polyA⁺ RNA blot analysis. The sizes of the transcripts are given in kilobase pairs below each transcript.

Fig. 2 [SEQ ID NOS; 1-104, 196-201] is the complete nucleotide sequence of cDNA-4 comprising the *RPS2* gene locus. The three reading frames are shown below the nucleotide sequence. The deduced amino acid

- 17 -

sequence of reading frame "a" is provided and contains 909 amino acids. The methionine encoded by the ATG start codon is circled in open reading frame "a" of Fig. 2. The A of the ATG start codon is nucleotide 31 of Fig. 2.

5 Fig. 3 [SEQ ID NOS: 105-106] is the nucleotide sequence of the *avrRpt2* gene and its deduced amino acid sequence. A potential ribosome binding site is underlined. An inverted repeat is indicated by horizontal arrows at the 3' end of the open reading
10 frame. The deduced amino acid sequence is provided below the nucleotide sequence of the open reading frame.

 Fig. 4 is a schematic summary of the complementation analysis that allowed functional confirmation that the DNA carried on p4104 and p4115
15 (encoding cDNA-4) confers RPS2 disease resistance activity to *Arabidopsis thaliana* plants previously lacking RPS2 disease resistance activity. Small vertical marks along the "genome" line represent restriction enzyme *EcoRI* recognition sites, and the numbers above
20 this line represent the size, in kilobase pairs (kb), of the resulting DNA fragments (see also Fig. 1E). Opposite "cDNAs" are the approximate locations of the coding sequences for RNA transcripts (See also Fig. 1F); arrowheads indicate the direction of transcription for
25 cDNAs 4, 5, and 6. For functional complementation experiments, *rps2-201C/rps2-201C* plants were genetically transformed with the *Arabidopsis thaliana* genomic DNA sequences indicated; these sequences were carried on the named plasmids (derivatives of the binary cosmid vector
30 pSLJ4541) and delivered to the plant via *Agrobacterium*-mediated transformation methods. The disease resistance phenotype of the resulting transformants following inoculation with *P. syringae* expressing *avrRpt2* is given as "Sus." (susceptible, no resistance response) or "Res."
35 (disease resistant).

- 18 -

Fig. 5A [SEQ ID NOS: 107-136;, AND 142] shows regions of sequence similarity between the L-6 protein of flax, N protein of tobacco, Prf protein of tomato, and rps2 protein of *Arabidopsis*.

5 Fig. 5B [SEQ ID NOS: 107, 108, 137-140] shows sequence similarity between the N and L-6 proteins.

Fig. 6 [SEQ ID NOS: 141 and 142] shows a sequence analysis of RPS2 polypeptide showing polypeptide regions corresponding to an N-terminal hydrophobic region, a
10 leucine zipper, NBSs (kinase-1a, kinase-2, and kinase-3 motifs), and a predicted membrane integrated region.

Fig. 7 [SEQ ID NOS: 143-146] shows the amino acid sequence of the RPS2 LRR (amino acids 505-867). The top line indicates the consensus sequences for the RPS2 LRR.
15 An "X" stands for an arbitrary amino acid sequence and an "a" stands for an aliphatic amino acid residue. The consensus sequence for the RPS2 LRR is closely related to the consensus for the yeast adenylate cyclase CYR1 LRR (PX Xa XXL XXL XXLXL XXNXaXXa). The amino acid residues
20 that match the consensus sequence are shown in bold. Although this figure shows 14 LRRs, the C-terminal boundary of the LRR is not very clear because the LRR closer to the C-terminus does not fit the consensus sequence very well.

25 Fig. 8 [SEQ ID NO: 3] shows a sequence analysis of RPS2, indicating regions with similarity to leucine zipper, P-loop, membrane-spanning, and leucine-rich repeat motifs. Regions with similarity to defined functional domains are indicated with a line over the
30 relevant amino acids. Potential N-glycosylation sequences are marked with a dot, and the location of the rps2-201 Thr to Pro mutation at amino acid 668 is marked with an asterisk.

Fig. 9 is a schematic representation of the
35 transient assay method. The top panel shows the

- 19 -

essential principles of the assay. The bottom panel shows a schematic representation of the actual transient assay procedure. *Psp* NP53121 is used because it is a weak *Arabidopsis* pathogen, but potent in causing the HR when carrying an avirulence gene. In the absence of an HR, the damage to plant cells infected with NP53121 is minimal, enhancing the difference of GUS accumulation in cells that undergo the HR in comparison to those that do not. Prior to bombardment, one half of an *Arabidopsis* leaf is infiltrated with *P. syringae* (stippled side of leaf); the other half of the leaf serves as a noninfected control, an "internal" reference for the infected side, and as a measure of transformation efficiency.

Fig. 10, panels A-B, are photographs showing the complementation of the *rps2* mutant phenotype using the biolistic transient expression assay. The left sides of *rps2*-101C mutant leaves were infiltrated with *Psp* 3121/*avrRpt2*. Infiltrated leaves were cobombarded with either 35S-uidA plus AGUS (Panel A) or 35S-uidA plus 35S-RPS2 (cDNA-2 clone 4) (Panel B). Note that in Panel B the infected side of the leaf shows less GUS activity than the uninfected side, indicating that the transformed cells on the infected side underwent an HR and that 35S-RPS2 complemented the mutant phenotype (see Fig. 9).

Fig. 11 is a schematic representation of pKEx4tr showing the structure of this cDNA expression vector. For convenience, the multiple cloning site contains the 8bp recognition sequences for *Pme*I and *Not*I and is flanked by T7 and T3 promoters. The region spanning the modified 35S promoter to the nopaline synthase 3' sequences (nos 3') was cloned into the *Hind* III-*Eco*RI site of pUC18, resulting in the loss of the *Eco*RI site.

Fig. 12 shows a nucleic acid sequence of the tomato *Prf* gene.

- 20 -

The Genetic Basis for Resistance to Pathogens

An overview of the interaction between a plant host and a microbial pathogen is presented. The invasion of a plant by a potential pathogen can have a range of outcomes delineated by the following outcomes: either the pathogen successfully proliferates in the host, causing associated disease symptoms, or its growth is halted by the host defenses. In some plant-pathogen interactions, the visible hallmark of an active defense response is the so-called hypersensitive response or "HR". The HR involves rapid necrosis of cells near the site of the infection and may include the formation of a visible dry brown lesion. Pathogens which elicit an HR on a given host are said to be avirulent on that host, the host is said to be resistant, and the plant-pathogen interaction is said to be incompatible. Strains which proliferate and cause disease on a particular host are said to be virulent; in this case the host is said to be susceptible, and the plant-pathogen interaction is said to be compatible.

"Classical" genetic analysis has been used successfully to help elucidate the genetic basis of plant-pathogen recognition for those cases in which a series of strains (races) of a particular fungal or bacterial pathogen are either virulent or avirulent on a series of cultivars (or different wild accessions) of a particular host species. In many such cases, genetic analysis of both the host and the pathogen revealed that many avirulent fungal and bacterial strains differ from virulent ones by the possession of one or more avirulence (avr) genes that have corresponding "resistance" genes in the host. This avirulence gene-resistance gene correspondence is termed the "gene-for-gene" model (Crute, et al., (1985) pp 197-309 in: *Mechanisms of Resistance to Plant Disease*. R.S.S. Fraser, ed.;

- 21 -

Ellingboe, (1981) *Annu. Rev. Phytopathol.* 19:125-143;
Flor, (1971) *Annu. Rev. Phytopathol.* 9:275-296; Keen and
Staskawicz, (1988) *supra*; and Keen et al. in: *Application
of Biotechnology to Plant Pathogen Control*. I. Chet, ed.,
5 John Wiley & Sons, 1993, pp. 65-88). According to a
simple formulation of this model, plant resistance genes
encode specific receptors for molecular signals generated
by avr genes. Signal transduction pathway(s) then carry
the signal to a set of target genes that initiate the HR
10 and other host defenses (Gabriel and Rolfe, (1990) *Annu.
Rev. Phytopathol.* 28:365-391). Despite this simple
predictive model, the molecular basis of the avr-
resistance gene interaction is still unknown.

One basic prediction of the gene-for-gene
15 hypothesis has been convincingly confirmed at the
molecular level by the cloning of a variety of bacterial
avr genes (Innes, et al., (1993) *J. Bacteriol.* 175:4859-
4869; Dong, et al., (1991) *Plant Cell* 3:61-72; Whelan et
al., (1991) *Plant Cell* 3:49-59; Staskawicz et al., (1987)
20 *J. Bacteriol.* 169:5789-5794; Gabriel et al., (1986)
P.N.A.S., USA 83:6415-6419; Keen and Staskawicz, (1988)
Annu. Rev. Microbiol. 42:421-440; Kobayashi et al.,
(1990) *Mol. Plant-Microbe Interact.* 3:94-102 and (1990)
Mol. Plant-Microbe Interact. 3:103-111). Many of these
25 cloned avirulence genes have been shown to correspond to
individual resistance genes in the cognate host plants
and have been shown to confer an avirulent phenotype when
transferred to an otherwise virulent strain. The *avrRpt2*
locus was isolated from *Pseudomonas syringae* pv. *tomato*
30 and sequenced by Innes et al. (Innes, R. et al. (1993) *J.
Bacteriol.* 175:4859-4869). Fig. 3 is the nucleotide
sequence and deduced amino acid sequence of the *avrRpt2*
gene.

Examples of known signals to which plants respond
35 when infected by pathogens include harpins from *Erwinia*

- 22 -

- (Wei et al. (1992) Science 257:85-88) and *Pseudomonas* (He et al. (1993) Cell 73:1255-1266); *avr4* (Joosten et al. (1994) Nature 367:384-386) and *avr9* peptides (van den Ackerveken et al (1992) Plant J. 2:359-366) from
- 5 *Cladosporium*; *PopA1* from *Pseudomonas* (Arlat et al. (1994) EMBO J. 13:543-553); *avrD*-generated lipopolysaccharide (Midland et al. (1993) J. Org. Chem. 58:2940-2945); and *NIP1* from *Rhynchosporium* (Hahn et al. (1993) Mol. Plant-Microbe Interact. 6:745-754).
- 10 Compared to *avr* genes, considerably less is known about plant resistance genes that correspond to specific *avr*-generated signals. The plant resistance gene, *RPS2* (*rps* for resistance to P*seudomonas* s*yringae*), the first
- 15 disease resistance genes corresponds to a specific *avr* gene (*avrRpt2*). Some of the work leading up to the cloning of *RPS2* is described in Yu, et al., (1993), Molecular Plant-Microbe Interactions 6:434-443 and in Kunkel, et al., (1993) Plant Cell 5:865-875.
- 20 An apparently unrelated avirulence gene which corresponds specifically to plant disease resistance gene, *Pto*, has been isolated from tomato (*Lycopersicon esculentum*) (Martin et al., (1993) Science 262:1432-1436). Tomato plants expressing the *Pto* gene are resistant to
- 25 infection by strains of *Pseudomonas syringae* pv. *tomato* that express the *avrPto* avirulence gene. The amino acid sequence inferred from the *Pto* gene DNA sequence displays strong similarity to serine-threonine protein kinases, implicating *Pto* in signal transduction. No similarity to
- 30 the tomato *Pto* locus or any known protein kinases was observed for *RPS2*, suggesting that *RPS2* is representative of a new class of plant disease resistance genes.
- The isolation of a race-specific resistance gene from *Zea mays* (corn) known as *Hm1* has been reported
- 35 (Johal and Briggs (1992) Science 258:985-987). *Hm1*

- 23 -

confers resistance against specific races of the fungal pathogen *Cochliobolus carbonum* by controlling degradation of a fungal toxin, a strategy that is mechanistically distinct from the avirulence-gene specific resistance of the *RPS2*-*avrRpt2* resistance mechanism.

The cloned *RPS2* gene of the invention can be used to facilitate the construction of plants that are resistant to specific pathogens and to overcome the inability to transfer disease resistance genes between species using classical breeding techniques (Keen et al., (1993), supra). There now follows a description of the cloning and characterization of an *Arabidopsis thaliana* *RPS2* genetic locus, the *RPS2* genomic DNA, and the *RPS2* cDNA. The *avrRpt2* gene and the *RPS2* gene, as well as mutants *rps2-101C*, *rps2-102C*, and *rps2-201C* (also designated *rps2-201*), are described in Dong, et al., (1991) Plant Cell 3:61-72; Yu, et al., (1993) supra; Kunkel et al., (1993) supra; Whalen et al., (1991), supra; and Innes et al., (1993), supra). A mutant designated *rps2-101N* has also been isolated. The identification and cloning of the *RPS2* gene is described below.

RPS2 Overcomes Sensitivity to Pathogens Carrying the *avrRpt2* Gene

To demonstrate the genetic relationship between an avirulence gene in the pathogen and a resistance gene in the host, it was necessary first to isolate an avirulence gene. By screening *Pseudomonas* strains that are known pathogens of crop plants related to *Arabidopsis*, highly virulent strains, *P. syringae* pv. *maculicola* (*Psm*) ES4326, *P. syringae* pv. *tomato* (*Pst*) DC3000, and an avirulent strain, *Pst* MM1065 were identified and analyzed as to their respective abilities to grow in wild type *Arabidopsis thaliana* plants (Dong et al., (1991) Plant

- 24 -

Cell, 3:61-72; Whalen et al., (1991) Plant Cell 3:49-59; MM1065 is designated JL1065 in Whalen et al.). *Psm* ES4326 or *Pst* DC3000 can multiply 10^4 fold in *Arabidopsis thaliana* leaves and cause water-soaked lesions that
5 appear over the course of two days. *Pst* MM1065 multiplies a maximum of 10 fold in *Arabidopsis thaliana* leaves and causes the appearance of a mildly chlorotic dry lesion after 48 hours. Thus, disease resistance is associated with severely inhibited growth of the
10 pathogen.

An avirulence gene (*avr*) of the *Pst* MM1065 strain was cloned using standard techniques as described in Dong et al. (1991), Plant Cell 3:61-72; Whalen et al., (1991) supra; and Innes et al., (1993), supra. The isolated
15 avirulence gene from this strain was designated *avrRpt2*. Normally, the virulent strain *Psm* ES4326 or *Pst* DC3000 causes the appearance of disease symptoms after 48 hours as described above. In contrast, *Psm* ES4326/*avrRpt2* or *Pst* DC3000/*avrRpt2* elicits the appearance of a visible
20 necrotic hypersensitivity response (HR) within 16 hours and multiplies 50 fold less than *Psm* ES4326 or *Pst* DC3000 in wild type *Arabidopsis thaliana* leaves (Dong et al., (1991), supra; and Whalen et al., (1991), supra). Thus, disease resistance in a wild type *Arabidopsis* plant
25 requires, in part, an avirulence gene in the pathogen or a signal generated by the avirulence gene.

The isolation of four *Arabidopsis thaliana* disease resistance mutants has been described using the cloned *avrRpt2* gene to search for the host gene required for
30 disease resistance to pathogens carrying the *avrRpt2* gene (Yu et al., (1993), supra; Kunkel et al., (1993), supra). The four *Arabidopsis thaliana* mutants failed to develop an HR when infiltrated with *Psm* ES4326/*avrRpt2* or *Pst* DC3000/*avrRpt2* as expected for plants having lost their
35 disease resistance capacity. In the case of one of these

- 25 -

mutants, approximately 3000 five to six week old M₂ ecotype Columbia (Col-0 plants) plants generated by ethyl methanesulfonic acid (EMS) mutagenesis were hand-inoculated with Psm ES4326/avrRpt2 and a single mutant, 5 *rps2-101C*, was identified (resistance to *Pseudomonas syringae*) (Yu et al., (1993), supra).

The second mutant was isolated using a procedure that specifically enriches for mutants unable to mount an HR (Yu et al., (1993), supra). When 10-day old 10 *Arabidopsis thaliana* seedlings growing on petri plates are infiltrated with *Pseudomonas syringae* pv. *phaseolicola* (Psp) NPS3121 versus Psp NPS3121/avrRpt2, about 90% of the plants infiltrated with Psp NPS3121 survive, whereas about 90%-95% of the plants infiltrated 15 with Psp NPS3121/avrRpt2 die. Apparently, vacuum infiltration of an entire small *Arabidopsis thaliana* seedling with Psp NPS3121/avrRpt2 elicits a systemic HR which usually kills the seedling. In contrast, seedlings infiltrated with Psp NPS3121 survive because Psp NPS3121 20 is a weak pathogen on *Arabidopsis thaliana*. The second disease resistance mutant was isolated by infiltrating 4000 EMS-mutagenized Columbia M₂ seedlings with Psp NPS3121/avrRpt2. Two hundred survivors were obtained. These were transplanted to soil and re-screened by hand 25 inoculation when the plants reached maturity. Of these 200 survivors, one plant failed to give an HR when hand-infiltrated with Psm ES4326/avrRpt2. This mutant was designated *rps2-102C* (Yu et al., (1993), supra).

A third mutant, *rps2-201C*, was isolated in a 30 screen of approximately 7500 M₂ plants derived from seed of *Arabidopsis thaliana* ecotype Col-0 that had been mutagenized with diepoxybutane (Kunkel et al., (1993), supra). Plants were inoculated by dipping entire leaf rosettes into a solution containing Pst DC3000/avrRpt2 35 bacteria and the surfactant Silwet L-77 (Whalen et al.,

- 26 -

(1991), supra), incubating plants in a controlled environment growth chamber for three to four days, and then visually observing disease symptom development. This screen revealed four mutant lines (carrying the
5 *rps2-201C*, *rps2-202C*, *rps2-203C*, and *rps2-204C* alleles), and plants homozygous for *rps2-201C* were a primary subject for further study (Kunkel et al., (1993), supra and the instant application).

Isolation of the fourth *rps2* mutant, *rps2-101N*,
10 has not yet been published. This fourth isolate is either a mutant or a susceptible *Arabidopsis* ecotype. Seeds of the *Arabidopsis* Nossen ecotype were gamma-irradiated and then sown densely in flats and allowed to germinate and grow through a nylon mesh. When the plants
15 were five to six weeks old, the flats were inverted, the plants were partially submerged in a tray containing a culture of *Psm* ES4326/*avrRpt2*, and the plants were vacuum infiltrated in a vacuum desiccator. Plants inoculated this way develop an HR within 24 hours. Using this
20 procedure, approximately 40,000 plants were screened and one susceptible plant was identified. Subsequent RFLP analysis of this plant suggested that it may not be a Nossen mutant but rather a different *Arabidopsis* ecotype that is susceptible to *Psm* ES4326/*avrRpt2*. This plant is
25 referred to as *rps2-101N*. The isolated mutants *rps2-101C*, *rps2-102C*, *rps2-201C*, and *rps2-101N* are referred to collectively as the "*rps2* mutants".

The *rps2* Mutants Fail to Specifically Respond to the Cloned Avirulence Gene, *avrRpt2*

30 The *RPS2* gene product is specifically required for resistance to pathogens carrying the avirulence gene, *avrRpt2*. A mutation in *Rps2* polypeptide that eliminates or reduces its function would be observable as the absence of a hypersensitive response upon infiltration of

- 27 -

the pathogen. The *rps2* mutants displayed disease symptoms or a null response when infiltrated with *Psm* ES4326/*avrRpt2*, *Pst* DC3000/*avrRpt2* or *Psp* NPS3121/*avrRpt2*, respectively. Specifically, no HR
5 response was elicited, indicating that the plants were susceptible and had lost resistance to the pathogen despite the presence of the *avrRpt2* gene in the pathogen.

Pathogen growth in *rps2* mutant plant leaves was similar in the presence and absence of the *avrRpt2* gene.
10 *Psm* ES4326 and *Psm* ES4326/*avrRpt2* growth in *rps2* mutants was compared and found to multiply equally well in the *rps2* mutants, at the same rate that *Psm* ES4326 multiplied in wild-type *Arabidopsis* leaves. Similar results were observed for *Pst* DC3000 and *Pst* DC3000/*avrRpt2* growth in
15 *rps2* mutants.

The *rps2* mutants displayed a HR when infiltrated with *Pseudomonas* pathogens carrying other *avr* genes, *Psm* ES4326/*avrB*, *Pst* DC3000/*avrB*, *Psm* ES4326/*avrRpm1*, *Pst* DC3000/*avrRpm1*. The ability to mount an HR to an *avr*
20 gene other than *avrRpt2* indicates that the *rps2* mutants isolated by selection with *avrRpt2* are specific to *avrRpt2*.

Mapping and Cloning of the RPS2 Gene

Genetic analysis of *rps2* mutants *rps2-101C*, *rps2-102C*, *rps-201C* and *rps-101N* showed that they all
25 corresponded to genes that segregated as expected for a single Mendelian locus and that all four were most likely allelic. The four *rps2* mutants were mapped to the bottom of chromosome IV using standard RFLP mapping procedures
30 including polymerase chain reaction (PCR)-based markers (Yu et al., (1993), supra; Kunkel et al., (1993), supra; and Mindrinos, M., unpublished). Segregation analysis showed that *rps2-101C* and *rps2-102C* are tightly linked to the PCR marker, PG11, while the RFLP marker M600 was used

- 28 -

to define the chromosome location of the *rps2-201C* mutation (Fig. 1A) (Yu et al., (1993), supra; Kunkel et al., (1993), supra). *RPS2* has subsequently been mapped to the centromeric side of PG11.

5 Heterozygous *RPS2/rps2* plants display a defense response that is intermediate between those displayed by the wild-type and homozygous *rps2/rps2* mutant plants (Yu, et al., (1993), supra; and Kunkel et al., (1993), supra). The heterozygous plants mounted an HR in response to *Psm*
10 ES4326/*avrRpt2* or *Pst* DC3000/*avrRpt2* infiltration; however, the HR appeared later than in wild type plants and required a higher minimum inoculum (Yu, et al., (1993), supra; and Kunkel et al., (1993), supra).

High Resolution Mapping of the *RPS2* Gene and *RPS2* cDNA

15 Isolation

To carry out map-based cloning of the *RPS2* gene, *rps2-101N/rps2-101N* was crossed with Landsberg erecta *RPS2/RPS2*. Plants of the F_1 generation were allowed to self pollinate (to "self") and 165 F_2 plants were selfed
20 to generate F_3 families. Standard RFLP mapping procedures showed that *rps2-101N* maps close to and on the centromeric side of the RFLP marker, PG11. To obtain a more detailed map position, *rps2-101N/rps2-101N* was crossed with a doubly marked Landsberg erecta strain
25 containing the recessive mutations, *cer2* and *ap2*. The genetic distance between *cer2* and *ap2* is approximately 15 cM, and the *rps2* locus is located within this interval. F_2 plants that displayed either a *CER2 ap2* or a *cer2 AP2* genotype were collected, selfed, and scored for *RPS2* by
30 inoculating at least 20 F_3 plants for each F_2 with *Psm* ES4326/*avrRpt2*. DNA was also prepared from a pool of approximately 20 F_3 plants for each F_2 line. The *CER2 ap2* and *cer2 AP2* recombinants were used to carry out a chromosome walk that is illustrated in Figure 1.

- 29 -

As shown in Figure 1, *RPS2* was mapped to a 28-35 kb region spanned by cosmid clones E4-4 and E4-6. This region contains at least six genes that produce detectable transcripts. There were no significant differences in the sizes of the transcripts or their level of expression in the *rps2* mutants as determined by RNA blot analysis. cDNA clones of each of these transcripts were isolated and five of these were sequenced. As is described below, one of these transcripts, cDNA-4, was shown to correspond to the *RPS2* locus. From this study, three independent cDNA clones (cDNA-4-4, cDNA-4-5, and cDNA-4-11) were obtained corresponding to *RPS2* from Columbia ecotype wild type plants. The apparent sizes of *RPS2* transcripts were 3.8 and 3.1 kb as determined by RNA blot analysis.

A fourth independent cDNA-4 clone (cDNA-4-2453) was obtained using map-based isolation of *RPS2* in a separate study. Yeast artificial chromosome (YAC) clones were identified that carry contiguous, overlapping inserts of *Arabidopsis thaliana* ecotype Col-0 genomic DNA from the M600 region spanning approximately 900 kb in the *RPS2* region. *Arabidopsis* YAC libraries were obtained from J. Ecker and E. Ward, supra and from E. Grill (Grill and Somerville (1991) Mol. Gen. Genet. 226:484-490). Cosmids designated "H" and "E" were derived from the YAC inserts and were used in the isolation of *RPS2* (Fig. 1).

The genetic and physical location of *RPS2* was more precisely defined using physically mapped RFLP, RAPD (random amplified polymorphic DNA) and CAPS (cleaved amplified polymorphic sequence) markers. Segregating populations from crosses between plants of genotype *RPS2/RPS2* (No-0 wild type) and *rps2-201/rps2-201* (Col-0 background) were used for genetic mapping. The *RPS2* locus was mapped using markers 17B7LE, PG11, M600 and other markers. For high-resolution genetic mapping, a

- 30 -

set of tightly linked RFLP markers was generated using insert end fragments from YAC and cosmid clones (Fig. 1) (Kunkel et al. (1993), supra; Konieczny and Ausubel (1993) Plant J. 4:403-410; and Chang et al. (1988) PNAS 5 USA 85:6856-6860). Cosmid clones E4-4 and E4-6 were then used to identify expressed transcripts (designated cDNA-4, -5, -6, -7, -8 of Fig 1F) from this region, including the cDNA-4-2453 clone.

RPS2 DNA Sequence Analysis

10 DNA sequence analysis of cDNA-4 from wild-type Col-0 plants and from mutants *rps2-101C*, *rps2-102C*, *rps2-201C* and *rps2-101N* showed that cDNA-4 corresponds to RPS2. DNA sequence analysis of *rps2-101C*, *rps2-102C* and *rps2-201C* revealed changes from the wild-type sequence as
15 shown in Table 1. The numbering system in Table 1 starts at the ATG start codon encoding the first methionine where A is nucleotide 1. DNA sequence analysis of cDNA-4 corresponding to mutant *rps2-102C* showed that it differed from the wild type sequence at amino acid residue 476.
20 Moreover, DNA sequence analysis of the cDNA corresponding to cDNA-4 from *rps2-101N* showed that it contained a 10 bp insertion at amino acid residue 581, a site within the leucine-rich repeat region which causes a shift in the RPS2 reading frame. Mutant *rps2-101C* contains a mutation
25 that leads to the formation of a chain termination codon. The DNA sequence of mutant allele *rps2-201C* revealed a mutation altering a single amino acid within a segment of the LRR region that also has similarity to the helix-loop-helix motif, further supporting the designation of
30 this locus as the RPS2 gene. The DNA and amino acid sequences are shown in Figure 2.

- 31 -

Table 1

Mutant	Wild type	position of mutation	Change
<i>rps2-101C</i>	703 TGA 705	704	TAA Stop Codon
5 <i>rps2-101N</i>	1741 GTG 1743	1741	GTGGAGTTGTATG Insertion
<i>rps2-102C</i>	1426 AGA 1428 arg	1427	AAA Amino acid 476 lys
10 <i>rps2-201C</i>	2002 ACC 2004 thr	2002	CCC Amino acid pro

DNA sequence analysis of cDNA-4 corresponding to *RPS2* from wild-type Col-0 plants revealed an open reading frame (between two stop codons) spanning 2,751 bp. There are 2,727 bp between the first methionine codon of this
 15 reading frame and the 3'-stop codon, which corresponds to a deduced 909 amino acid polypeptide (See open reading frame "a" of Fig. 2). The amino acid sequence has a relative molecular weight of 104,460 and a pI of 6.51.

As discussed below, *RPS2* belongs to a new class of
 20 disease resistance genes; the structure of the *Rps2* polypeptide does not resemble the protein structure of the product of the previously cloned and publicized avirulence gene-specific plant disease resistance gene, *Pto*, which has a putative protein kinase domain. From
 25 the above analysis of the deduced amino acid sequence, *RPS2* contains several distinct protein domains conserved in other proteins from both eukaryotes and prokaryotes. These domains include, but are not limited, to Leucine Rich Repeats (LRR) (Kobe and Deisenhofer, (1994) Nature
 30 366:751-756); nucleotide binding site, e.g. the kinase 1a motif (P-loop) (Saraste et al. (1990) Trends in Biological Sciences TIBS 15:430-434; Helix-Loop-Helix (Murre et al. (1989) Cell 56:777-783; and Leucine Zipper

- 32 -

(Rodrigues and Park (1993) Mol. Cell Biol. 13:6711-6722). The amino acid sequence of Rps2 contains a LRR motif (LRR motif from amino acid residue 505 to amino acid residue 867), which is present in many known proteins and which is thought to be involved in protein-protein interactions and may thus allow interaction with other proteins that are involved in plant disease resistance. The N-terminal portion of the Rps2 polypeptide LRR is, for example, related to the LRR of yeast (*Saccharomyces cerevisiae*) adenylate cyclase, CYR1. A region predicted to be a transmembrane spanning domain (Klein et al. (1985) Biochim., Biophys. Acta 815:468-476) is located from amino acid residue 350 to amino acid residue 365, N-terminal to the LRR. An ATP/GTP binding site motif (P-loop) is predicted to be located between amino acid residue 177 and amino acid residue 194, inclusive. The motifs are discussed in more detail below.

From the above analysis of the deduced amino acid sequence, the Rps2 polypeptide may have a membrane-receptor structure which consists of an N-terminal extracellular region and a C-terminal cytoplasmic region. Alternatively, the topology of the Rps2 may be the opposite: an N-terminal cytoplasmic region and a C-terminal extracellular region. LRR motifs are extracellular in many cases and the Rps2 LRR contains five potential N-glycosylation sites.

Identification of RPS2 by Functional Complementation

Complementation of *rps2-201* homozygotes with genomic DNA corresponding to *Arabidopsis thaliana* functionally confirmed that the genomic region encoding cDNA-4 carries RPS2 activity. Cosmids were constructed that contained overlapping contiguous sequences of wild type *Arabidopsis thaliana* DNA from the RPS2 region contained in YACs EW11D4, EW9C3, and YUP11F1 of Fig. 1

- 33 -

and Fig. 4. The cosmid vectors were constructed from pSLJ4541 (obtained from J. Jones, Sainsbury Institute, Norwich, England) which contains sequences that allow the inserted sequence to be integrated into the plant genome via *Agrobacterium*-mediated transformation (designated "binary cosmid"). "H" and "E" cosmids (Fig. 1) were used to identify clones carrying DNA from the *Arabidopsis thaliana* genomic *RPS2* region.

More than forty binary cosmids containing inserted *RPS2* region DNA were used to transform *rps2-201* homozygous mutants utilizing *Agrobacterium*-mediated transformation (Chang et al. ((1990) p. 28, Abstracts of the Fourth International Conference on *Arabidopsis* Research, Vienna, Austria). Transformants which remained susceptible (determined by methods including the observed absence of an HR following infection to *P. syringae* pv. *phaseolicola* strain 3121 carrying *avrRpt2* and Psp 3121 without *avrRpt2*) indicated that the inserted DNA did not contain functional *RPS2*. These cosmids conferred the "Sus." or susceptible phenotype indicated in Fig. 4. Transformants which had acquired *avrRpt2*-specific disease resistance (determined by methods including the display of a strong hypersensitive response (HR) when inoculated with Psp 3121 with *avrRpt2*, but not following inoculation with Psp 3121 without *avrRpt2*) suggested that the inserted DNA contained a functional *RPS2* gene capable of conferring the "Res." or resistant phenotype indicated in Fig. 4. Transformants obtained using the pD4 binary cosmid displayed a strong resistance phenotype as described above. The presence of the insert DNA in the transformants was confirmed by classical genetic analysis (the tight genetic linkage of the disease resistance phenotype and the kanamycin resistance phenotype conferred by the cotransformed selectable marker) and Southern analysis. These results indicated that *RPS2* is

- 34 -

encoded by a segment of the 18 kb *Arabidopsis thaliana* genomic region carried on cosmid pD4 (Fig. 4).

To further localize the *RPS2* locus and confirm its ability to confer a resistance phenotype on the *rps2-201* homozygous mutants, a set of six binary cosmids containing partially overlapping genomic DNA inserts were tested. The overlapping inserts pD2, pD4, pD14, pD15, pD27, and pD47 were chosen based on the location of the transcription corresponding to the five cDNA clones in the *RPS2* region (Fig. 4). These transformation experiments utilized a vacuum infiltration procedure (Bechtold et al. (1993) C.R. Acad. Sci. Paris 316:1194-1199) for *Agrobacterium*-mediated transformation. *Agrobacterium*-mediated transformations with cosmids pD2, pD14, pD15, pD39, and pD46 were performed using a root transformation/regeneration protocol (Valveekens et al. (1988), PNAS 85:5536-5540). The results of pathogen inoculation experiments assaying for *RPS2* activity in these transformants is indicated in Fig. 4.

These experiments were further confirmed using a modification of the vacuum filtration procedure. In particular, the procedure of Bechtold et al. (supra) was modified such that plants were grown in peat-based potting soil covered with a screen, primary inflorescences were removed, and plants with secondary inflorescences (approximately 3 to 15 cm in length) were inverted directly into infiltration medium, infiltrated, and then grown to seed harvest without removal from soil (detailed protocol available on the AAtDB computer database (43)). The presence of introduced sequences in the initial pD4 transformant was verified by DNA blot analysis with a pD4 vector and insert sequences (separately) as probes. The presence of the expected sequences in transformants obtained with the vacuum infiltration protocol was also confirmed by DNA blot

- 35 -

analysis. Root transformation experiments (19) were performed with an easily regenerable *rps2-201/rps2-201* x No-0 mapping population. Transformants were obtained for pD4 with in plant transformation, for pD2, 14, 16, 39, 5 and 49 with root transformation, and for pD2, 4, 14, 15, 27, and 47 with vacuum infiltration as modified.

Additional transformation experiments utilized binary cosmids carrying the complete coding region and more than 1 kb of upstream genomic sequence for only 10 cDNA-4 or cDNA-6. Using the vacuum infiltration transformation method, three independent transformants were obtained that carried the wild-type cDNA-6 genomic region in a *rps2-201c* homozygous background (pAD431 of Fig. 4). None of these plants displayed *avrRpt2*- 15 dependent disease resistance. Homozygous *rps2-201c* mutants were transformed with wild-type genomic cDNA-4 (p4104 and p4115, each carrying Col-0 genomic sequences corresponding to all of the cDNA-4 open reading frame, plus approximately 1.7 kb of 5' upstream sequence and 20 approximately 0.3 kb of 3' sequence downstream of the stop codon). These p4104 and p4115 transformants displayed a disease resistance phenotype similar to the wild-type *RPS2* homozygotes from which the *rps2* were derived. Additional mutants (*rps2-101N* and *rps2-101C* 25 homozygotes) also displayed *avrRpt2*-dependent resistance when transformed with the cDNA-4 genomic region.

RPS2 Sequences Allow Detection of Other Resistance Genes

DNA blot analysis of *Arabidopsis thaliana* genomic DNA using *RPS2* cDNA as the probe showed that *Arabidopsis* 30 contains several DNA sequences that hybridize to *RPS2* or a portion thereof, suggesting that there are several related genes in the *Arabidopsis* genome.

From the aforementioned description and the nucleic acid sequence shown in Fig. 2, it is possible to

- 36 -

isolate other plant disease resistance genes having about 50% or greater sequence identity to the *RPS2* gene. Detection and isolation can be carried out with an oligonucleotide probe containing the *RPS2* gene or a
5 portion thereof greater than 9 nucleic acids in length, and preferably greater than about 18 nucleic acids in length. Probes to sequences encoding specific structural features of the *Rps2* polypeptide are preferred as they provide a means of isolating disease resistance genes
10 having similar structural domains. Hybridization can be done using standard techniques such as are described in Ausubel et al., *Current Protocols in Molecular Biology*, John Wiley & Sons, (1989).

For example, high stringency conditions for
15 detecting the *RPS2* gene include hybridization at about 42°C, and about 50% formamide; a first wash at about 65°C, about 2X SSC, and 1% SDS; followed by a second wash at about 65°C and about 0.1% x SSC. Lower stringency conditions for detecting *RPS* genes having about 50%
20 sequence identity to the *RPS2* gene are detected by, for example, hybridization at about 42°C in the absence of formamide; a first wash at about 42°C, about 6X SSC, and about 1% SDS; and a second wash at about 50°C, about 6X SSC, and about 1% SDS. An approximately 350 nucleotide
25 DNA probe encoding the middle portion of the LRR region of *Rps2* was used as a probe in the above example. Under lower stringency conditions, a minimum of 5 DNA bands were detected in *Bam*HI digested *Arabidopsis thaliana* genomic DNA as sequences having sufficient sequence
30 identity to hybridize to DNA encoding the middle portion of the LRR motif of *Rps2*. Similar results were obtained using a probe containing a 300 nucleotide portion of the *RPS2* gene encoding the extreme N-terminus of *Rps2* outside of the LRR motif.

- 37 -

Isolation of other disease resistance genes is performed by PCR amplification techniques well known to those skilled in the art of molecular biology using oligonucleotide primers designed to amplify only
5 sequences flanked by the oligonucleotides in genes having sequence identity to *RPS2*. The primers are optionally designed to allow cloning of the amplified product into a suitable vector.

The RPS Disease-Resistance Gene Family

10 As discussed above, we have discovered that the *Arabidopsis RPS2* gene described herein is representative of a new class of plant resistance genes. Analysis of the derived amino acid sequence for *RPS2* revealed several regions of similarity with known polypeptide motifs (see,
15 e.g., Schneider et al., Genes Dev. 6:797 (1991)). Most prominent among these is a region of multiple, leucine-rich repeats (LRRs). The LRR motif has been implicated in protein-protein interactions and ligand binding in a diverse array of proteins (see, e.g., Kornfield et al.,
20 Annu. Rev. Biochem. 64:631 (1985); Alber, Curr. Opin. Gen. Dev. 2:205 (1992); Lupas et al., Science 252:1162 (1991); Saraste et al., Trend Biochem. Sci. 15:430 (1990)). In one example, LRRs form the hormone binding sites of mammalian gonadotropin hormone receptors (see,
25 e.g., Lupas et al., Science 252:1162 (1991)) and, in another example, a domain of yeast adenylate cyclase that interacts with the RAS2 protein (Kornfield et al., Annu. Rev. Biochem. 64:631 (1985)). In *RPS2*, the LRR domain spans amino acids 503-867 and contains fourteen repeat
30 units of length 22-26 amino acids. A portion of each repeat resembles the LRR consensus sequence (I/L/V)XXLXXLXX(I/L)XL. In Figure 7, the LRRs from *RPS2* are shown, as well as an *RPS2* consensus sequence. Within the *RPS2* LRR region, five (of six) sequences matching the

- 38 -

N-glycosylation consensus sequence [NX(S/T)] were observed (Figure 8, marked with a dot). In particular, N-glycosylation is predicted to occur at amino acids 158, 543, 666, 757, 778, 787. Interestingly, the single
5 nucleotide difference between functional *RPS2* and mutant allele *rps2-201* is within the LRR coding region, and this mutation disrupts one of the potential glycosylation sites.

Also observed in the deduced amino acid sequence
10 for *RPS2* is a second potential protein-protein interaction domain, a leucine zipper (see, e.g., von Heijne, J. Mol. Biol. 225:487 (1992)), at amino acids 30-57. This region contains four contiguous heptad repeats that match the leucine zipper consensus sequence
15 (I/R)XDLXXX. Leucine zippers facilitate the dimerization of transcription factors by formation of coiled-coil structures, but no sequences suggestive of an adjacent DNA binding domain (such as a strongly basic region or a potential zinc-finger) were detected in *RPS2*. Coiled-
20 coil regions also promote specific interactions between proteins that are not transcription factors (see, e.g., Ward et al., Plant Mol. Biol. 14:561 (1990); Ecker, Methods 1:186 (1990); Grill et al., Mol. Gen. Genet. 226:484 (1991)), and computer database similarity
25 searches with the region spanning amino acids 30-57 of *RPS2* revealed highest similarity to the coiled-coil regions of numerous myosin and paramyosin proteins.

A third *RPS2* motif was found at the sequence GPGGVGKT at deduced amino acids 182-189. This portion of
30 *RPS2* precisely matches the generalized consensus for the phosphate-binding loop (P-loop) of numerous ATP- and GTP-binding proteins (see, e.g., Saraste et al., supra). The postulated *RPS2* P-loop is similar to those found in RAS proteins and ATP synthase β -subunits (Saraste et al.,
35 supra), but surprisingly is most similar to the published

- 39 -

P-loop sequences for the *nifH* and *chvD* genes, respectively. The presence of this P-loop sequence strongly suggests nucleotide triphosphate binding as one aspect of RPS2 function. This domain is also referred to
5 as a kinase-1a motif (or a nucleotide binding site, or NBS). Other conserved NBSs are present in the RPS2 sequence; these NBSs include a kinase-2 motif at amino acids 258-262 and a kinase-3a motif at amino acids 330-335.

10 Finally, inspection of the RPS2 sequence reveals a fourth RPS2 motif, a potential membrane-spanning domain located at amino acids 340-360. Within this region, a conserved GLPLAL motif is found at amino acids 347-352. The presence of the membrane-spanning domain raises the
15 possibility that the RPS2 protein is membrane localized, with the N-terminal leucine zipper and P-loop domains residing together on the opposite side of the membrane from the LRR region. An orientation in which the C-terminal LRR domain is extracellular is suggested by the
20 fact that five of the six potential N-linked glycosylation sites occur C-terminal to the proposed membrane-spanning domain, as well as by the overall more positive charge of the N-terminal amino acid residues (see, e.g., Kornfield et al., supra; von Heijne, supra).
25 A number of proteins that contain LRRs are postulated or known to be membrane-spanning receptors in which the LRRs are displayed extracellularly as a ligand-binding domain (see, e.g., Lopez et al., Proc. Natl. Acad. Sci. 84:5615 (1987); Braun et al., EMBO J. 10:1885 (1991); Schneider
30 et al., supra).

The plant kingdom contains hundreds of resistance genes that are necessarily divergent since they control different resistance specificities. However, plant defense responses such as production of activated oxygen
35 species, PR-protein gene expression, and the

- 40 -

hypersensitive response are common to diverse plant-pathogen interactions. This implies that there are points of convergence in the defense signal transduction pathways downstream of initial pathogen recognition, and
5 also suggests that similar functional motifs may exist among diverse resistance gene products. Indeed, *RPS2* is dissimilar from previously described disease resistance genes such as *Hm1* or *Pto* (see, e.g., Johal et al., supra; Martin et al., supra), and thus represents a new class of
10 genes having disease resistance capabilities.

Isolation of Other Members of the RPS Disease-Resistance Gene Family Using Conserved Motif Probes and Primers

We have discovered that the *RPS2* motifs described above are conserved in other disease-resistance
15 genes, including, without limitation, the N protein, the L6 protein, and the Prf protein. As shown in Fig. 5(A and B), we have determined that the L6 polypeptide of flax, the N polypeptide of tobacco, and the Prf polypeptide of tomato each share unique regions of
20 similarity (including, but not limited to, the leucine-rich repeats, the membrane-spanning domain, the leucine zipper, and the P-loop and other NBS domains).

On the basis of this discovery, the isolation of virtually any member of the RPS gene family is made
25 possible using standard techniques. In particular, using all or a portion of the amino acid sequence of a conserved RPS motif (for example, the amino acid sequences defining any RPS P-loop, NBS, leucine-rich repeat, leucine zipper, or membrane-spanning region), one
30 may readily design RPS oligonucleotide probes, including RPS degenerate oligonucleotide probes (i.e., a mixture of all possible coding sequences for a given amino acid sequence). These oligonucleotides may be based upon the sequence of either strand of the DNA comprising the

- 41 -

motif. General methods for designing and preparing such probes are provided, for example, in Ausubel et al., supra and *Guide to Molecular Cloning Techniques*, 1987, S. L. Berger and A. R. Kimmel, eds., Academic Press, New York. These oligonucleotides are useful for RPS gene isolation, either through their use as probes capable of hybridizing to RPS complementary sequences or as primers for various polymerase chain reaction (PCR) cloning strategies.

Hybridization techniques and procedures are well known to those skilled in the art and are described, for example, in Ausubel et al., supra and *Guide to Molecular Cloning Techniques*, 1987, S. L. Berger and A. R. Kimmel, eds., Academic Press, New York. If desired, a combination of different oligonucleotide probes may be used for the screening of the recombinant DNA library. The oligonucleotides are labelled with ^{32}P using methods known in the art, and the detectably-labelled oligonucleotides are used to probe filter replicas from a recombinant plant DNA library. Recombinant DNA libraries may be prepared according to methods well known in the art, for example, as described in Ausubel et al., supra. Positive clones may, if desired, be rescreened with additional oligonucleotide probes based upon other RPS conserved regions. For example, an RPS clone identified based on hybridization with a P-loop-derived probe may be confirmed by re-screening with a leucine-rich repeat-derived oligonucleotide.

As discussed above, RPS oligonucleotides may also be used as primers in PCR cloning strategies. Such PCR methods are well known in the art and described, for example, in *PCR Technology*, H.A. Erlich, ed., Stockton Press, London, 1989; *PCR Protocols: A Guide to Methods and Applications*, M.A. Innis, D.H. Gelfand, J.J. Sninsky, and T.J. White, eds., Academic Press, Inc., New York,

- 42 -

1990; and Ausubel et al., supra. If desired, members of the RPS disease-resistance gene family may be isolated using the PCR "RACE" technique, or Rapid Amplification of cDNA Ends (see, e.g., Innis et al., supra). By this
 5 method, oligonucleotide primers based on an RPS conserved domain are oriented in the 3' and 5' directions and are used to generate overlapping PCR fragments. These overlapping 3'- and 5'-end RACE products are combined to produce an intact full-length cDNA. This method is
 10 described in Innis et al., supra; and Frohman et al., Proc. Natl. Acad. Sci. 85:8998, 1988.

Any number of probes and primers according to the invention may be designed based on the conserved RPS motifs described herein. Preferred motifs are boxed in
 15 the sequences shown in Fig. 5(A or B). In particular, oligonucleotides according to the invention may be based on the conserved P-loop domain, the amino acids of which are shown below:

MOTIF 1

20	L6	G MGGIGKTTTA [SEQ ID NO: 110]
	N	G MGGVGKTTIA [SEQ ID NO: 111]
	PrfP	G MPGLGKTTLA [SEQ ID NO: 112]
	RPS2	G PGGVGKTTLM [SEQ ID NO: 113]

From these sequences, appropriate oligonucleotides are
 25 designed and prepared using standard methods. Particular examples of RPS oligonucleotides based on the P-loop domain are as follows (N is A, C, T, or G).

Based on MOTIF 1:

30	5'	GGNATGGGNGGNNTNGGNAA(A or G)ACNAC 3' [SEQ ID NO: 158]
	5'	NCGNG(A/T)NGTNA(T/G)(G/A/T)A(T/A)NCGNA 3' [SEQ ID NO: 159]
	5'	GG(T or A)NT(T or G or C)GG(T or A)AA(G or A)AC(T or C or A)AC 3' [SEQ ID NO: 160]

- 43 -

5' GGNATGGGNGGNNTNGGNAA(A or G)ACNAC 3' [SEQ ID NO: 158]

5' N(G or A)(C or T)N(A or G)(A or G or T)NGTNGT(C or T)TTNCCNANNCCN(G or L)(G or C)N(G or A)(T or G)NCC 3' [SEQ ID NO: 161]

5' GGN(C or A)(T or C)N(G or C)(G or C)NGGNNTNGGNAA (A or G)ACNAC 3' [SEQ ID NO: 162]

Other conserved RPS motifs useful for oligonucleotide design are shown below. These motifs are also depicted in the sequence of Fig. 5(A or B).

MOTIF 2

L6	FKILVV LDDVD [SEQ ID NO: 114]
N	KKVLIV LDDID [SEQ ID NO: 115]
15 PrfP	KRFLIL IDDVW [SEQ ID NO: 116]
RPS2	KRFLLL LDDVW [SEQ ID NO: 117]

MOTIF 3

L6	SRFIIT SR [SEQ ID NO: 118]
N	SRIIIT TR [SEQ ID NO: 119]
20 PrfP	SRIILT TR [SEQ ID NO: 120]
RPS2	CKVMFT TR [SEQ ID NO: 121]

MOTIF 4

L6	GLPLTLK V [SEQ ID NO: 122]
N	GLPLALK V [SEQ ID NO: 123]
25 PrfP	GLPLSVV L [SEQ ID NO: 124]
RPS2	GLPLALI T [SEQ ID NO: 125]

MOTIF 5

L6	KISYDAL [SEQ ID NO: 126]
N	KISYDGL [SEQ ID NO: 127]
30 PrfP	GFSYKNL [SEQ ID NO: 128]
RPS2	KFSYDNL [SEQ ID NO: 129]

- 44 -

From the above motifs and the sequence motifs designated in Figure 5A and B, appropriate oligonucleotides are designed and prepared. Particular examples of such RPS oligonucleotides are as follows (N is A, T, C, or G).

- 5 Based on MOTIF 2:
- 5' T(T or C)GA(T or C)GA(T or C)(A or G)T(T or G
 or C)(T or G)(A or G)(T or G or C)(G or A)A
 3' [SEQ ID NO: 163]
- 10 5' T(T or C)CCA(G or C or A)A(T or C)(G or
 A)TC(A or G)TCNA 3' [SEQ ID NO: 164]
- 5' (C or G or A)(T or C)(C or A)NA(T or C)(G or
 A)TC(G or A)TCNA(G or A or T)NA(G or A or
 C)NANNA(G or A)NA 3' [SEQ ID NO: 165]
- 15 5' (T or A)(T or A)N(A or C)(A or G)(A or G)(T
 or G or A)TN(T or C)TNNTN(G or T or C)TN(A or
 T or C)TNGA(T or C)GA 3' [SEQ ID NO: 166]
- Based on MOTIF 3:
- 5' NCGNG(A or T)NGTNA(T or G)(G or A or T)A(T or
 A)NCGNGA 3' [SEQ ID NO: 167]
- 20 5' NCGNG(A or T)NGTNA(T or G)(G or A or T)A(T or
 A)NCGNGA 3' [SEQ ID NO: 167]
- 5' NC(G or T)N(G or C)(A or T)NGTNA(A or G or
 T)(A or G or T)AT(A or G or T)AATNG 3' [SEQ ID
 NO: 168].
- 25 Based on MOTIF 4:
- 5' NA(G or A)NGGNA(G or A)NCC 3' [SEQ ID NO: 169]
- 5' GG(T or A)(T or C)T(T or G or C)CC(T or A)(T
 or C)T(T or G or C)GC(T or C or A)(T or C)T
 3' [SEQ ID NO: 170]
- 30 5' A(A or G)(T or G or A)GC(G or C or A)A(G or
 A)(T or A)GG(G or C or A)A(G or A)(A or G or
 T or C)C C 3' [SEQ ID NO: 171]
- 5' NA(G or A)NGGNA(G or A)NCC 3' [SEQ ID NO:
 169]
- 35 5' N(A or G)NN(T or A)(T or C)NA(G or C or A)N(C
 or G)(A or T or C)NA(G or A)NGGNA(G or A)NCC
 3' [SEQ ID NO: 172]

- 45 -

5' GGN(T or C)TNCCN(T or C)TN(G or A or T)(C or G)N(T or G or C)T 3' [SEQ ID NO: 173]

Based on MOTIF 5:

5 A(A or G)(A or G)TT(A or G)TC(A or G)TA(G or A or T)(G or C)(T or A)(G or A)A(T or A)(C or T)TT 3' [SEQ ID NO: 174]

5' A(G or A)N(T or C)(T or C)NT(C or T)(A or G)TAN(G or C)(A or G)NANN(C or T)(C or T) 3' [SEQ ID NO: 175]

10 5' (G or A)(G or A)N(A or T)T(A or C or T)(T or A)(G or C)NTA(T or C)(G or A)AN(A or G)(A or C or G)N(T or C)T 3' [SEQ ID NO: 176]

Based on MOTIF 6:

15 5' GTNTT(T or C)(T or C)TN(T or A)(G or C)NTT(T or C)(A or C)G(A or G)GG 3' [SEQ ID NO: 177]

Based on MOTIF 7:

5' CCNAT(A or C or T)TT(T or C)TA(T or C)(G or A)(T or A)(G or T or C)GTNGA(T or C)CC 3' [SEQ ID NO: 178]

20 Based on MOTIF 8:

5' GTNGGNAT(A or C or T)GA(T or C)(G or A)(A or C)NCA 3' [SEQ ID NO: 179]

Based on MOTIF 9:

25 5' (G or A)AA(G or A)CANGC(A or G or T)AT(G or A)TCNA(G or A)(G or A)AA 3' [SEQ ID NO: 180]

5' TT(T or C)(T or C)TNGA(T or C)AT(A or C or T)GCNTG(T or C)TT 3' [SEQ ID NO: 181]

Based on MOTIF 10:

30 5' CCCAT(G or A)TC(T or C)(T or C)(T or G)NA(T or G or A)N(T or A)(G or A)(G or A)TC(A or G)TGCAT 3' [SEQ ID NO: 182]

5' ATGCA(T or C)GA(T or C)(T or C)(T or A)N(A or C or T)TN(A or C)(A or G)(A or G)GA(T or C)ATGGG 3' [SEQ ID NO: 183]

35 Based on MOTIF 11:

- 46 -

5' NA(G or A)N(G or C)(A or T)(T or C)T(T or C)NA(A or G)(C or T)TT 3' [SEQ ID NO: 184]

5' (A or T)(G or C)NAA(A or G)(T or C)TN(A or G)A(A or G)(A or T)(G or C)N(T or C)T 3' [SEQ ID NO: 185]

Based on MOTIF 12:

5' (A or G or T)(A or T)(A or T)(C or T)TCNA(G or A)N(G or C)(A or T)N(T or C)(G or T)NA(G or A)NCC 3' [SEQ ID NO: 186]

10 5' GGN(T or C)TN(A or C)(G or A)N(A or T)(G or L)N(T or C)TNGA 3' [SEQ ID NO: 187]

Once a clone encoding a candidate RPS family gene is identified, it is then determined whether such gene is capable of conferring disease-resistance to a plant host using the methods described herein or other methods well known in the art of molecular plant pathology.

A Biolistic Transient Expression Assay For Identification of Plant Resistance Genes

We have developed a functional transient expression system capable of providing a rapid and broadly applicable method for identifying and characterizing virtually any gene for its ability to confer disease-resistance to a plant cell. In brief, the assay system involves delivering by biolistic transformation a candidate plant disease-resistance gene to a plant tissue sample (e.g., a piece of tissue from a leaf) and then evaluating the expression of the gene within the tissue by appraising the presence or absence of a disease-resistance response (e.g., the hypersensitive response). This assay provides a method for identifying disease-resistance genes from a wide variety of plant species, including ones that are not amenable to genetic or transgenic studies.

- 47 -

The principle of the assay is depicted in the top portion of Figure 9. In general, plant cells carrying a mutation in the resistance gene of interest are utilized. Prior to biolistic transformation, the plant tissue is
5 infiltrated with a phytopathogenic bacterium carrying the corresponding avirulence gene. In addition, a gene to be assayed for its resistance gene activity is co-introduced by biolistics with a reporter gene. The expression of the cobombarded reporter gene serves as an indicator for
10 viability of the transformed cells. Both genes are expressed under the control of a strong and constitutive promoter. If the gene to be assayed does not complement the resistance gene function, the plant cells do not undergo a hypersensitive response (HR) and, therefore,
15 survive (Fig. 9, top panel, right). In this case, cells accumulate a large amount of the reporter gene product. If, on the other hand, a resistance gene is introduced, the plant cells recognize the signal from the avirulence-gene-carrying bacterium and undergo the HR because the
20 expressed resistance gene product complements the function (Fig. 9, top panel, left). In this case, the plant cells do not have enough time to accumulate a large amount of reporter gene product before their death. Given the transformation efficiency estimated by a proper
25 control (such as the uninfected half of the leaf), measuring the accumulation of reporter gene product can thus indicate whether the gene to be assayed complements the resistance gene function.

In one working example, we now demonstrate the
30 effectiveness of the transient expression assay, using the bacterial avirulence gene *avrRpt2* and the corresponding *Arabidopsis thaliana* resistance gene *RPS2* (Fig. 9, bottom panel). In brief, *rps2* mutant leaves, preinfected with *P. syringae* carrying *avrRpt2*, were co-
35 bombarded with two plasmids, one of which contained the

- 48 -

RPS2 gene and the other the *Escherichia coli uidA* gene encoding β -glucuronidase (GUS; Jefferson et al., 1986, supra). Both the *RPS2* and *uidA* genes are located downstream of the strong constitutive 35S promoter from cauliflower mosaic virus (Odell et al., infra). If the 35S-*RPS2* construct complements the *rps2* mutation, the transformed cells rapidly undergo programmed cell death in response to the *P. syringae* carrying *avrRpt2*, and relatively little GUS activity accumulates. If the *rps2* mutation is not complemented, cell death does not occur and high levels of GUS activity accumulate. These differences in GUS activity are detected histochemically. Because the cDNA library used to identify *RPS2* was constructed in the expression vector pKEx4tr, the 35S-*RPS2* cDNA construct in pKEx4tr could be used directly in the transient assay. As shown in Fig. 11, pKEx4tr is a cDNA expression vector designed for the unidirectional insertion of cDNA inserts. Inserted cDNA is expressed under the control of the 35S cauliflower mosaic virus promoter.

Our results are shown in Fig. 9, lower panel. In this experiment, we infected one side of a leaf of an *rps2* mutant plant with *P. syringae* pv. *phaselolicola* 3121 carrying *avrRpt2* (*Psp* 3121/*avrRpt2*). *Psp* 3121 is a weak pathogen of *A. thaliana* and *Psp* 3121/*avrRpt2* can elicit an HR in a plant carrying the resistance gene *RPS2* (e.g., a wild type plant). Leaves of 5-week-old *Arabidopsis* plants were infiltrated with an appropriate bacterial suspension at a dose of 2×10^8 /ml by hand infiltration as described (Dong et al., supra). After an incubation period (typically 2-4 hours), the leaves were bombarded using a Bio-Rad PDS-1000/He apparatus (1100 psi) after 2-4 hr of infection. Gold particles were prepared according to the instructions of the manufacturer. For each bombardment, 1.4 μ g of pKEx4tr-G, 0.1 μ g of a

- 49 -

plasmid to be tested, and 0.5 mg of 1 μ m gold particles were used. After the bombardment, the leaves were incubated in a humidity chamber at 22°C for 1 day and then subjected to a histochemical GUS staining using 5-bromo-4-chloro-3-indiyl glucuronidase (X-Gluc) at 37°C for 12 hr (Jefferson, 1987, supra). This staining method with X-gluc stains cells expressing GUS enzyme with a blue color. The uninfected side of the leaf serves as a control for transformation efficiency of the leaf because in a single leaf, transformation efficiency (i.e., density of transformed cells) is similar on both sides of the leaf. If transformed cells on the infected side are rapidly killed, staining of the cells on the infected side is weaker than staining on the uninfected side.

When the resistance gene *RPS2* was co-introduced, the transformed cells on the infected side of the leaf showed much weaker staining than ones on the uninfected side (Fig. 10). In contrast, when an unrelated gene was co-introduced, the transformed cells on the infected side showed similar staining intensity to ones on the uninfected side (Fig. 10).

Thus, as summarized in the Table 2, *35S-RPS4* (cDNA 4), but not cDNA-5 or cDNA-6, complemented the HR phenotype of *rps2-101C*. (See Figure 1)

25

Table 2

<u>Gene Tested</u>	<u>Response (Decreased GUS Activity)^a</u>
ΔGUS (35S-uidA containing internal uidA deletion)	-
cDNA-5 (35S-AB11)	
cDNA-4 (35S-RPS2)	+
cDNA-6 (35S-CK1)	

30

- 50 -

^aWhen decreased GUS activity was observed on the infiltrated side of the leaf, the response was scored as plus (Fig. 10).

Both *RPS2* cDNA-4 clones 4 and 11, corresponding to the
5 two *RPS2* different transcript sizes, complemented the
rps2 mutant phenotype, indicating that both transcripts
encode a functional product. Moreover, *35S-RPS2* also
complemented mutants *rps2-102C*, *rps2-101N*, and *rps2-201C*,
further confirming that the *rps2-101C*, *rps2-102C*, *rps2-*
10 *201C* and *rps2-101N* mutations are all allelic. In short,
the cloned *RPS2* gene complemented the *rps2* mutation in
this transient expression assay, and complementation by
RPS2 was observed in all four available *rps2* mutant
stains.

15 Next we used the transient assay system to test
the specificity of the cloned *RPS2* gene for an *avrRpt2*-
generated signal (i.e., the "gene-for-gene" specificity
of a *P. syringae* avirulence gene and a corresponding *A.*
thaliana resistance gene (*avrRpm1* and *RPM1*,
20 respectively)). This experiment involved the use of an
rps2-101 rpm1 double mutant that cannot mount an HR when
challenged with *P. syringae* carrying *avrRpt2* or the
unrelated avirulence gene *avrRpm1* (Debener et al., Plant
Journal 1:289-302, 1991). As summarized in Table 3,
25 complementation of the *rps2* mutant phenotype by *35S-RPS2*
was only observed in the presence of a signal generated
by *avrRpt2*, indicating that *RPS2* does not simply
sensitize the plant resistance response in a nonspecific
manner.

- 51 -

Table 3

<u>avr Gene</u>	<u>Construct Cobombarded with 35S-uidA</u>	<u>Response^a</u>
None (vector only)	Δ GUS ^b	-
5 <i>avrRpt2</i>	Δ GUS	-
<i>avrRpm1</i>	Δ GUS	-
None (vector only)	35S-RPS2	-
<i>avrRpt2</i>	35S-RPS2	+
<i>avrRpm1</i>	35S-RPS2	-

10 ^aWhen decreased GUS activity was observed on the infiltrated side of the leaf, the response was scored as plus. (Figure 10, panel B)

^b Δ GUS is 35S-uidA containing an internal deletion in the uidA gene.

15 Also as shown in Table 3, the RPS2 gene complemented the mutant phenotype when leaves were infected with *Psp* 3121/*avrRpt2* but not with *Psp* 3121/*avrRpm1*. Therefore, the RPS2 gene complemented only the *rps2* mutation; it did not the *rpm1* mutation.

20 We have also discovered that overexpression of an *rps* gene family member, e.g., *rps2* but not other genes, in the transient assay leads to apparent cell death, obviating the need to know the corresponding avirulence gene for a putative resistance gene that has been cloned.

25 Using this assay, any plant disease-resistance gene may be identified from a cDNA expression library. In one particular example, a cDNA library is constructed in an expression vector and then introduced as described herein into a plant cultivar or its corresponding mutant
30 plant lacking the resistance gene of interest.

 Preferably, the cDNA library is divided into small pools, and each pool co-introduced with a reporter gene. If a pool contains a resistance gene clone (i.e., the pool

- 52 -

"complements" the resistance gene function), the positive pool is divided into smaller pools and the same procedure is repeated until identification of a single positive clone is ultimately achieved. This approach facilitates the cloning of any resistance gene of interest without genetic crosses or the creation of transgenics.

We now describe the cloning of another member of the RPS gene family, the Prf gene of tomato.

The initial step for the cloning of the Prf gene came from classical genetic analysis which showed that Prf was tightly linked to the tomato Pto gene (Salmeron et al., The Plant Cell 6:511-520, 1994). This prompted construction of a cosmid contig of 200 kb in length which encompassed the Pto locus. DNA probes from this contig were used to screen a tomato cDNA library constructed using tomato leaf tissue that had been infected with Pst expressing the avrPto avirulence gene as source material. Two classes of cDNAs were identified based on cross-hybridization of clones to each other. While one class corresponded to members of the Pto gene family, the other class displayed no hybridization to Pto family members. Taking the assumption (based on the aforementioned genetic analysis) that Prf might reside extremely close to the Pto gene, cDNAs from the second class were analyzed further as candidate Prf clones. These clones were hybridized to filters containing DNAs from six independent prf mutant lines that had been isolated by diepoxybutane or fast neutron treatment. In one of the fast neutron mutants, the cDNA probe revealed a 1.1 kb deletion in the genomic DNA, suggesting that the cDNA clone might in fact represent Prf. Wild-type DNA corresponding to the deletion was cloned from Prf/Prf tomato. A 5 kb region was sequenced and found to potentially encode a protein containing P-loop and leucine-rich repeat motifs, supporting the hypothesis

- 53 -

that this DNA encoded Prf. The corresponding DNA was cloned and sequenced from the fast neutron mutant plant. Sequencing this DNA confirmed the mutation to be a simple 1.1 kb deletion excising DNA between the potential P-loop and leucine-rich repeat coding regions. The gene is expressed based on RT-PCR analysis which has shown that an mRNA is transcribed from this region. The identity of the cloned DNA as the Prf gene is based on both the existence of the deletion mutation and the predicted protein sequence, which reveals patches of strong similarity to other cloned disease resistance gene products throughout the amino-terminal half (as described herein). A partial sequence of the Prf gene is shown in Figure 12.

15 RPS Expression in Transgenic Plant Cells and Plants

The expression of the RPS2 genes in plants susceptible to pathogens carrying *avrRpt2* is achieved by introducing into a plant a DNA sequence containing the RPS2 gene for expression of the Rps2 polypeptide. A number of vectors suitable for stable transfection of plant cells or for the establishment of transgenic plants are available to the public; such vectors are described in, e.g., Pouwels et al., *Cloning Vectors: A Laboratory Manual*, 1985, Supp. 1987); Weissbach and Weissbach, *Methods for Plant Molecular Biology*, Academic Press, 1989; and Gelvin et al., *Plant Molecular Biology Manual*, Kluwer Academic Publishers, 1990. Typically, plant expression vectors include (1) one or more cloned plant genes under the transcriptional control of 5' and 3' regulatory sequences and (2) a dominant selectable marker. Such plant expression vectors may also contain, if desired, a promoter regulatory region (e.g., a regulatory region controlling inducible or constitutive, environmentally- or developmentally-regulated, or cell-

- 54 -

or tissue-specific expression), a transcription initiation start site, a ribosome binding site, an RNA processing signal, a transcription termination site, and/or a polyadenylation signal.

5 An example of a useful plant promoter which could be used to express a plant resistance gene according to the invention is a caulimovirus promoter, e.g., the cauliflower mosaic virus (CaMV) 35S promoter. These promoters confer high levels of expression in most plant
10 tissues, and the activity of these promoters is not dependent on virtually encoded proteins. CaMV is a source for both the 35S and 19S promoters. In most tissues of transgenic plants, the CaMV 35S promoter is a strong promoter (see, e.g., Odel et al., Nature 313:810,
15 (1985)). The CaMV promoter is also highly active in monocots (see, e.g., Dekeyser et al., Plant Cell 2:591, (1990); Terada and Shimamoto, Mol. Gen. Genet. 220:389, (1990)).

Other useful plant promoters include, without
20 limitation, the nonpaline synthase promoter (An et al., Plant Physiol. 88:547, (1988)) and the octopine synthase promoter (Fromm et al., Plant Cell 1:977, (1989)).

For certain applications, it may be desirable to produce the *RPS2* gene product or the *avrRpt2* gene product
25 in an appropriate tissue, at an appropriate level, or at an appropriate developmental time. Thus, there are a variety of gene promoters, each with its own distinct characteristics embodied in its regulatory sequences, shown to be regulated in response to the environment,
30 hormones, and/or developmental cues. These include gene promoters that are responsible for (1) heat-regulated gene expression (see, e.g., Callis et al., Plant Physiol. 88: 965, (1988)), (2) light-regulated gene expression (e.g., the pea *rbcS-3A* described by Kuhlemeier et al.,
35 Plant Cell 1: 471, (1989); the maize *rbcS* promoter

- 55 -

described by Schaffner and Sheen, Plant Cell 3: 997, (1991); or the chlorophyll a/b-binding protein gene found in pea described by Simpson et al., EMBO J. 4: 2723, (1985)), (3) hormone-regulated gene expression (e.g., the
5 abscisic acid responsive sequences from the *Em* gene of wheat described Marcotte et al., Plant Cell 1:969, (1989)), (4) wound-induced gene expression (e.g., of *wunI* described by Siebertz et al., Plant Cell 1: 961, (1989)), or (5) organ-specific gene expression (e.g., of the
10 tuber-specific storage protein gene described by Roshal et al., EMBO J. 6:1155, (1987); the 23-kDa zein gene from maize described by Schernthaner et al., EMBO J. 7: 1249, (1988); or the French bean β -phaseolin gene described by Bustos et al., Plant Cell 1:839, (1989)).

15 Plant expression vectors may also optionally include RNA processing signals, e.g, introns, which have been shown to be important for efficient RNA synthesis and accumulation (Callis et al., Genes and Dev. 1: 1183, (1987)). The location of the RNA splice sequences can
20 influence the level of transgene expression in plants. In view of this fact, an intron may be positioned upstream or downstream of an Rps2 polypeptide-encoding sequence in the transgene to modulate levels of gene expression.

25 In addition to the aforementioned 5' regulatory control sequences, the expression vectors may also include regulatory control regions which are generally present in the 3' regions of plant genes (Thornburg et al., Proc. Natl Acad. Sci USA 84: 744, (1987); An et
30 al., Plant Cell 1: 115, (1989)). For example, the 3' terminator region may be included in the expression vector to increase stability of the mRNA. One such terminator region may be derived from the PI-II terminator region of potato. In addition, other commonly

- 56 -

used terminators are derived from the octopine or nopaline synthase signals.

The plant expression vector also typically contains a dominant selectable marker gene used to
5 identify the cells that have become transformed. Useful selectable marker genes for plant systems include genes encoding antibiotic resistance genes, for example, those encoding resistance to hygromycin, kanamycin, bleomycin, G418, streptomycin or spectinomycin. Genes required for
10 photosynthesis may also be used as selectable markers in photosynthetic-deficient strains. Finally, genes encoding herbicide resistance may be used as selectable markers; useful herbicide resistance genes include the *bar* gene encoding the enzyme phosphinothricin
15 acetyltransferase, which confers resistance to the broad spectrum herbicide Basta® (Hoechst AG, Frankfurt, Germany).

Efficient use of selectable markers is facilitated by a determination of the susceptibility of a plant cell
20 to a particular selectable agent and a determination of the concentration of this agent which effectively kills most, if not all, of the transformed cells. Some useful concentrations of antibiotics for tobacco transformation include, e.g., 75-100 µg/ml (kanamycin), 20-50 µg/ml
25 (hygromycin), or 5-10 µg/ml (bleomycin). A useful strategy for selection of transformants for herbicide resistance is described, e.g., in Vasil I.K., *Cell Culture and Somatic Cell Genetics of Plants*, Vol I, II, III Laboratory Procedures and Their Applications Academic
30 Press, New York, 1984.

It should be readily apparent to one skilled in the field of plant molecular biology that the level of gene expression is dependent not only on the combination of promoters, RNA processing signals and terminator

- 57 -

elements, but also on how these elements are used to increase the levels of gene expression.

The above exemplary techniques may be used for the expression of any gene in the RPS family.

5 Plant Transformation

Upon construction of the plant expression vector, several standard methods are known for introduction of the recombinant genetic material into the host plant for the generation of a transgenic plant. These methods

10 include (1) *Agrobacterium*-mediated transformation (*A. tumefaciens* or *A. rhizogenes*) (see, e.g., Lichtenstein and Fuller In: *Genetic Engineering*, vol 6, PWJ Rigby, ed, London, Academic Press, 1987; and Lichtenstein, C.P., and Draper, J., In: *DNA Cloning*, Vol II, D.M. Glover, ed,

15 Oxford, IRI Press, 1985), (2) the particle delivery system (see, e.g., Gordon-Kamm et al., *Plant Cell* 2:603, (1990); or BioRad Technical Bulletin 1687, *supra*), (3) microinjection protocols (see, e.g., Green et al., *Plant Tissue and Cell Culture*, Academic Press, New York, 1987),

20 (4) polyethylene glycol (PEG) procedures (see, e.g., Draper et al., *Plant Cell Physiol* 23:451, (1982); or e.g., Zhang and Wu, *Theor. Appl. Genet.* 76:835, (1988)), (5) liposome-mediated DNA uptake (see, e.g., Freeman et al., *Plant Cell Physiol* 25: 1353, (1984)), (6)

25 electroporation protocols (see, e.g., Gelvin et al *supra*; Dekeyser et al. *supra*; or Fromm et al *Nature* 319: 791, (1986)), and (7) the vortexing method (see, e.g., Kindle, K., *Proc. Natl. Acad. Sci., USA* 87:1228, (1990)).

The following is an example outlining an

30 *Agrobacterium*-mediated plant transformation. The general process for manipulating genes to be transferred into the genome of plant cells is carried out in two phases. First, all the cloning and DNA modification steps are done in *E. coli*, and the plasmid containing the gene

- 58 -

construct of interest is transferred by conjugation into *Agrobacterium*. Second, the resulting *Agrobacterium* strain is used to transform plant cells. Thus, for the generalized plant expression vector, the plasmid contains
5 an origin of replication that allows it to replicate in *Agrobacterium* and a high copy number origin of replication functional in *E. coli*. This permits facile production and testing of transgenes in *E. coli* prior to transfer to *Agrobacterium* for subsequent introduction
10 into plants. Resistance genes can be carried on the vector, one for selection in bacteria, e.g., streptomycin, and the other that will express in plants, e.g., a gene encoding for kanamycin resistance or an herbicide resistance gene. Also present are restriction
15 endonuclease sites for the addition of one or more transgenes operably linked to appropriate regulatory sequences and directional T-DNA border sequences which, when recognized by the transfer functions of *Agrobacterium*, delimit the region that will be
20 transferred to the plant.

In another example, plant cells may be transformed by shooting into the cell tungsten microprojectiles on which cloned DNA is precipitated. In the Biolistic Apparatus (Bio-Rad, Hercules, CA) used for the shooting,
25 a gunpowder charge (22 caliber Power Piston Tool Charge) or an air-driven blast drives a plastic macroprojectile through a gun barrel. An aliquot of a suspension of tungsten particles on which DNA has been precipitated is placed on the front of the plastic macroprojectile. The
30 latter is fired at an acrylic stopping plate that has a hole through it that is too small for the macroprojectile to go through. As a result, the plastic macroprojectile smashes against the stopping plate and the tungsten microprojectiles continue toward their target through the
35 hole in the plate. For the instant invention the target

- 59 -

can be any plant cell, tissue, seed, or embryo. The DNA introduced into the cell on the microprojectiles becomes integrated into either the nucleus or the chloroplast.

Transfer and expression of transgenes in plant
5 cells is now routine practice to those skilled in the art. It has become a major tool to carry out gene expression studies and to attempt to obtain improved plant varieties of agricultural or commercial interest.

10 Transgenic Plant Regeneration

Plant cells transformed with a plant expression vector can be regenerated, e.g., from single cells, callus tissue or leaf discs according to standard plant tissue culture techniques. It is well known in the art
15 that various cells, tissues and organs from almost any plant can be successfully cultured to regenerate an entire plant; such techniques are described, e.g., in Vasil supra; Green et al., supra; Weissbach and Weissbach, supra; and Gelvin et al., supra.

20 In one possible example, a vector carrying a selectable marker gene (e.g., kanamycin resistance), a cloned *RPS2* gene under the control of its own promoter and terminator or, if desired, under the control of exogenous regulatory sequences such as the 35S CaMV
25 promoter and the nopaline synthase terminator is transformed into *Agrobacterium*. Transformation of leaf tissue with vector-containing *Agrobacterium* is carried out as described by Horsch et al. (Science 227: 1229, (1985)). Putative transformants are selected after a few
30 weeks (e.g., 3 to 5 weeks) on plant tissue culture media containing kanamycin (e.g. 100 µg/ml). Kanamycin-resistant shoots are then placed on plant tissue culture media without hormones for root initiation. Kanamycin-resistant plants are then selected for greenhouse growth.

- 60 -

If desired, seeds from self-fertilized transgenic plants can then be sowed in a soil-less media and grown in a greenhouse. Kanamycin-resistant progeny are selected by sowing surfaced sterilized seeds on hormone-free
5 kanamycin-containing media. Analysis for the integration of the transgene is accomplished by standard techniques (see, e.g., Ausubel et al. supra; Gelvin et al. supra).

Transgenic plants expressing the selectable marker are then screened for transmission of the transgene DNA
10 by standard immunoblot and DNA and RNA detection techniques. Each positive transgenic plant and its transgenic progeny are unique in comparison to other transgenic plants established with the same transgene. Integration of the transgene DNA into the plant genomic
15 DNA is in most cases random and the site of integration can profoundly effect the levels, and the tissue and developmental patterns of transgene expression. Consequently, a number of transgenic lines are usually screened for each transgene to identify and select plants
20 with the most appropriate expression profiles.

Transgenic lines are evaluated for levels of transgene expression. Expression at the RNA level is determined initially to identify and quantitate expression-positive plants. Standard techniques for RNA
25 analysis are employed and include PCR amplification assays using oligonucleotide primers designed to amplify only transgene RNA templates and solution hybridization assays using transgene-specific probes (see, e.g., Ausubel et al., supra). The RNA-positive plants are then
30 analyzed for protein expression by Western immunoblot analysis using Rps2 polypeptide-specific antibodies (see, e.g., Ausubel et al., supra). In addition, *in situ* hybridization and immunocytochemistry according to standard protocols can be done using transgene-specific

- 61 -

nucleotide probes and antibodies, respectively, to localize sites of expression within transgenic tissue.

Once the Rps2 polypeptide has been expressed in any cell or in a transgenic plant (e.g., as described above), it can be isolated using any standard technique, e.g., affinity chromatography. In one example, an anti-Rps2 antibody (e.g., produced as described in Ausubel et al., supra, or by any standard technique) may be attached to a column and used to isolate the polypeptide. Lysis and fractionation of Rps2-producing cells prior to affinity chromatography may be performed by standard methods (see, e.g., Ausubel et al., supra). Once isolated, the recombinant polypeptide can, if desired, be further purified, e.g., by high performance liquid chromatography (see, e.g., Fisher, *Laboratory Techniques In Biochemistry And Molecular Biology*, Work and Burdon, eds., Elsevier, 1980).

These general techniques of polypeptide expression and purification can also be used to produce and isolate useful Rps2 fragments or analogs.

Antibody Production

Using a polypeptide described above (e.g., the recombinant protein or a chemically synthesized RPS peptide based on its deduced amino acid sequence), polyclonal antibodies which bind specifically to an RPS polypeptide may be produced by standard techniques (see, e.g., Ausubel et al., supra) and isolated, e.g., following peptide antigen affinity chromatography. Monoclonal antibodies can also be prepared using standard hybridoma technology (see, e.g., Kohler et al., *Nature* 256: 495, 1975; Kohler et al., *Eur. J. Immunol.* 6: 292, 1976; Hammerling et al., in *Monoclonal Antibodies and T Cell Hybridomas*, Elsevier, N.Y., 1981; and Ausubel et al., supra).

- 62 -

Once produced, polyclonal or monoclonal antibodies are tested for specific RSP polypeptide recognition by Western blot or immunoprecipitation analysis (by methods described in Ausubel et al., supra). Antibodies which specifically recognize a RPS polypeptide are considered to be useful in the invention; such antibodies may be used, e.g., for screening recombinant expression libraries as described in Ausubel et al., supra. Exemplary peptides (derived from Rps2) for antibody production include:

LKFSYDNLESDDL [SEQ ID NO: 188]

GVYGPGGVGKTTLMQS [SEQ ID NO: 189]

GGLPLALITLGGAM [SEQ ID NO: 190]

Use

Introduction of RPS2 into a transformed plant cell provides for resistance to bacterial pathogens carrying the *avrRpt2* avirulence gene. For example, transgenic plants of the instant invention expressing RPS2 might be used to alter, simply and inexpensively, the disease resistance of plants normally susceptible to plant pathogens carrying the avirulence gene, *avrRpt2*.

The invention also provides for broad-spectrum pathogen resistance by mimicking the natural mechanism of host resistance. First, the RPS2 transgene is expressed in plant cells at a sufficiently high level to initiate the plant defense response constitutively in the absence of signals from the pathogen. The level of expression associated with plant defense response initiation is determined by measuring the levels of defense response gene expression as described in Dong et al., supra.

Second, the RPS2 transgene is expressed by a controllable promoter such as a tissue-specific promoter, cell-type specific promoter or by a promoter that is induced by an external signal or agent thus limiting the temporal and

- 63 -

tissue expression of a defense response. Finally, the *RPS2* gene product is co-expressed with the *avrRpt2* gene product. The *RPS2* gene is expressed by its natural promoter, by a constitutively expressed promoter such as
5 the CaMV 35S promoter, by a tissue-specific or cell-type specific promoter, or by a promoter that is activated by an external signal or agent. Co-expression of *RPS2* and *avrRpt2* will mimic the production of gene products associated with the initiation of the plant defense
10 response and provide resistance to pathogens in the absence of specific resistance gene-avirulence gene corresponding pairs in the host plant and pathogen.

The invention also provides for expression in plant cells of a nucleic acid having the sequence of Fig.
15 2 or the expression of a degenerate variant thereof encoding the amino acid sequence of open reading frame "a" of Fig. 2.

The invention further provides for the isolation of nucleic acid sequences having about 50% or greater
20 sequence identity to *RPS2* by using the *RPS2* sequence of Fig. 2 or a portion thereof greater than 9 nucleic acids in length, and preferably greater than about 18 nucleic acids in length as a probe. Appropriate reduced hybridization stringency conditions are utilized to
25 isolate DNA sequences having about 50% or greater sequence identity to the *RPS2* sequence of Fig. 2.

Also provided by the invention are short conserved regions characteristic of *RPS* disease resistance genes. These conserved regions provide oligonucleotide sequences
30 useful for the production of hybridization probes and PCR primers for the isolation of other plant disease-resistance genes.

Both the *RPS2* gene and related *RPS* family genes provide disease resistance to plants, especially crop
35 plants, most especially important crop plants such as

- 64 -

tomato, pepper, maize, wheat, rice and legumes such as soybean and bean, or any plant which is susceptible to pathogens carrying an avirulence gene, e.g., the *avrRpt2* avirulence gene. Such pathogens include, but are not
5 limited to, *Pseudomonas syringae* strains.

The invention also includes any biologically active fragment or analog of an Rps2 polypeptide. By "biologically active" is meant possessing any in vivo activity which is characteristic of the Rps2 polypeptide
10 shown in Fig. 2. A useful Rps2 fragment or Rps2 analog is one which exhibits a biological activity in any biological assay for disease resistance gene product activity, for example, those assays described by Dong et al. (1991), supra; Yu et al. (1993) supra; Kunkel et al.
15 (1993) supra; and Whalen et al. (1991). In particular, a biologically active Rps2 polypeptide fragment or analog is capable of providing substantial resistance to plant pathogens carrying the *avrRpt2* avirulence gene. By substantial resistance is meant at least partial
20 reduction in susceptibility to plant pathogens carrying the *avrRpt2* gene.

Preferred analogs include Rps2 polypeptides (or biologically active fragments thereof) whose sequences differ from the wild-type sequence only by conservative
25 amino acid substitutions, for example, substitution of one amino acid for another with similar characteristics (e.g., valine for glycine, arginine for lysine, etc.) or by one or more non-conservative amino acid substitutions, deletions, or insertions which do not abolish the
30 polypeptide's biological activity.

Analogous can differ from naturally occurring Rps2 polypeptide in amino acid sequence or can be modified in ways that do not involve sequence, or both. Analogous of the invention will generally exhibit at least 70%,
35 preferably 80%, more preferably 90%, and most preferably

- 65 -

95% or even 99%, homology with a segment of 20 amino acid residues, preferably 40 amino acid residues, or more preferably the entire sequence of a naturally occurring Rps2 polypeptide sequence.

5 Alterations in primary sequence include genetic variants, both natural and induced. Also included are analogs that include residues other than naturally occurring L-amino acids, e.g., D-amino acids or non-naturally occurring or synthetic amino acids, e.g., β or
10 γ amino acids. Also included in the invention are Rps2 polypeptides modified by in vivo chemical derivatization of polypeptides, including acetylation, methylation, phosphorylation, carboxylation, or glycosylation.

 In addition to substantially full-length
15 polypeptides, the invention also includes biologically active fragments of the polypeptides. As used herein, the term "fragment", as applied to a polypeptide, will ordinarily be at least 20 residues, more typically at least 40 residues, and preferably at least 60 residues in
20 length. Fragments of Rps2 polypeptide can be generated by methods known to those skilled in the art. The ability of a candidate fragment to exhibit a biological activity of Rps2 can be assessed by those methods described herein. Also included in the invention are
25 Rps2 polypeptides containing residues that are not required for biological activity of the peptide, e.g., those added by alternative mRNA splicing or alternative protein processing events.

Other embodiments are within the following claims.

- 66 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: Ausubel, Frederick M.
Staskawicz, Brian J.
Brent, Andrew F.
Dahlbeck, Douglas
Katagiri, Fumiaki
Kunkel, Barbara N.
Mindrinos, Michael N.
Yu, Guo-Liang

(ii) TITLE OF INVENTION: RPS2 GENE AND USES THEREOF

(iii) NUMBER OF SEQUENCES: 201

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Fish & Richardson
(B) STREET: 225 Franklin Street Suite 3100
(C) CITY: Boston
(D) STATE: MA
(E) COUNTRY: USA
(F) ZIP: 02110-2904

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30B

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: US 08/227,360
(B) FILING DATE: 13-APR-1994
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Clark, Paul T.
(B) REGISTRATION NUMBER: 30,162
(C) REFERENCE/DOCKET NUMBER: 00786/230001

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (617) 542-5070
(B) TELEFAX: (617) 542-8906
(C) TELEX: 100254

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2903 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

AAGTAAAAGA AAGAGCGAGA AATCATCGAA ATGGATTCA TCTCATCTCT TATCGTTGGC	60
TGTGCTCAGG TGTTGTGTGA ATCTATGAAT ATGGCCGAGA GAAGAGGACA TAAGACTGAT	120

- 67 -

CTTAGACAAG	CCATCACTGA	TCTTGAAACA	GCCATCGGTG	ACTTGAAGGC	CATACGTGAT	180
GACCTGACTT	TACGGATCCA	ACAAGACGGT	CTAGAGGGAC	GAAGCTGCTC	AAATCGTGCC	240
AGAGAGTGGC	TTAGTGCGGT	GCAAGTAACG	GAGACTAAAA	CAGCCCTACT	TTTAGTGAGG	300
TTTAGGCGTC	GGGAACAGAG	GACGCGAATG	AGGAGGAGAT	ACCTCAGTTG	TTTCGGTTGT	360
GGCGACTACA	AACTGTGCAA	GAAGGTTTCT	GCCATATTGA	AGAGCATTGG	TGAGCTGAGA	420
GAACGCTCTG	AAGCTATCAA	AACAGATGGC	GGGTCAATTC	AAGTAACTTG	TAGAGAGATA	480
CCCATCAAGT	CCGTTGTCTG	AAATACCACG	ATGATGGAAC	AGGTTTTGGA	ATTTCTCAGT	540
GAAGAAGAAG	AAAGAGGAAT	CATTGGTGTT	TATGGACCTG	GTGGGGTTGG	GAAGACAACG	600
TTAATGCAGA	GCATTAACAA	CGAGCTGATC	ACAAAAGGAC	ATCAGTATGA	TGTA CTGATT	660
TGGGTTCAAA	TGTCCAGAGA	ATTCGGCGAG	TGTACAATTC	AGCAAGCCGT	TGGAGCACGG	720
TTGGGTTTAT	CTTGGGACGA	GAAGGAGACC	GGCGAAAACA	GAGCTTTGAA	GATATACAGA	780
GCTTTGAGAC	AGAAACGTTT	CTTGTTGTTG	CTAGATGATG	TCTGGGAAGA	GATAGACTTG	840
GAGAAACTG	GAGTTCCTCG	ACCTGACAGG	GAAAACAAAT	GCAAGGTGAT	GTTACACACA	900
CGGTCTATAG	CATTATGCAA	CAATATGGGT	GCGGAATACA	AGTTGAGAGT	GGAGTTTCTG	960
GAGAAGAAAC	ACGCGTGGGA	GCTGTTCTGT	AGTAAGGTAT	GGAGAAAAGA	TCTTTTAGAG	1020
TCATCATCAA	TTCGCCGGCT	CGCGGAGATT	ATAGTGAGTA	AATGTGGAGG	ATTGCCACTA	1080
GCGTTGATCA	CTTTAGGAGG	AGCCATGGCT	CATAGAGAGA	CAGAAGAAGA	GTGGATCCAT	1140
GCTAGTGAAG	TTCTGACTAG	ATTTCCAGCA	GAGATGAAGG	GTATGAACTA	TGTATTTGCC	1200
CTTTTGAAAT	TCAGCTACGA	CAACCTCGAG	AGTGATCTGC	TTCGGTCTTG	TTTCTTGATC	1260
TGCGCTTTAT	TCCCAGAAGA	ACATTCTATA	GAGATCGAGC	AGCTTGTTGA	GTA CTGGGTC	1320
GGCGAAGGGT	TTCTCACCAG	CTCCCATGGC	GTTAACACCA	TTTACAAGGG	ATATTTTCTC	1380
ATTGGGGATC	TGAAAGCGGC	ATGTTTGTTG	GAAACCGGAG	ATGAGAAAAC	ACAGGTGAAG	1440
ATGCATAATG	TGGTCAGAAG	CTTGCATTG	TGGATGGCAT	CTGAACAGGG	GACTTATAAG	1500
GAGCTGATCC	TAGTTGAGCC	TAGCATGGGA	CATACTGAAG	CTCCTAAAGC	AGAAA CTGG	1560
CGACAAGCGT	TGGTGATCTC	ATTGTTAGAT	AACAGAATCC	AGACCTTGCC	TGAAAACTC	1620
ATATGCCCCG	AACTGACAAC	ACTGATGCTC	CAACAGAACA	GCTCTTTGAA	GAAGATTCCA	1680
ACAGGGTTTT	TCATGCATAT	GCCTGTTCTC	AGAGTCTTGG	ACTTGTCGTT	CACAAGTATC	1740
ACTGAGATTC	CGTTGTCTAT	CAAGTATTTG	GTGGAGTTGT	ATCATCTGTC	TATGTCAGGA	1800
ACAAAGATAA	GTGTATTGCC	ACAGGAGCTT	GGGAATCTTA	GAAA ACTGAA	GCATCTGGAC	1860
CTACAAAGAA	CTCAGTTTCT	TCAGACGATC	CCACGAGATG	CCATATGTTG	GCTGAGCAAG	1920
CTCGAGGTTT	TGA ACTTGTA	CTACAGTTAC	GCCGGTTGGG	AACTGCAGAG	CTTTGGAGAA	1980
GATGAAGCAG	AAGAACTCGG	ATTCGCTGAC	TTGGAATACT	TGGAAAACCT	AACCACACTC	2040

- 68 -

```

GGTATCACTG TTCTCTCATT GGAGACCCTA AAAACTCTCT TCGAGTTCGG TGCTTTGCAT      2100
AAACATATAC AGCATCTCCA CGTTGAAGAG TGCAATGAAC TCCTCTACTT CAATCTCCCA      2160
TCACTCACTA ACCATGGCAG GAACCTGAGA AGACTTAGCA TTAAAAGTTG CCATGACTTG      2220
GAGTACCTGG TCACACCCGC AGATTTTGAA AATGATTGGC TTCCGAGTCT AGAGGTTCTG      2280
ACGTTACACA GCCTTCACAA CTTAACCAGA GTGTGGGGAA ATTCTGTAAG CCAAGATTGT      2340
CTGCGGAATA TCCGTTGCAT AAACATTTCA CACTGCAACA AGCTGAAGAA TGTCTCATGG      2400
GTTCAGAAAC TCCCAAAGCT AGAGGTGATT GAACTGTTCTG ACTGCAGAGA GATAGAGGAA      2460
TTGATAAGCG AACACGAGAG TCCATCCGTC GAAGATCCAA CATTGTTCCC AAGCCTGAAG      2520
ACCTTGAGAA CTAGGGATCT GCCAGAACTA AACAGCATCC TCCCATCTCG ATTTTCATTC      2580
CAAAAAGTTG AAACATTAGT CATCACAAAT TGCCCCAGAG TTAAGAAACT GCCGTTTCAG      2640
GAGAGGAGGA CCCAGATGAA CTTGCCAACA GTTTATTGTG AGGAGAAATG GTGGAAGCA      2700
CTGGA AAAAG ATCAACCAA CGAAGAGCTT TGTTATTTAC CGCGCTTTGT TCCAAATTGA      2760
TATAAGAGCT AAGAGCACTC TGTACAAATA TGTCCATTCA TAAGTAGCAG GAAGCCAGGA      2820
AGGTTGTTCC AGTGAAGTCA TCACTTTCC ACATAGCCAC AAAACTAGAG ATTATGTAAT      2880
CATAAAAACC AAACATATCCG CGA                                          2903

```

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 885 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

```

Lys Lys Glu Arg  Glu Ile Ile Glu Met  Asp Phe Ile Ser Ser Leu Ile
 1          5          10          15
Val Gly Cys Ala Gln Val Leu Cys Glu Ser Met Asn Met Ala Glu Arg
 20          25          30
Arg Gly His Lys Thr Asp Leu Arg Gln Ala Ile Thr Asp Leu Arg Ile
 35          40          45
Gln Gln Asp Gly Leu Glu Gly Arg Ser Cys Ser Asn Arg Ala Arg Glu
 50          55          60
Trp Leu Ser Ala Val Gln Val Thr Glu Thr Lys Thr Ala Leu Leu Leu
 65          70          75          80
Val Arg Phe Arg Arg Arg Glu Gln Arg Thr Arg Met Arg Arg Arg Tyr
 85          90          95

```

- 69 -

Leu Ser Cys Phe Gly Cys Ala Asp Tyr Lys Leu Cys Lys Lys Val Ser
 100 105 110
 Ala Ile Leu Lys Ser Ile Gly Glu Leu Arg Glu Arg Ser Glu Ala Ile
 115 120 125
 Lys Thr Asp Gly Gly Ser Ile Gln Val Thr Cys Arg Glu Ile Pro Ile
 130 135 140
 Lys Ser Val Val Gly Asn Thr Thr Met Met Glu Gln Val Leu Glu Phe
 145 150 155 160
 Leu Ser Glu Glu Glu Glu Arg Gly Ile Ile Gly Val Tyr Gly Pro Gly
 165 170 175
 Gly Val Gly Lys Thr Thr Leu Met Gln Ser Ile Asn Asn Glu Leu Ile
 180 185 190
 Thr Lys Gly His Gln Tyr Asp Val Leu Ile Trp Val Gln Met Ser Arg
 195 200 205
 Glu Phe Gly Glu Cys Thr Ile Gln Gln Ala Val Gly Ala Arg Leu Gly
 210 215 220
 Leu Ser Trp Asp Glu Lys Glu Thr Gly Glu Asn Arg Ala Leu Lys Ile
 225 230 235 240
 Tyr Arg Ala Leu Arg Gln Lys Arg Phe Leu Leu Leu Leu Asp Asp Val
 245 250 255
 Trp Glu Glu Ile Asp Leu Glu Lys Thr Gly Val Pro Arg Pro Asp Arg
 260 265 270
 Glu Asn Lys Cys Lys Val Met Phe Thr Thr Arg Ser Ile Ala Leu Cys
 275 280 285
 Asn Asn Met Gly Ala Glu Tyr Lys Leu Arg Val Glu Phe Leu Glu Lys
 290 295 300
 Lys His Ala Trp Glu Leu Phe Cys Ser Lys Val Trp Arg Lys Asp Leu
 305 310 315 320
 Leu Glu Ser Ser Ser Ile Arg Arg Leu Ala Glu Ile Ile Val Ser Lys
 325 330 335
 Cys Gly Gly Leu Pro Leu Ala Leu Ile Thr Leu Gly Gly Ala Met Ala
 340 345 350
 His Arg Glu Thr Glu Glu Glu Trp Ile His Ala Ser Glu Val Leu Thr
 355 360 365
 Arg Phe Pro Ala Glu Met Lys Gly Met Asn Tyr Val Phe Ala Leu Leu
 370 375 380
 Lys Phe Ser Tyr Asp Asn Leu Glu Ser Asp Leu Leu Arg Ser Cys Phe
 385 390 395 400
 Leu Tyr Cys Ala Leu Phe Pro Glu Glu His Ser Ile Glu Ile Glu Gln
 405 410 415
 Leu Val Glu Tyr Trp Val Gly Glu Gly Phe Leu Thr Ser Ser His Gly
 420 425 430

- 70 -

Val Asn Thr Ile Tyr Lys Gly Tyr Phe Leu Ile Gly Asp Leu Lys Ala
 435 440 445
 Ala Cys Leu Leu Glu Thr Gly Asp Glu Lys Thr Gln Val Lys Met His
 450 455 460
 Asn Val Val Arg Ser Phe Ala Leu Trp Met Ala Ser Glu Gln Gly Thr
 465 470 475 480
 Tyr Lys Glu Leu Ile Leu Val Glu Pro Ser Met Gly His Thr Glu Ala
 485 490 495
 Pro Lys Ala Glu Asn Trp Arg Gln Ala Leu Val Ile Ser Leu Leu Asp
 500 505 510
 Asn Arg Ile Gln Thr Leu Pro Glu Lys Leu Ile Cys Pro Lys Leu Thr
 515 520 525
 Thr Leu Met Leu Gln Gln Asn Ser Ser Leu Lys Lys Ile Pro Thr Gly
 530 535 540
 Phe Phe Met His Met Pro Val Leu Arg Val Leu Asp Leu Ser Phe Thr
 545 550 555 560
 Ser Ile Thr Glu Ile Pro Leu Ser Ile Lys Tyr Leu Val Glu Leu Tyr
 565 570 575
 His Leu Ser Met Ser Gly Thr Lys Ile Ser Val Leu Pro Gln Glu Leu
 580 585 590
 Gly Asn Leu Arg Lys Leu Lys His Leu Asp Leu Gln Arg Thr Gln Phe
 595 600 605
 Leu Gln Thr Ile Pro Arg Asp Ala Ile Cys Trp Leu Ser Lys Leu Glu
 610 615 620
 Val Leu Asn Leu Tyr Tyr Ser Tyr Ala Gly Trp Glu Leu Gln Ser Phe
 625 630 635 640
 Gly Glu Asp Glu Ala Glu Glu Leu Gly Phe Ala Asp Leu Glu Tyr Leu
 645 650 655
 Glu Asn Leu Thr Thr Leu Gly Ile Thr Val Leu Ser Leu Glu Thr Leu
 660 665 670
 Lys Thr Leu Phe Glu Phe Gly Ala Leu His Lys His Ile Gln His Leu
 675 680 685
 His Val Glu Glu Cys Asn Glu Leu Leu Tyr Phe Asn Leu Pro Ser Leu
 690 695 700
 Thr Asn His Gly Arg Asn Leu Arg Arg Leu Ser Ile Lys Ser Cys His
 705 710 715 720
 Asp Leu Glu Tyr Leu Val Thr Pro Ala Asp Phe Glu Asn Asp Trp Leu
 725 730 735
 Pro Ser Leu Glu Val Leu Thr Leu His Ser Leu His Asn Leu Arg Cys
 740 745 750
 Ile Asn Ile Ser His Cys Asn Lys Leu Lys Asn Val Ser Trp Val Gln
 755 760 765

- 71 -

Lys Leu Pro Lys Leu Glu Val Ile Glu Leu Phe Asp Cys Arg Glu Ile
 770 775 780
 Glu Glu Leu Ile Ser Glu His Glu Ser Pro Ser Val Glu Asp Pro Thr
 785 790 795 800
 Leu Phe Pro Ser Leu Lys Thr Leu Arg Thr Arg Asp Leu Pro Glu Leu
 805 810 815
 Asn Ser Ile Leu Pro Ser Arg Phe Ser Phe Gln Lys Val Glu Thr Leu
 820 825 830
 Val Ile Thr Asn Cys Pro Arg Val Lys Lys Leu Pro Phe Gln Glu Arg
 835 840 845
 Arg Thr Gln Met Asn Leu Pro Thr Val Tyr Cys Glu Glu Lys Trp Trp
 850 855 860
 Lys Ala Leu Glu Lys Asp Gln Pro Asn Glu Glu Leu Cys Tyr Leu Pro
 865 870 875 880
 Arg Phe Val Pro Asn
 885

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Glu His Ser Val Gln Ile Cys Pro Phe Ile Ser Ser Arg Lys Pro Gly
 1 5 10 15
 Arg Leu Phe Gln
 20

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 6 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Ser His Gln Leu Ser Thr
 1 5

- 72 -

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 11 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: not relevant
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Arg Leu Cys Asn His Lys Asn Gln Thr Ile Arg
1 5 10

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 28 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: not relevant
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Ser Lys Arg Lys Ser Glu Lys Ser Ser Lys Trp Ile Ser Ser His Leu
1 5 10 15
Leu Ser Leu Ala Val Leu Arg Cys Cys Val Asn Leu
20 25

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 25 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: not relevant
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Ile Trp Arg Arg Glu Glu Asp Ile Arg Leu Ile Leu Asp Lys Pro Ser
1 5 10 15
Leu Ile Leu Lys Gln Pro Ser Val Thr
20 25

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:

- 73 -

- (A) LENGTH: 6 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Arg Pro Tyr Val Met Thr
1 5

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 8 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Leu Tyr Gly Ser Asn Lys Thr Val
1 5

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 17 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Arg Asp Glu Ala Ala Gln Ile Val Pro Glu Ser Gly Leu Val Arg Cys
1 5 10 15
Lys

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 8 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

- 74 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Arg Arg Leu Lys Gln Pro Tyr Phe
1 5

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Gly Leu Gly Val Gly Asn Arg Gly Arg Glu
1 5 10

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Gly Gly Asp Thr Ser Val Val Ser Val Val Pro Thr Thr Asn Cys Ala
1 5 10 15

Arg Arg Phe Leu Pro Tyr
20

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Arg Ala Leu Val Ser

- 75 -

1 5

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

Glu	Asn	Ala	Leu	Lys	Leu	Ser	Lys	Gln	Met	Ala	Gly	Gln	Phe	Lys
1				5					10					15

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Leu	Val	Glu	Arg	Tyr	Pro	Ser	Ser	Pro	Leu	Ser	Glu	Ile	Pro	Arg
1				5					10					15

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 29 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

Trp	Asn	Arg	Phe	Trp	Asn	Phe	Ser	Val	Lys	Lys	Lys	Lys	Glu	Glu	Ser
1				5					10						15
Leu	Val	Phe	Met	Asp	Leu	Val	Gly	Leu	Gly	Arg	Gln	Arg			
			20					25							

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7 amino acids

- 76 -

- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Cys Arg Ala Leu Thr Thr Ser
1 5

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 9 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Ser Gln Lys Asp Ile Ser Met Met Tyr
1 5

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 36 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

Phe Gly Phe Lys Cys Pro Glu Asn Ser Ala Ser Val Gln Phe Ser Lys
1 5 10 15

Pro Leu Glu His Gly Trp Val Tyr Leu Gly Thr Arg Arg Arg Pro Ala
20 25 30

Lys Thr Glu Leu
35

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 5 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

- 77 -

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Arg Tyr Thr Glu Leu
1 5

(2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 8 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: not relevant
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

Asp Arg Asn Val Ser Cys Cys Cys
1 5

(2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 6 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: not relevant
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

Met Met Ser Gly Lys Arg
1 5

(2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 17 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: not relevant
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

Thr Trp Arg Lys Leu Glu Phe Leu Asp Leu Thr Gly Lys Thr Asn Ala
1 5 10 15

- 78 -

Arg

(2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 6 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

Cys Ser Arg His Gly Leu
1 5

(2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 11 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

His Tyr Ala Thr Ile Trp Val Arg Asn Thr Ser
1 5 10

(2) INFORMATION FOR SEQ ID NO:27:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 23 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

Glu Trp Ser Phe Trp Arg Arg Asn Thr Arg Gly Ser Cys Ser Val Val
1 5 10 15

Arg Tyr Gly Glu Lys Ile Phe
20

(2) INFORMATION FOR SEQ ID NO:28:

- (i) SEQUENCE CHARACTERISTICS:

- 79 -

- (A) LENGTH: 11 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

Ser His His Gln Phe Ala Gly Ser Arg Arg Leu
1 5 10

(2) INFORMATION FOR SEQ ID NO:29:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 7 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

Val Asn Val Glu Asp Cys His
1 5

(2) INFORMATION FOR SEQ ID NO:30:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 19 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

Glu Glu Pro Trp Leu Ile Glu Arg Gln Lys Lys Ser Gly Ser Met Leu
1 5 10 15

Val Lys Phe

(2) INFORMATION FOR SEQ ID NO:31:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 6 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

- 80 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

Leu Asp Phe Gln Gln Arg
1 5

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 6 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: not relevant
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

Thr Met Tyr Leu Pro Phe
1 5

(2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 27 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: not relevant
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

Asn Ser Ala Thr Thr Thr Ser Arg Val Ile Cys Phe Gly Leu Val Ser
1 5 10 15

Cys Thr Ala Leu Tyr Ser Gln Lys Asn Ile Leu
20 25

(2) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 33 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: not relevant
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

Arg Ser Ser Ser Leu Leu Ser Thr Gly Ser Ala Lys Gly Phe Ser Pro

- 81 -

1	5	10	15
Ala	Pro	Met	Ala
	Leu	Thr	Pro
		Phe	Thr
			Arg
			Asp
			Ile
			Phe
			Ser
			Leu
			Gly
	20	25	30

Ile

(2) INFORMATION FOR SEQ ID NO:35:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 14 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

Lys	Arg	His	Val	Cys	Trp	Lys	Pro	Glu	Met	Arg	Lys	His	Arg
1				5					10				

(2) INFORMATION FOR SEQ ID NO:36:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 22 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

Arg	Cys	Ile	Met	Trp	Ser	Glu	Ala	Leu	His	Cys	Gly	Trp	His	Leu	Asn
1				5					10					15	

Arg	Gly	Leu	Ile	Arg	Ser
					20

(2) INFORMATION FOR SEQ ID NO:37:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

Leu	Ser	Leu	Ala	Trp	Asp	Ile	Leu	Lys	Leu	Leu	Lys	Gln	Lys	Thr	Gly
1				5					10					15	

- 82 -

Asp Lys Arg Trp
20

(2) INFORMATION FOR SEQ ID NO:38:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 15 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

Ile Thr Glu Ser Arg Pro Cys Leu Lys Asn Ser Tyr Ala Arg Asn
 1 5 10 15

(2) INFORMATION FOR SEQ ID NO:39:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 7 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

Cys Ser Asn Arg Thr Ala Leu
 1 5

(2) INFORMATION FOR SEQ ID NO:40:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 46 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

Arg Arg Phe Gln Gln Gly Phe Ser Cys Ile Cys Leu Phe Ser Glu Ser
 1 5 10 15

Trp Thr Cys Arg Ser Gln Val Ser Leu Arg Phe Arg Cys Leu Ser Ser
 20 25 30

Ile Trp Trp Ser Cys Ile Ile Cys Leu Cys Gln Glu Gln Arg
 35 40 45

- 83 -

(2) INFORMATION FOR SEQ ID NO:41:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: not relevant
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

Val Tyr Cys His Arg Ser Leu Gly Ile Leu Glu Asn
1 5 10

(2) INFORMATION FOR SEQ ID NO:42:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: not relevant
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

Ser Ile Trp Thr Tyr Lys Glu Leu Ser Phe Phe Arg Arg Ser His Glu
1 5 10 15
Met Pro Tyr Val Gly
20

(2) INFORMATION FOR SEQ ID NO:43:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: not relevant
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

Ala Ser Ser Arg Phe
1 5

(2) INFORMATION FOR SEQ ID NO:44:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 32 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: not relevant

- 84 -

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

Thr	Cys	Thr	Thr	Val	Thr	Pro	Val	Gly	Asn	Cys	Arg	Ala	Leu	Glu	Lys
1				5				10						15	

Met	Lys	Gln	Lys	Asn	Ser	Asp	Ser	Leu	Thr	Trp	Asn	Thr	Trp	Lys	Thr
			20					25					30		

(2) INFORMATION FOR SEQ ID NO:45:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

Pro	His	Ser	Val	Ser	Leu	Phe	Ser	His	Trp	Arg	Pro
1				5				10			

(2) INFORMATION FOR SEQ ID NO:46:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 38 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

Lys	Leu	Ser	Ser	Ser	Ser	Val	Leu	Cys	Ile	Asn	Ile	Tyr	Ser	Ile	Ser
1				5				10						15	

Thr	Leu	Lys	Ser	Ala	Met	Asn	Ser	Ser	Thr	Ser	Ile	Ser	His	His	Ser
			20					25					30		

Leu	Thr	Met	Ala	Gly	Thr
					35

(2) INFORMATION FOR SEQ ID NO:47:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 amino acids
- (B) TYPE: amino acid

- 85 -

- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

Glu Asp Leu Ala Leu Lys Val Ala Met Thr Trp Ser Thr Trp Ser His
1 5 10 15

Pro Gln Ile Leu Lys Met Ile Gly Phe Arg Val
20 25

(2) INFORMATION FOR SEQ ID NO:48:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 7 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

Arg Tyr Thr Ala Phe Thr Thr
1 5

(2) INFORMATION FOR SEQ ID NO:49:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 7 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

Pro Glu Cys Gly Glu Ile Leu
1 5

(2) INFORMATION FOR SEQ ID NO:50:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

- 86 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

Ala Lys Ile Val Cys Gly Ile Ser Val Ala
1 5 10

(2) INFORMATION FOR SEQ ID NO:51:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

Thr Phe His Thr Ala Thr Ser
1 5

(2) INFORMATION FOR SEQ ID NO:52:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 6 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

Phe Arg Asn Ser Gln Ser
1 5

(2) INFORMATION FOR SEQ ID NO:53:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 8 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

Leu Asn Cys Ser Thr Ala Glu Arg
1 5

(2) INFORMATION FOR SEQ ID NO:54:

- 87 -

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 16 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

Ala	Asn	Thr	Arg	Val	His	Pro	Ser	Lys	Ile	Gln	His	Cys	Ser	Gln	Ala
1				5					10					15	

(2) INFORMATION FOR SEQ ID NO:55:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 7 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

Glu	Leu	Gly	Ile	Cys	Gln	Asn
1				5		

(2) INFORMATION FOR SEQ ID NO:56:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 15 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

Thr	Ala	Ser	Ser	His	Leu	Asp	Phe	His	Ser	Lys	Lys	Leu	Lys	His
1				5					10					15

(2) INFORMATION FOR SEQ ID NO:57:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

- 88 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

Ser Ser Gln Ile Ala Pro Glu Leu Arg Asn Cys Arg Phe Arg Arg Gly
1 5 10 15

Gly Pro Arg

(2) INFORMATION FOR SEQ ID NO:58:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 47 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

Thr Cys Gln Gln Phe Ile Val Arg Arg Asn Gly Gly Lys His Trp Lys
1 5 10 15

Lys Ile Asn Gln Thr Lys Ser Phe Val Ile Tyr Arg Ala Leu Phe Gln
20 25 30

Ile Asp Ile Arg Ala Lys Ser Thr Leu Tyr Lys Tyr Val His Ser
35 40 45

(2) INFORMATION FOR SEQ ID NO:59:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:

Val Ala Gly Ser Gln Glu Gly Cys Ser Ser Glu Val Ile Asn Phe Pro
1 5 10 15

His Ser His Lys Thr Arg Asp Tyr Val Ile Ile Lys Thr Lys Leu Ser
20 25 30

Ala

(2) INFORMATION FOR SEQ ID NO:60:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 amino acids
- (B) TYPE: amino acid

- 89 -

- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:60:

Val	Lys	Glu	Arg	Ala	Arg	Asn	His	Arg	Asn	Gly	Phe	His	Leu	Ile	Ser
1				5				10					15		
Tyr	Arg	Trp	Leu	Cys	Ser	Gly	Val	Val							
			20				25								

(2) INFORMATION FOR SEQ ID NO:61:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:61:

Ile	Tyr	Glu	Tyr	Gly	Gly	Glu	Lys	Arg	Thr
1			5					10	

(2) INFORMATION FOR SEQ ID NO:62:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 5 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:62:

Leu	Glu	Gly	His	Thr
1			5	

(2) INFORMATION FOR SEQ ID NO:63:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 23 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

- 90 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:63:

Pro	Asp	Phe	Thr	Asp	Pro	Thr	Arg	Arg	Ser	Arg	Gly	Thr	Lys	Leu	Leu
1				5					10					15	
Lys Ser Cys Gln Arg Val Ala															
20															

(2) INFORMATION FOR SEQ ID NO:64:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 7 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:64:

Cys	Gly	Ala	Ser	Asn	Gly	Asp
1				5		

(2) INFORMATION FOR SEQ ID NO:65:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 8 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:65:

Asn	Ser	Pro	Thr	Phe	Ser	Glu	Val
1				5			

(2) INFORMATION FOR SEQ ID NO:66:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 35 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:66:

Ala	Ser	Gly	Thr	Glu	Asp	Ala	Asn	Glu	Glu	Glu	Ile	Pro	Gln	Leu	Phe
1				5				10						15	

- 91 -

Arg Leu Cys Arg Leu Gln Thr Val Gln Glu Gly Phe Cys His Ile Glu
 20 25 30

Glu His Trp
 35

(2) INFORMATION FOR SEQ ID NO:67:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 5 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:67:

Ala Glu Arg Thr Leu
 1 5

(2) INFORMATION FOR SEQ ID NO:68:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 13 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:68:

Ser Tyr Gln Asn Arg Trp Arg Val Asn Ser Ser Asn Leu
 1 5 10

(2) INFORMATION FOR SEQ ID NO:69:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 22 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:69:

Arg Asp Thr His Gln Val Arg Cys Arg Lys Tyr His Asp Asp Gly Thr
 1 5 10 15

Gly Phe Gly Ile Ser Gln
 20

- 92 -

(2) INFORMATION FOR SEQ ID NO:70:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:70:

Arg Arg Arg Lys Arg Asn His Trp Cys Leu Trp Thr Trp Trp Gly Trp
 1 5 10 15

Glu Asp Asn Val Asn Ala Glu His
 20

(2) INFORMATION FOR SEQ ID NO:71:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 10 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:71:

Gln Arg Ala Asp His Lys Arg Thr Ser Val
 1 5 10

(2) INFORMATION FOR SEQ ID NO:72:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 55 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:72:

Cys Thr Asp Leu Gly Ser Asn Val Gln Arg Ile Arg Arg Val Tyr Asn
 1 5 10 15

Ser Ala Ser Arg Trp Ser Thr Val Gly Phe Ile Leu Gly Arg Glu Gly
 20 25 30

Asp Arg Arg Lys Gln Ser Phe Glu Asp Ile Gln Ser Phe Glu Thr Glu
 35 40 45

- 93 -

Thr Phe Leu Val Val Ala Arg
50 55

(2) INFORMATION FOR SEQ ID NO:73:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 15 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:73:

Cys Leu Gly Arg Asp Arg Leu Gly Glu Asn Trp Ser Ser Ser Thr
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:74:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 9 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:74:

Arg Asp Arg Arg Arg Val Asp Pro Cys
1 5

(2) INFORMATION FOR SEQ ID NO:75:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 41 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:75:

Gln Gly Lys Gln Met Gln Gly Asp Val His Asp Thr Val Tyr Ser Ile
1 5 10 15

Met Gln Gln Tyr Gly Cys Gly Ile Gln Val Glu Ser Gly Val Ser Gly
20 25 30

Glu Glu Thr Arg Val Gly Ala Val Leu
35 40

- 94 -

(2) INFORMATION FOR SEQ ID NO:76:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: not relevant
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:76:

Gly Met Glu Lys Arg Ser Phe Arg Val Ile Ile Asn Ser Pro Ala Arg
1 5 10 15
Gly Asp Tyr Ser Glu
 20

(2) INFORMATION FOR SEQ ID NO:77:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 17 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: not relevant
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:77:

Met Trp Arg Ile Ala Thr Ser Val Asp His Phe Arg Arg Ser His Gly
1 5 10 15
Ser

(2) INFORMATION FOR SEQ ID NO:78:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 24 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: not relevant
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:78:

Ile Ser Ser Arg Asp Glu Gly Tyr Glu Leu Cys Ile Cys Pro Phe Glu
1 5 10 15
Ile Gln Leu Arg Gln Pro Arg Glu
 20

- 95 -

(2) INFORMATION FOR SEQ ID NO:79:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 24 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: not relevant
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:79:

Ser Ala Ser Val Leu Phe Leu Val Leu Arg Phe Ile Pro Arg Arg Thr
1 5 10 15
Phe Tyr Arg Asp Arg Ala Ala Cys
 20

(2) INFORMATION FOR SEQ ID NO:80:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 14 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: not relevant
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:80:

Val Leu Gly Arg Arg Arg Val Ser His Gln Leu Pro Trp Arg
1 5 10

(2) INFORMATION FOR SEQ ID NO:81:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 22 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: not relevant
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:81:

His His Leu Gln Gly Ile Phe Ser His Trp Gly Ser Glu Ser Gly Met
1 5 10 15
Phe Val Gly Asn Arg Arg
 20

(2) INFORMATION FOR SEQ ID NO:82:

- (i) SEQUENCE CHARACTERISTICS:

- 96 -

- (A) LENGTH: 7 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:82:

Glu Asn Thr Gly Glu Asp Ala
1 5

(2) INFORMATION FOR SEQ ID NO:83:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 43 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:83:

Lys Thr His Met Pro Glu Thr Asp Asn Thr Asp Ala Pro Thr Glu Gly
1 5 10 15

Leu Phe Glu Glu Asp Ser Asn Arg Val Phe His Ala Tyr Ala Cys Ser
20 25 30

Gln Ser Leu Gly Leu Val Val His Lys Tyr His
35 40

(2) INFORMATION FOR SEQ ID NO:84:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 11 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:84:

Cys Gly Gln Lys Leu Cys Ile Val Asp Gly Ile
1 5 10

(2) INFORMATION FOR SEQ ID NO:85:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 5 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant

- 97 -

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:85:

Gly Ala Asp Pro Ser
1 5

(2) INFORMATION FOR SEQ ID NO:86:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 14 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:86:

Ser Arg Lys Leu Ala Thr Ser Val Gly Asp Leu Ile Val Arg
1 5 10

(2) INFORMATION FOR SEQ ID NO:87:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 6 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:87:

Gln Asn Pro Asp Leu Ala
1 5

(2) INFORMATION FOR SEQ ID NO:88:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:88:

- 98 -

Asp Ser Val Val Tyr Gln Val Phe Gly Gly Val Val Ser Ser Val Tyr
 1 5 10 15
 Val Arg Asn Lys Asp Lys Cys Ile Ala Thr Gly Ala Trp Glu Ser
 20 25 30

(2) INFORMATION FOR SEQ ID NO:89:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 47 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:89:

Lys Thr Glu Ala Ser Gly Pro Thr Lys Asn Ser Val Ser Ser Asp Asp
 1 5 10 15
 Pro Thr Arg Cys His Met Leu Ala Glu Gln Ala Arg Gly Ser Glu Leu
 20 25 30
 Val Leu Gln Leu Arg Arg Leu Gly Thr Ala Glu Leu Trp Arg Arg
 35 40 45

(2) INFORMATION FOR SEQ ID NO:90:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 7 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:90:

Ser Arg Arg Thr Arg Ile Arg
 1 5

(2) INFORMATION FOR SEQ ID NO:91:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 30 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:91:

- 99 -

Leu Gly Ile Leu Gly Lys Pro Asn His Thr Arg Tyr His Cys Ser Leu
1 5 10 15

Ile Gly Asp Pro Lys Asn Ser Leu Arg Val Arg Cys Phe Ala
20 25 30

(2) INFORMATION FOR SEQ ID NO:92:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 7 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:92:

Thr Tyr Thr Ala Ser Pro Arg
1 5

(2) INFORMATION FOR SEQ ID NO:93:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:93:

Thr Pro Leu Leu Gln Ser Pro Ile Thr His
1 5 10

(2) INFORMATION FOR SEQ ID NO:94:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 8 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:94:

Pro Trp Gln Glu Pro Glu Lys Thr
1 5

(2) INFORMATION FOR SEQ ID NO:95:

- (i) SEQUENCE CHARACTERISTICS:

- 100 -

- (A) LENGTH: 10 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:95:

Leu Gly Val Pro Gly His Thr Arg Arg Phe
 1 5 10

(2) INFORMATION FOR SEQ ID NO:96:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 58 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:96:

Leu Ala Ser Glu Ser Arg Gly Ser Asp Val Thr Gln Pro Ser Gln Leu
 1 5 10 15

Asn Gln Ser Val Gly Lys Phe Cys Lys Pro Arg Leu Ser Ala Glu Tyr
 20 25 30

Pro Leu His Lys His Phe Thr Leu Gln Gln Ala Glu Glu Cys Leu Met
 35 40 45

Gly Ser Glu Thr Pro Lys Ala Arg Gly Asp
 50 55

(2) INFORMATION FOR SEQ ID NO:97:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 33 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:97:

Thr Val Arg Leu Gln Arg Asp Arg Gly Ile Asp Lys Arg Thr Arg Glu
 1 5 10 15

Ser Ile Arg Arg Arg Ser Asn Ile Val Pro Lys Pro Glu Asp Leu Glu
 20 25 30

- 101 -

Asn

(2) INFORMATION FOR SEQ ID NO:98:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 18 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:98:

Gly Ser Ala Arg Thr Lys Gln His Pro Pro Ile Ser Ile Phe Ile Pro
1 5 10 15

Lys Ser

(2) INFORMATION FOR SEQ ID NO:99:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:99:

Asn Ile Ser His His Lys Leu Pro Gln Ser
1 5 10

(2) INFORMATION FOR SEQ ID NO:100:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 18 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:100:

Glu Thr Ala Val Ser Gly Glu Glu Asp Pro Asp Glu Leu Ala Asn Ser
1 5 10 15

Leu Leu

- 102 -

(2) INFORMATION FOR SEQ ID NO:101:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 4 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:101:

Thr Ser His His
1

(2) INFORMATION FOR SEQ ID NO:102:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 13 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:102:

Glu Leu Arg Ala Leu Cys Thr Asn Met Ser Ile His Lys
1 5 10

(2) INFORMATION FOR SEQ ID NO:103:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 23 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:103:

Gln Glu Ala Arg Lys Val Val Pro Val Lys Ser Ser Thr Phe His Ile
1 5 10 15

Ala Thr Lys Leu Glu Ile Met
20

(2) INFORMATION FOR SEQ ID NO:104:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 6 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant

- 103 -

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:104:

Lys Pro Asn Tyr Pro Arg
1 5

(2) INFORMATION FOR SEQ ID NO:105:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1491 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:105:

ATCGATTGAT CTCTGGCTCA GTGCGAGTAG TCCATTGAG AGCAGTCGTA GCCCCGCGTG	60
GCGCATCATG GAGCTATTTG GAATTTTCGC AGGGTTATCG ATTCGTAGTG GGAACCCATT	120
CATTGTTTGG AACCACCAAC GGACGACTTA ACAAGCTCCC CGAGGTGCAT GATGAAAATT	180
GCTCCAGTTG CCATAAATCA CAGCCCGCTC AGCAGGGAGG TCCCGTCACA CGCGGCACCC	240
ACTCAGGCAA AGCAAACCAA CTTTCAATCT GAAGCTGGCG ATTTAGATGC AAGAAAAAGT	300
AGCGCTTCAA GCCCGGAAAC CCGCGCATT CTGCTACTA AGACAGTACT CGGGAGACAC	360
AAGATAGAGG TTCCGGCCTT TGGAGGGTGG TTCAAAAAGA AATCATCTAA GCACGAGACG	420
GGCGGTTCAA GTGCCAACGC AGATAGTTCT AGCGTGGCTT CCGATTCCAC CGAAAAACCT	480
TTGTTCCGTC TCACGCACGT TCCTTACGTA TCCCAAGGTA ATGAGCGAAT GGGATGTTGG	540
TATGCCTGCG CAAGAATGGT TGGCCATTCT GTCGAAGCTG GGCCTCGCCT AGGGCTGCCG	600
GAGCTCTATG AGGGAAGGGA GCGGCCAGCT GGGCTACAAG ATTTTTCAGA TG TAGAAAGG	660
TTTATTACACA ATGAAGGATT AACTCGGGTA GACCTTCCAG ACAATGAGAG ATTTACACAC	720
GAAGAGTTGG GTGCACTGTT GTATAAGCAC GGGCCGATTA TATTGGGTG GAAAACTCCG	780
AATGACAGCT GGCACATGTC GGTCTCACT GGTGTCGATA AAGAGACGTC GTCCATTACT	840
TTTACGATC CCCGACAGG GCGGACCTA GCAATGCCGC TCGATTACTT TAATCAGCGA	900
TTGGCATGGC AGGTTCCACA CGCAATGCTC TACCGCTAAG TAGCAGGGTA TCTTACGTCG	960
GCGGCATCAT GACAAGCCCA TGATGCCGCC AGCAGCTACC TGAATGCCGT CTGGCTTTTT	1020
GGTCCCTATT GTCGTATCCG GAAGATGACG TCAAAGAATC TCGGCAAGAG CTTTCTTGCT	1080

- 104 -

CGACTCCTCA GCTTCCGGAT CGATCAGGTC GCTTGCCAGA GCGCGCTTGT CCATGAGCAT 1140
 CTGCCACAGC TGCTGGTCGA TGGTGTCTTC AGCTAAAGGG ATTTTGACGA CAACCATGCG 1200
 CAACTGCCCCG TTGCGATACG CTCGATCCTG AAGCCCCGGT GTCCATGGCA GCCCCAAGAA 1260
 AAAGACATAG TTCGCCGCTG TGAGGTTGTA GCCTGTGCCG GCGGCCGACC TGGTCCCGAT 1320
 AACACCCTG CAGTCCGGAT CCTGCTGGAA AGCATCAATC GCCTTCTGCC GCTTCTTGGG 1380
 CGAGTCACTG CCCACCAACG TCACGCACCC GACGCCAAGC TTGAGGCAGT GCTCCCGCAA 1440
 CGTGGCCACG GATTCTGAT ACTCGCAGAA GAGGATCACC TTGTCGTCGA C 1491

(2) INFORMATION FOR SEQ ID NO:106:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 255 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:106:

Met Lys Ile Ala Pro Val Ala Ile Asn His Ser Pro Leu Ser Arg Glu
 1 5 10 15
 Val Pro Ser His Ala Ala Pro Thr Gln Ala Lys Gln Thr Asn Leu Gln
 20 25 30
 Ser Glu Ala Gly Asp Leu Asp Ala Arg Lys Ser Ser Ala Ser Ser Pro
 35 40 45
 Glu Thr Arg Ala Leu Leu Ala Thr Lys Thr Val Leu Gly Arg His Lys
 50 55 60
 Ile Glu Val Pro Ala Phe Gly Gly Trp Phe Lys Lys Lys Ser Ser Lys
 65 70 75 80
 His Glu Thr Gly Gly Ser Ser Ala Asn Ala Asp Ser Ser Ser Val Ala
 85 90 95
 Ser Asp Ser Thr Glu Lys Pro Leu Phe Arg Leu Thr His Val Pro Tyr
 100 105 110
 Val Ser Gln Gly Asn Glu Arg Met Gly Cys Trp Tyr Ala Cys Ala Arg
 115 120 125
 Met Val Gly His Ser Val Glu Ala Gly Pro Arg Leu Gly Leu Pro Glu
 130 135 140
 Leu Tyr Glu Gly Arg Glu Ala Pro Ala Gly Leu Gln Asp Phe Ser Asp
 145 150 155 160
 Val Glu Arg Phe Ile His Asn Glu Gly Leu Thr Arg Val Asp Leu Pro
 165 170 175
 Asp Asn Glu Arg Phe Thr His Glu Glu Leu Gly Ala Leu Leu Tyr Lys

- 105 -

	180		185		190
His Gly Pro Ile Ile Phe Gly Trp Lys Thr Pro Asn Asp Ser Trp His					
195		200		205	
Met Ser Val Leu Thr Gly Val Asp Lys Glu Thr Ser Ser Ile Thr Phe					
210		215		220	
His Asp Pro Arg Gln Gly Pro Asp Leu Ala Met Pro Leu Asp Tyr Phe					
225		230		235	240
Asn Gln Arg Leu Ala Trp Gln Val Pro His Ala Met Leu Tyr Arg					
	245		250		255

(2) INFORMATION FOR SEQ ID NO:107:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1209 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:107:

Met Asn Pro Ser Gly Ser Phe Pro Ser Val Glu Tyr Glu Val Phe Leu					
1	5		10		15
Ser Phe Arg Gly Pro Asp Thr Arg Glu Gln Phe Thr Asp Phe Leu Tyr					
	20		25		30
Gln Ser Leu Arg Arg Tyr Lys Ile His Thr Phe Arg Asp Asp Asp Glu					
	35		40		45
Leu Leu Lys Gly Lys Glu Ile Gly Pro Asn Leu Leu Arg Ala Ile Asp					
	50		55		60
Gln Ser Lys Ile Tyr Val Pro Ile Ile Ser Ser Gly Tyr Ala Asp Ser					
65	70		75		80
Lys Trp Cys Leu Met Glu Leu Ala Glu Ile Val Arg Arg Gln Glu Glu					
	85		90		95
Asp Pro Arg Arg Ile Ile Leu Pro Ile Phe Tyr Met Val Asp Pro Ser					
	100		105		110
Asp Val Arg His Gln Thr Gly Cys Tyr Lys Lys Ala Phe Arg Lys His					
	115		120		125
Ala Asn Lys Phe Asp Gly Gln Thr Ile Gln Asn Trp Lys Asp Ala Leu					
	130		135		140
Lys Lys Val Gly Asp Leu Lys Gly Trp His Ile Gly Lys Asn Asp Lys					
145	150		155		160
Gln Gly Ala Ile Ala Asp Lys Val Ser Ala Asp Ile Trp Ser His Ile					
	165		170		175

- 106 -

Ser Lys Glu Asn Leu Ile Leu Glu Thr Asp Glu Leu Val Gly Ile Asp
 180 185 190
 Asp His Ile Thr Ala Val Leu Glu Lys Leu Ser Leu Asp Ser Glu Asn
 195 200 205
 Val Thr Met Val Gly Leu Tyr Gly Met Gly Gly Ile Gly Lys Thr Thr
 210 215 220
 Thr Ala Lys Ala Val Tyr Asn Lys Ile Ser Ser Cys Phe Asp Cys Cys
 225 230 235 240
 Cys Phe Ile Asp Asn Ile Arg Glu Thr Gln Glu Lys Asp Gly Val Val
 245 250 255
 Val Leu Gln Lys Lys Leu Val Ser Glu Ile Leu Arg Ile Asp Ser Gly
 260 265 270
 Ser Val Gly Phe Asn Asn Asp Ser Gly Gly Arg Lys Thr Ile Lys Glu
 275 280 285
 Arg Val Ser Arg Phe Lys Ile Leu Val Val Leu Asp Asp Val Asp Glu
 290 295 300
 Lys Phe Lys Phe Glu Asp Met Leu Gly Ser Pro Lys Asp Phe Ile Ser
 305 310 315 320
 Gln Ser Arg Phe Ile Ile Thr Ser Arg Ser Met Arg Val Leu Gly Thr
 325 330 335
 Leu Asn Glu Asn Gln Cys Lys Leu Tyr Glu Val Gly Ser Met Ser Lys
 340 345 350
 Pro Arg Ser Leu Glu Leu Phe Ser Lys His Ala Phe Lys Lys Asn Thr
 355 360 365
 Pro Pro Ser Ser Tyr Tyr Glu Thr Leu Ala Asn Asp Val Val Asp Thr
 370 375 380
 Thr Ala Gly Leu Pro Leu Thr Leu Lys Val Ile Gly Ser Leu Leu Phe
 385 390 395 400
 Lys Gln Glu Ile Ala Val Trp Glu Asp Thr Leu Glu Gln Leu Arg Arg
 405 410 415
 Thr Leu Asn Leu Asp Glu Val Tyr Asp Arg Leu Lys Ile Ser Tyr Asp
 420 425 430
 Ala Leu Asn Pro Glu Ala Lys Glu Ile Phe Leu Asp Ile Ala Cys Phe
 435 440 445
 Phe Ile Gly Gln Asn Lys Glu Glu Pro Tyr Tyr Met Trp Thr Asp Cys
 450 455 460
 Asn Phe Tyr Pro Ala Ser Asn Ile Ile Phe Leu Ile Gln Arg Cys Met
 465 470 475 480
 Ile Gln Val Gly Asp Asp Asp Glu Phe Lys Met His Asp Gln Leu Arg
 485 490 495
 Asp Met Gly Arg Glu Ile Val Arg Arg Glu Asp Val Leu Pro Trp Lys
 500 505 510

- 107 -

Ser Arg Ile Trp Ser Ala Glu Glu Gly Ile Asp Leu Leu Leu Asn Lys
 515 520 525
 Arg Lys Gly Ser Ser Lys Val Lys Ala Ile Ser Ile Pro Trp Gly Val
 530 535 540
 Lys Tyr Glu Phe Lys Ser Glu Cys Phe Leu Asn Leu Ser Glu Leu Arg
 545 550 555 560
 Tyr Leu His Ala Arg Glu Ala Met Leu Thr Gly Asp Phe Asn Asn Leu
 565 570 575
 Leu Pro Asn Leu Lys Trp Leu Glu Leu Pro Phe Tyr Lys His Gly Glu
 580 585 590
 Asp Asp Pro Pro Leu Thr Asn Tyr Thr Met Lys Asn Leu Ile Ile Val
 595 600 605
 Ile Leu Glu His Ser His Ile Thr Ala Asp Asp Trp Gly Gly Trp Arg
 610 615 620
 His Met Met Lys Met Ala Glu Arg Leu Lys Val Val Arg Leu Ala Ser
 625 630 635 640
 Asn Tyr Ser Leu Tyr Gly Arg Arg Val Arg Leu Ser Asp Cys Trp Arg
 645 650 655
 Phe Pro Lys Ser Ile Glu Val Leu Ser Met Thr Ala Ile Glu Met Asp
 660 665 670
 Glu Val Asp Ile Gly Glu Leu Lys Lys Leu Lys Thr Leu Val Leu Lys
 675 680 685
 Pro Cys Pro Ile Gln Lys Ile Ser Gly Gly Thr Phe Gly Met Leu Lys
 690 695 700
 Gly Leu Arg Glu Leu Cys Leu Glu Phe Asn Trp Gly Thr Asn Leu Arg
 705 710 715 720
 Glu Val Val Ala Asp Ile Gly Gln Leu Ser Ser Leu Lys Val Leu Lys
 725 730 735
 Thr Gly Ala Lys Glu Val Glu Ile Asn Glu Phe Pro Leu Gly Leu Lys
 740 745 750
 Thr Glu Leu Ser Thr Ser Ser Arg Ile Pro Asn Asn Leu Ser Gln Leu
 755 760 765
 Leu Asp Leu Glu Val Leu Lys Val Tyr Asp Cys Lys Asp Gly Phe Asp
 770 775 780
 Met Pro Pro Ala Ser Pro Ser Glu Asp Glu Ser Ser Val Trp Trp Lys
 785 790 795 800
 Val Ser Lys Leu Lys Ser Leu Gln Leu Glu Lys Thr Arg Ile Asn Val
 805 810 815
 Asn Val Val Asp Asp Ala Ser Ser Gly Gly His Leu Pro Arg Tyr Leu
 820 825 830
 Leu Pro Thr Ser Leu Thr Tyr Leu Lys Ile Tyr Gln Cys Thr Glu Pro
 835 840 845

- 108 -

Thr Trp Leu Pro Gly Ile Glu Asn Leu Glu Asn Leu Thr Ser Leu Glu
 850 855 860
 Val Asn Asp Ile Phe Gln Thr Leu Gly Gly Asp Leu Asp Gly Leu Gln
 865 870 875 880
 Gly Leu Arg Ser Leu Glu Ile Leu Arg Ile Arg Lys Val Asn Gly Leu
 885 890 895
 Ala Arg Ile Lys Gly Leu Lys Asp Leu Leu Cys Ser Ser Thr Cys Lys
 900 905 910
 Leu Arg Lys Phe Tyr Ile Thr Glu Cys Pro Asp Leu Ile Glu Leu Leu
 915 920 925
 Pro Cys Glu Leu Gly Val Gln Thr Val Val Val Pro Ser Met Ala Glu
 930 935 940
 Leu Thr Ile Arg Asp Cys Pro Arg Leu Glu Val Gly Pro Met Ile Arg
 945 950 955 960
 Ser Leu Pro Lys Phe Pro Met Leu Lys Lys Leu Asp Leu Ala Val Ala
 965 970 975
 Asn Ile Thr Lys Glu Glu Asp Leu Asp Ala Ile Gly Ser Leu Glu Glu
 980 985 990
 Leu Val Ser Leu Glu Leu Glu Leu Asp Asp Thr Ser Ser Gly Ile Glu
 995 1000 1005
 Arg Ile Val Ser Ser Ser Lys Leu Gln Lys Leu Thr Thr Leu Val Val
 1010 1015 1020
 Lys Val Pro Ser Leu Arg Glu Ile Glu Gly Leu Glu Glu Leu Lys Ser
 1025 1030 1035 1040
 Leu Gln Asp Leu Tyr Leu Glu Gly Cys Thr Ser Leu Gly Arg Leu Pro
 1045 1050 1055
 Leu Glu Lys Leu Lys Glu Leu Asp Ile Gly Gly Cys Pro Asp Leu Thr
 1060 1065 1070
 Glu Leu Val Gln Thr Val Val Ala Val Pro Ser Leu Arg Gly Leu Thr
 1075 1080 1085
 Ile Arg Asp Cys Pro Arg Leu Glu Val Gly Pro Met Ile Gln Ser Leu
 1090 1095 1100
 Pro Lys Phe Pro Met Leu Asn Glu Leu Thr Leu Ser Met Val Asn Ile
 1105 1110 1115 1120
 Thr Lys Glu Asp Glu Leu Glu Val Leu Gly Ser Leu Glu Glu Leu Asp
 1125 1130 1135
 Ser Leu Glu Leu Thr Leu Asp Asp Thr Cys Ser Ser Ile Glu Arg Ile
 1140 1145 1150
 Ser Phe Leu Ser Lys Leu Gln Lys Leu Thr Thr Leu Ile Val Glu Val
 1155 1160 1165
 Pro Ser Leu Arg Glu Ile Glu Gly Leu Ala Glu Leu Lys Ser Leu Arg
 1170 1175 1180

- 109 -

Ile Leu Tyr Leu Glu Gly Cys Thr Ser Leu Glu Arg Leu Trp Pro Asp
 1185 1190 1195 1200

Gln Gln Gln Leu Gly Ser Leu Lys Asn
 1205

(2) INFORMATION FOR SEQ ID NO:108:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1143 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:108:

Met Ala Ser Ser Ser Ser Ser Arg Trp Ser Tyr Asp Val Phe Leu
 1 5 10 15
 Ser Phe Arg Gly Glu Asp Thr Arg Lys Thr Phe Thr Ser His Leu Tyr
 20 25 30
 Glu Val Leu Asn Asp Lys Gly Ile Lys Thr Phe Gln Asp Asp Lys Arg
 35 40 45
 Leu Glu Tyr Gly Ala Thr Ile Pro Gly Glu Leu Cys Lys Ala Ile Glu
 50 55 60
 Glu Ser Gln Phe Ala Ile Val Val Phe Ser Glu Asn Tyr Ala Thr Ser
 65 70 75 80
 Arg Trp Cys Leu Asn Glu Leu Val Lys Ile Met Glu Cys Lys Thr Arg
 85 90 95
 Phe Lys Gln Thr Val Ile Pro Ile Phe Tyr Asp Val Asp Pro Ser His
 100 105 110
 Val Arg Asn Gln Lys Glu Ser Phe Ala Lys Ala Phe Glu Glu His Glu
 115 120 125
 Thr Lys Tyr Lys Asp Asp Val Glu Gly Ile Gln Arg Trp Arg Ile Ala
 130 135 140
 Leu Asn Glu Ala Ala Asn Leu Lys Gly Ser Cys Asp Asn Arg Asp Lys
 145 150 155 160
 Thr Asp Ala Asp Cys Ile Arg Gln Ile Val Asp Gln Ile Ser Ser Lys
 165 170 175
 Leu Cys Lys Ile Ser Leu Ser Tyr Leu Gln Asn Ile Val Gly Ile Asp
 180 185 190
 Thr His Leu Glu Lys Ile Glu Ser Leu Leu Glu Ile Gly Ile Asn Gly
 195 200 205
 Val Arg Ile Met Gly Ile Trp Gly Met Gly Gly Val Gly Lys Thr Thr
 210 215 220
 Ile Ala Arg Ala Ile Phe Asp Thr Leu Leu Gly Arg Met Asp Ser Ser

- 110 -

225		230		235		240
Tyr Gln Phe Asp Gly Ala Cys Phe Leu Lys Asp Ile Lys Glu Asn Lys						
		245		250		255
Arg Gly Met His Ser Leu Gln Asn Ala Leu Leu Ser Glu Leu Leu Arg						
		260		265		270
Glu Lys Ala Asn Tyr Asn Asn Glu Glu Asp Gly Lys His Gln Met Ala						
		275		280		285
Ser Arg Leu Arg Ser Lys Lys Val Leu Ile Val Leu Asp Asp Ile Asp						
		290		295		300
Asn Lys Asp His Tyr Leu Glu Tyr Leu Ala Gly Asp Leu Asp Trp Phe						
		305		310		315
Gly Asn Gly Ser Arg Ile Ile Ile Thr Thr Arg Asp Lys His Leu Ile						
		325		330		335
Glu Lys Asn Asp Ile Ile Tyr Glu Val Thr Ala Leu Pro Asp His Glu						
		340		345		350
Ser Ile Gln Leu Phe Lys Gln His Ala Phe Gly Lys Glu Val Pro Asn						
		355		360		365
Glu Asn Phe Glu Lys Leu Ser Leu Glu Val Val Asn Tyr Ala Lys Gly						
		370		375		380
Leu Pro Leu Ala Leu Lys Val Trp Gly Ser Leu Leu His Asn Leu Arg						
		385		390		395
Leu Thr Glu Trp Lys Ser Ala Ile Glu His Met Lys Asn Asn Ser Tyr						
		405		410		415
Ser Gly Ile Ile Asp Lys Leu Lys Ile Ser Tyr Asp Gly Leu Glu Pro						
		420		425		430
Lys Gln Gln Glu Met Phe Leu Asp Ile Ala Cys Phe Leu Arg Gly Glu						
		435		440		445
Glu Lys Asp Tyr Ile Leu Gln Ile Leu Glu Ser Cys His Ile Gly Ala						
		450		455		460
Glu Tyr Gly Leu Arg Ile Leu Ile Asp Lys Ser Leu Val Phe Ile Ser						
		465		470		475
Glu Tyr Asn Gln Val Gln Met His Asp Leu Ile Gln Asp Met Gly Lys						
		485		490		495
Tyr Ile Val Asn Phe Gln Lys Asp Pro Gly Glu Arg Ser Arg Leu Trp						
		500		505		510
Leu Ala Lys Glu Val Glu Glu Val Met Ser Asn Asn Thr Gly Thr Met						
		515		520		525
Ala Met Glu Ala Ile Trp Val Ser Ser Tyr Ser Ser Thr Leu Arg Phe						
		530		535		540
Ser Asn Gln Ala Val Lys Asn Met Lys Arg Leu Arg Val Phe Asn Met						
		545		550		555
Gly Arg Ser Ser Thr His Tyr Ala Ile Asp Tyr Leu Pro Asn Asn Leu						

565

570

575

Arg	Cys	Phe	Val	Cys	Thr	Asn	Tyr	Pro	Trp	Glu	Ser	Phe	Pro	Ser	Thr
			580					585					590		
Phe	Glu	Leu	Lys	Met	Leu	Val	His	Leu	Gln	Leu	Arg	His	Asn	Ser	Leu
		595					600					605			
Arg	His	Leu	Trp	Thr	Glu	Thr	Lys	His	Leu	Pro	Ser	Leu	Arg	Arg	Ile
	610					615					620				
Asp	Leu	Ser	Trp	Ser	Lys	Arg	Leu	Thr	Arg	Thr	Pro	Asp	Phe	Thr	Gly
625					630					635					640
Met	Pro	Asn	Leu	Glu	Tyr	Val	Asn	Leu	Tyr	Gln	Cys	Ser	Asn	Leu	Glu
				645					650					655	
Glu	Val	His	His	Ser	Leu	Gly	Cys	Cys	Ser	Lys	Val	Ile	Gly	Leu	Tyr
			660					665					670		
Leu	Asn	Asp	Cys	Lys	Ser	Leu	Lys	Arg	Phe	Pro	Cys	Val	Asn	Val	Glu
		675					680					685			
Ser	Leu	Glu	Tyr	Leu	Gly	Leu	Arg	Ser	Cys	Asp	Ser	Leu	Glu	Lys	Leu
	690					695					700				
Pro	Glu	Ile	Tyr	Gly	Arg	Met	Lys	Pro	Glu	Ile	Gln	Ile	His	Met	Gln
705					710					715					720
Gly	Ser	Gly	Ile	Arg	Glu	Leu	Pro	Ser	Ser	Ile	Phe	Gln	Tyr	Lys	Thr
				725					730					735	
His	Val	Thr	Lys	Leu	Leu	Leu	Trp	Asn	Met	Lys	Asn	Leu	Val	Ala	Leu
			740					745					750		
Pro	Ser	Ser	Ile	Cys	Arg	Leu	Lys	Ser	Leu	Val	Ser	Leu	Ser	Val	Ser
		755					760					765			
Gly	Cys	Ser	Lys	Leu	Glu	Ser	Leu	Pro	Glu	Glu	Ile	Gly	Asp	Leu	Asp
	770					775					780				
Asn	Leu	Arg	Val	Phe	Asp	Ala	Ser	Asp	Thr	Leu	Ile	Leu	Arg	Pro	Pro
785					790					795					800
Ser	Ser	Ile	Ile	Arg	Leu	Asn	Lys	Leu	Ile	Ile	Leu	Met	Phe	Arg	Gly
				805					810					815	
Phe	Lys	Asp	Gly	Val	His	Phe	Glu	Phe	Pro	Pro	Val	Ala	Glu	Gly	Leu
			820					825					830		
His	Ser	Leu	Glu	Tyr	Leu	Asn	Leu	Ser	Tyr	Cys	Asn	Leu	Ile	Asp	Gly
		835					840					845			
Gly	Leu	Pro	Glu	Glu	Ile	Gly	Ser	Leu	Ser	Ser	Leu	Lys	Lys	Leu	Asp
	850					855					860				
Leu	Ser	Arg	Asn	Asn	Phe	Glu	His	Leu	Pro	Ser	Ser	Ile	Ala	Gln	Leu
865					870					875					880
Gly	Ala	Leu	Gln	Ser	Leu	Asp	Leu	Lys	Asp	Cys	Gln	Arg	Leu	Thr	Gln
				885					890					895	
Leu	Pro	Glu	Leu	Pro	Pro	Glu	Leu	Asn	Glu	Leu	His	Val	Asp	Cys	His

- 112 -

900								905				910			
Met	Ala	Leu	Lys	Phe	Ile	His	Tyr	Leu	Val	Thr	Lys	Arg	Lys	Lys	Leu
		915					920					925			
His	Arg	Val	Lys	Leu	Asp	Asp	Ala	His	Asn	Asp	Thr	Met	Tyr	Asn	Leu
	930					935					940				
Phe	Ala	Tyr	Thr	Met	Phe	Gln	Asn	Ile	Ser	Ser	Met	Arg	His	Asp	Ile
945					950					955					960
Ser	Ala	Ser	Asp	Ser	Leu	Ser	Leu	Thr	Val	Phe	Thr	Gly	Gln	Pro	Tyr
				965					970					975	
Pro	Glu	Lys	Ile	Pro	Ser	Trp	Phe	His	His	Gln	Gly	Trp	Asp	Ser	Ser
			980					985					990		
Val	Ser	Val	Asn	Leu	Pro	Glu	Asn	Trp	Tyr	Ile	Pro	Asp	Lys	Phe	Leu
		995					1000					1005			
Gly	Phe	Ala	Val	Cys	Tyr	Ser	Arg	Ser	Leu	Ile	Asp	Thr	Thr	Ala	His
	1010					1015					1020				
Leu	Ile	Pro	Val	Cys	Asp	Asp	Lys	Met	Ser	Arg	Met	Thr	Gln	Lys	Leu
1025					1030					1035					1040
Ala	Leu	Ser	Glu	Cys	Asp	Thr	Glu	Ser	Ser	Asn	Tyr	Ser	Glu	Trp	Asp
				1045					1050					1055	
Ile	His	Phe	Phe	Phe	Val	Pro	Phe	Ala	Gly	Leu	Trp	Asp	Thr	Ser	Lys
			1060					1065					1070		
Ala	Asn	Gly	Lys	Thr	Pro	Asn	Asp	Tyr	Gly	Ile	Ile	Arg	Leu	Ser	Phe
		1075					1080					1085			
Ser	Gly	Glu	Glu	Lys	Met	Tyr	Gly	Arg	Leu	Arg	Leu	Tyr	Lys	Glu	Gly
	1090					1095					1100				
Pro	Glu	Val	Asn	Ala	Leu	Leu	Gln	Met	Arg	Glu	Asn	Ser	Asn	Glu	Pro
1105					1110					1115					1120
Thr	Glu	His	Ser	Thr	Gly	Ile	Arg	Arg	Thr	Gln	Tyr	Asn	Asn	Arg	Thr
				1125					1130					1135	
Ser	Phe	Tyr	Glu	Leu	Ile	Asn									
			1140												

(2) INFORMATION FOR SEQ ID NO:109:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 429 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:109:

Leu Arg Ser Lys Leu Asp Leu Ile Ile Asp Leu Lys His Gln Ile Glu
1 5 10 15

- 113 -

Ser Val Lys Glu Gly Leu Leu Cys Leu Arg Ser Phe Ile Asp His Phe
 20 25 30
 Ser Glu Ser Tyr Val Glu His Asp Glu Ala Cys Gly Leu Ile Ala Arg
 35 40 45
 Val Ser Val Met Ala Tyr Lys Ala Glu Tyr Val Ile Asp Ser Cys Leu
 50 55 60
 Ala Tyr Ser His Pro Leu Trp Tyr Lys Val Leu Trp Ile Ser Glu Val
 65 70 75 80
 Leu Glu Asn Ile Lys Leu Val Asn Lys Val Val Gly Glu Thr Cys Glu
 85 90 95
 Arg Arg Asn Thr Glu Val Thr Val His Glu Val Ala Lys Thr Thr Thr
 100 105 110
 Asn Val Ala Pro Ser Phe Ser Ala Tyr Thr Gln Arg Ala Asn Glu Glu
 115 120 125
 Met Glu Gly Phe Gln Asp Thr Ile Asp Glu Leu Lys Asp Lys Leu Leu
 130 135 140
 Gly Gly Ser Pro Glu Leu Asp Val Ile Ser Ile Val Gly Met Pro Gly
 145 150 155 160
 Leu Gly Lys Thr Thr Leu Ala Lys Lys Ile Tyr Asn Asp Pro Glu Val
 165 170 175
 Thr Ser Arg Phe Asp Val His Ala Gln Cys Val Val Thr Gln Leu Tyr
 180 185 190
 Ser Trp Arg Glu Leu Leu Leu Thr Ile Leu Asn Asp Val Leu Glu Pro
 195 200 205
 Ser Asp Arg Asn Glu Lys Glu Asp Gly Glu Ile Ala Asp Glu Leu Arg
 210 215 220
 Arg Phe Leu Leu Thr Lys Arg Phe Leu Ile Leu Ile Asp Asp Val Trp
 225 230 235 240
 Asp Tyr Lys Val Trp Asp Asn Leu Cys Met Cys Phe Ser Asp Val Ser
 245 250 255
 Asn Arg Ser Arg Ile Ile Leu Thr Thr Arg Leu Asn Asp Val Ala Glu
 260 265 270
 Tyr Val Lys Cys Glu Ser Asp Pro His His Leu Arg Leu Phe Arg Asp
 275 280 285
 Asp Glu Ser Trp Thr Leu Leu Gln Lys Glu Val Phe Gln Gly Glu Ser
 290 295 300
 Cys Pro Pro Glu Leu Glu Asp Val Gly Phe Glu Ile Ser Lys Ser Cys
 305 310 315 320
 Arg Gly Leu Pro Leu Ser Val Val Leu Val Ala Gly Val Leu Lys Gln
 325 330 335
 Lys Lys Lys Thr Leu Asp Ser Trp Lys Val Val Glu Gln Ser Leu Ser
 340 345 350

- 114 -

Ser	Gln	Arg 355	Ile	Gly	Ser	Leu	Glu 360	Glu	Ser	Ile	Ser	Ile 365	Ile	Gly	Phe
Ser	Tyr 370	Lys	Asn	Leu	Pro	His 375	Tyr	Leu	Lys	Pro	Cys 380	Phe	Leu	Tyr	Phe
Gly 385	Gly	Phe	Leu	Gln	Gly 390	Lys	Asp	Ile	His	Asp 395	Ser	Lys	Met	Thr	Lys 400
Leu	Trp	Val	Ala	Glu 405	Glu	Phe	Val	Gln	Ala 410	Asn	Asn	Glu	Lys	Gly 415	Gln
Glu	Asp	Thr	Arg 420	Thr	Arg	Phe	Leu	Gly 425	Arg	Ser	Tyr	Trp			

(2) INFORMATION FOR SEQ ID NO:110:

(i) SEQUENCE CHARACTERISTICS:

- SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 11 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:110:

Gly Met Gly Gly Ile Gly Lys Thr Thr Thr Ala
1 5 10

(2) INFORMATION FOR SEQ ID NO:111:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 11 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:111:

Gly Met Gly Gly Val Gly Lys Thr Thr Ile Ala
1 5 10

(2) INFORMATION FOR SEQ ID NO:112:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 11 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:112:

- 115 -

Gly Met Pro Gly Leu Gly Lys Thr Thr Leu Ala
1 5 10

(2) INFORMATION FOR SEQ ID NO:113:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 11 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:113:

Gly Pro Gly Gly Val Gly Lys Thr Thr Leu Met
1 5 10

(2) INFORMATION FOR SEQ ID NO:114:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 11 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:114:

Phe Lys Ile Leu Val Val Leu Asp Asp Val Asp
1 5 10

(2) INFORMATION FOR SEQ ID NO:115:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 11 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:115:

Lys Lys Val Leu Ile Val Leu Asp Asp Ile Asp
1 5 10

(2) INFORMATION FOR SEQ ID NO:116:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 11 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

- 116 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:116:

Lys Arg Phe Leu Ile Leu Ile Asp Asp Val Trp
1 5 10

(2) INFORMATION FOR SEQ ID NO:117:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 11 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:117:

Lys Arg Phe Leu Leu Leu Leu Asp Asp Val Trp
1 5 10

(2) INFORMATION FOR SEQ ID NO:118:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 8 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:118:

Ser Arg Phe Ile Ile Thr Ser Arg
1 5

(2) INFORMATION FOR SEQ ID NO:119:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 8 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:119:

Ser Arg Ile Ile Ile Thr Thr Arg
1 5

(2) INFORMATION FOR SEQ ID NO:120:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 8 amino acids
(B) TYPE: amino acid

- 117 -

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:120:

Ser Arg Ile Ile Leu Thr Thr Arg
1 5

(2) INFORMATION FOR SEQ ID NO:121:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 3 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:121:

Thr Thr Arg
1

(2) INFORMATION FOR SEQ ID NO:122:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 8 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:122:

Gly Leu Pro Leu Thr Leu Lys Val
1 5

(2) INFORMATION FOR SEQ ID NO:123:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 8 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:123:

Gly Leu Pro Leu Ala Leu Lys Val
1 5

- 118 -

(2) INFORMATION FOR SEQ ID NO:124:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 8 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:124:

Gly Leu Pro Leu Ser Val Val Leu
1 5

(2) INFORMATION FOR SEQ ID NO:125:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 8 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:125:

Gly Leu Pro Leu Ala Leu Ile Thr
1 5

(2) INFORMATION FOR SEQ ID NO:126:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 7 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:126:

Lys Ile Ser Tyr Asp Ala Leu
1 5

(2) INFORMATION FOR SEQ ID NO:127:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 7 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

- 119 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:127:

Lys Ile Ser Tyr Asp Gly Leu
1 5

(2) INFORMATION FOR SEQ ID NO:128:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 7 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:128:

Gly Phe Ser Tyr Lys Asn Leu
1 5

(2) INFORMATION FOR SEQ ID NO:129:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 7 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:129:

Val Phe Leu Ser Phe Arg Gly
1 5

(2) INFORMATION FOR SEQ ID NO:130:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 8 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:130:

Pro Ile Phe Tyr Met Val Asp Pro
1 5

(2) INFORMATION FOR SEQ ID NO:131:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 8 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

- 120 -

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:131:

Pro Ile Phe Tyr Asp Val Asp Pro
1 5

(2) INFORMATION FOR SEQ ID NO:132:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 6 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:132:

Val Gly Ile Asp Asp His
1 5

(2) INFORMATION FOR SEQ ID NO:133:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 6 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:133:

Val Gly Ile Asp Thr His
1 5

(2) INFORMATION FOR SEQ ID NO:134:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 7 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:134:

Phe Leu Asp Ile Ala Cys Phe
1 5

(2) INFORMATION FOR SEQ ID NO:135:

- 121 -

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 9 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: protein

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:135:

Met His Asp Gln Leu Arg Asp Met Gly
1 5

- (2) INFORMATION FOR SEQ ID NO:136:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 9 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: protein

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:136:

Met His Asp Leu Ile Gln Asp Met Gly
1 5

- (2) INFORMATION FOR SEQ ID NO:137:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 9 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: protein

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:137:

Met His Asp Leu Ile Gln Asp Met Gly
1 5

- (2) INFORMATION FOR SEQ ID NO:138:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 6 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: protein

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:138:

- 122 -

Ser Lys Leu Glu Ser Leu
1 5

(2) INFORMATION FOR SEQ ID NO:139:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 8 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:139:

Gly Leu His Ser Leu Glu Tyr Leu
1 5

(2) INFORMATION FOR SEQ ID NO:140:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 8 base pairs
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:140:

Gly Leu Arg Ser Leu Glu Ile Leu
1 5

(2) INFORMATION FOR SEQ ID NO:141:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 3432 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:141:

ACAAGTAAAA GAAAGAGCGA GAAATCATCG AAATGGATTT CATCTCATCT CTTATCGTTG	60
GCTGTGCTCA GGTGTTGTGT GAATCTATGA ATATGGCGGA GAGAAGAGGA CATAAGACTG	120
ATCTTAGACA AGCCATCACT GATCTTGAAA CAGCCATCGG TGA CT TGAAG GCCATACGTG	180
ATGACCTGAC TTTACGGATC CAACAAGACG GTCTAGAGGG ACGAAGCTGC TCAAATCGTG	240
CCAGAGAGTG GCTTAGTGCG GTGCAAGTAA CGGAGACTAA AACAGCCCTA CTTTTAGTGA	300
GGTTTAGGCG TCGGGAACAG AGGACGCGAA TGAGGAGGAG ATACCTCAGT TGTTTCGGTT	360
GTGCCGACTA CAACTGTGC AAGAAGGTTT CTGCCATATT GAAGAGCATT GGTGAGCTGA	420
GAGAACGCTC TGAAGCTATC AAAACAGATG GCGGGTCAAT TCAAGTAACT TG TAGAGAGA	480
TACCCATCAA GTCCGTTGTC GGAAATACCA CGATGATGGA ACAGGTTTTG GAATTTCTCA	540

- 123 -

GTGAAGAAGA AGAAAGAGGA ATCATTGGTG TTTATGGACC TGGTGGGGTT GGGAAGACAA	600
CGTTAATGCA GAGCATTAAAC AACGAGCTGA TCACAAAAGG ACATCAGTAT GATGTACTGA	660
TTTGGGTTCA AATGTCCAGA GAATTCGGCG AGTGTTACAAT TCAGCAAGCC GTTGGAGCAC	720
GGTTGGGTTT ATCTTGGGAC GAGAAGGAGA CCGGCGAAAA CAGAGCTTTG AAGATATACA	780
GAGCTTTGAG ACAGAAACGT TTCTTGTGTG TGCTAGATGA GTCTGGGAAG ACATAGACTT	840
GGAGAAAACCT GGAGTTCCTC GACCTTGACA GGGAAAACAA ATGCAAGGTG ATGTTCACGA	900
CACGGTCTAT AGCATTATGC AACAATATGG GTGCGGAATA CAAGTTGAGA GTGGAGTTTC	960
TGGAGAAGAA ACACGCGTGG GAGCTGTTCT GTAGTAAGGT ATGGAGAAAA GATCTTTTAG	1020
AGTCATCATC AATTCGCCGG CTCGCGGAGA TTATAGTGAG TAAATGTGGA GGATTGCCAC	1080
TAGCGTTGAT CACTTTAGGA GGAGCCATGG CTCATAGAGA GACAGAAGAA GAGTGGATCC	1140
ATGCTAGTGA AGTTCTGACT AGATTTCAG CAGAGATGAA GGGTATGAAC TATGTATTTG	1200
CCCTTTTGAA ATTCAGCTAC GACAACCTCG AGAGTGATCT GCTTCGGTCT TGTTCCTGT	1260
ACTGCGCTTT ATTCCCAGAA GAACATTGTA TAGAGATCGA GCAGCTTGTT CAGTACTGGG	1320
TCGGCGAAGG GTTCTCACC AGCTCCCATG GCGTTAACAC CATTTACAAG GGATATTTTC	1380
TCATTGGGGA TCTGAAAGCG GCATGTTTGT TGGAAACCGG AGATGAGAAA ACACAGGTGA	1440
AGATGCATAA TGTGGTCAGA AGCTTTGCAT TGTGGATGGC ATCTGAACAG GGGACTTATA	1500
AGGAGCTGAT CCTAGTTGAG CCTAGCATGG GACATACTGA AGCTCCTAAA GCAGAAAACCT	1560
GGCGACAAGC TTGGTGATCT CATTGTTAGA TAACAGAATC CAGACCTTGC CTGAAAAACT	1620
CATATGCCCCG AAACAGACAA CACTGATGCT CCAACAGAAC AGCTCTTTGA AGAAGATTCC	1680
AACAGGGTTT TTCATGCATA TGCCTGTTCT CAGAGTCTTG GACTTGTCGT TCACAAGTAT	1740
CACTGAGATT CCGTTGTCTA TCAAGTATTT GGTGGAGTTG TATCATCTGT CTATGTCAGG	1800
AACAAAGATA AGTGTATTGC CACAGGAGCT TGGGAATCTT AGAAAACCTGA AGCATCTGGA	1860
CCTACAAAGA ACTCAGTTTC TTCAGACGAT CCCACGAGAT GCCATATGTT GGCTGAGCAA	1920
GCTCGAGGTT CTGAACTTGT ACTACAGTTA CGCCGGTTGG GAACTGCAGA GCTTTGGAGA	1980
AGATGAAGCA GAAGAACTCG GATTCGCTGA CTTGGAATAC TTGGAAAACC TAACCACACT	2040
CGGTATCACT GTTCTCTCAT TGGAGACCCT AAAAATCTC TTCGAGTTCTG GTGCTTTGCA	2100
TAAACATATA CAGCATCTCC ACGTTGAAGA GTGCAATGAA CTCCTCTACT TCAATCTCCC	2160
ATCACTCACT AACCATGGCA GGAACCTGAG AAGACTTAGC ATTAAAAGTT GCCATGACTT	2220
GGAGTACCTG GTCACACCCG CAGATTTTGA AAATGATTGG CTTCCGAGTC TAGAGGTTCT	2280
GACGTTACAC AGCCTTCACA ACTTAACCAG AGTGTGGGGA AATTCTGTAA GCCAAGATTG	2340
TCTGCGGAAT ATCCGTTGCA TAAACATTTT ACACTGCAAC AAGCTGAAGA ATGTCTCATG	2400
GGTTCAGAAA CTCCCAAAGC TAGAGGTGAT TGAAGTGTTC GACTGCAGAG AGATAGAGGA	2460

- 124 -

ATTGATAAGC GAACACGAGA GTCCATCCGT CGAAGATCCA ACATTGTTCC CAAGCCTGAA 2520
 GACCTTGAGA ACTAGGGATC TGCCAGAACT AAACAGCATC CTCCCATCTC GATTTTCATT 2580
 CCAAAAAGTT GAAACATTAG TCATCACAAA TTGCCCCAGA GTTAAGAAAC TGCCGTTTCA 2640
 GGAGAGGAGG ACCCAGATGA ACTTGCCAAC AGTTTATTGT GAGGAGAAAT GGTGGAAAGC 2700
 ACTGGAAAAA GTTGAAACAT TAGTCATCAC AAATTGCCCC AGAGTTAAGA AACTGCCGTT 2760
 TCAGGAGAGG AGGACCCAGA TGAAGTTGCC AACAGTTTAT TGTGAGGAGA AATGGTGGAA 2820
 AGCACTGGAA AAAGATCAAC CAAACGAAGA GCTTTGTTAT TTACCGCGCT TTGTTCCAAA 2880
 TTGATATAAG AGCTAAGAGC ACTCTGTACA AATATGTCCA TTCATAAGTA GCAGGAAGCC 2940
 AGGAAGGTTG TTCCAGTGAA GTCATCAACT TTCCACTAGA CCACAAAACCT AGAGATTATG 3000
 TAATCATAAA AACCAAATA TCCGCGATCA AATAGATCTC ACGACTATGA GGACGAAGAC 3060
 TCACCGAGTA TCGTCGATAT AGAACTCCA AGCTCCAGTT CCGATCAGTG AAGACGAACA 3120
 AGTTTATCAG ATCTCTGCAA CAATTCTGGG AATCGTCACC TCAGATTAGA CCTCCAGTAA 3180
 GAAGTGAGAA AGCATGGACG ACGACTGTGA AGAATTGAGC TAATGAGCTG AACCGGATCC 3240
 GGTGAAATTG CAGAACCGGA TCGGAGAAGA AGAATTTTGC ATTTGTGCAT CTTTATTTTT 3300
 AATTGTTACG TTTGAGCCCC AATAATCATA GATATTGTAG TGAAGACCAA ATTCATGGT 3360
 GGATCAATCA AATTGTATTT TCAAATTTTC GTAGTGTAAT AACCGAAAAA GGAATAAAAA 3420
 GGTCACCTGAG TA 3432

(2) INFORMATION FOR SEQ ID NO:142:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 909 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:142:

Met Asp Phe Ile Ser Ser Leu Ile Val Gly Cys Ala Gln Val Leu Cys
 1 5 10 15
 Glu Ser Met Asn Met Ala Glu Arg Arg Gly His Lys Thr Asp Leu Arg
 20 25 30
 Gln Ala Ile Thr Asp Leu Glu Thr Ala Ile Gly Asp Leu Lys Ala Ile
 35 40 45
 Arg Asp Asp Leu Thr Leu Arg Ile Gln Gln Asp Gly Leu Glu Gly Arg
 50 55 60
 Ser Cys Ser Asn Arg Ala Arg Glu Trp Leu Ser Ala Val Gln Val Thr
 65 70 75 80
 Glu Thr Lys Thr Ala Leu Leu Leu Val Arg Phe Arg Arg Arg Glu Gln

- 125 -

85										90					95						
Arg	Thr	Arg	Met	Arg	Arg	Arg	Tyr	Leu	Ser	Cys	Phe	Gly	Cys	Ala	Asp						
			100					105					110								
Tyr	Lys	Leu	Cys	Lys	Lys	Val	Ser	Ala	Ile	Leu	Lys	Ser	Ile	Gly	Glu						
		115					120					125									
Leu	Arg	Glu	Arg	Ser	Glu	Ala	Ile	Lys	Thr	Asp	Gly	Gly	Ser	Ile	Gln						
	130					135					140										
Val	Thr	Cys	Arg	Glu	Ile	Pro	Ile	Lys	Ser	Val	Val	Gly	Asn	Thr	Thr						
145					150					155					160						
Met	Met	Glu	Gln	Val	Leu	Glu	Phe	Leu	Ser	Glu	Glu	Glu	Glu	Arg	Gly						
			165						170					175							
Ile	Ile	Gly	Val	Tyr	Gly	Pro	Gly	Gly	Val	Gly	Lys	Thr	Thr	Leu	Met						
		180					185						190								
Gln	Ser	Ile	Asn	Asn	Glu	Leu	Ile	Thr	Lys	Gly	His	Gln	Tyr	Asp	Val						
		195					200					205									
Leu	Ile	Trp	Val	Gln	Met	Ser	Arg	Glu	Phe	Gly	Glu	Cys	Thr	Ile	Gln						
	210					215					220										
Gln	Ala	Val	Gly	Ala	Arg	Leu	Gly	Leu	Ser	Trp	Asp	Glu	Lys	Glu	Thr						
225					230					235					240						
Gly	Glu	Asn	Arg	Ala	Leu	Lys	Ile	Tyr	Arg	Ala	Leu	Arg	Gln	Lys	Arg						
			245						250					255							
Phe	Leu	Leu	Leu	Leu	Asp	Asp	Val	Trp	Glu	Glu	Ile	Asp	Leu	Glu	Lys						
			260					265					270								
Thr	Gly	Val	Pro	Arg	Pro	Asp	Arg	Glu	Asn	Lys	Cys	Lys	Val	Met	Phe						
		275					280					285									
Thr	Thr	Arg	Ser	Ile	Ala	Leu	Cys	Asn	Asn	Met	Gly	Ala	Glu	Tyr	Lys						
	290					295					300										
Leu	Arg	Val	Glu	Phe	Leu	Glu	Lys	Lys	His	Ala	Trp	Glu	Leu	Phe	Cys						
305					310					315					320						
Ser	Lys	Val	Trp	Arg	Lys	Asp	Leu	Leu	Glu	Ser	Ser	Ser	Ile	Arg	Arg						
			325						330					335							
Leu	Ala	Glu	Ile	Ile	Val	Ser	Lys	Cys	Gly	Gly	Leu	Pro	Leu	Ala	Leu						
		340						345					350								
Ile	Thr	Leu	Gly	Gly	Ala	Met	Ala	His	Arg	Glu	Thr	Glu	Glu	Glu	Trp						
		355					360					365									
Ile	His	Ala	Ser	Glu	Val	Leu	Thr	Arg	Phe	Pro	Ala	Glu	Met	Lys	Gly						
	370					375					380										
Met	Asn	Tyr	Val	Phe	Ala	Leu	Leu	Lys	Phe	Ser	Tyr	Asp	Asn	Leu	Glu						
385					390					395					400						
Ser	Asp	Leu	Leu	Arg	Ser	Cys	Phe	Leu	Tyr	Cys	Ala	Leu	Phe	Pro	Glu						
			405						410				415								
Glu	His	Ser	Ile	Glu	Ile	Glu	Gln	Leu	Val	Glu	Tyr	Trp	Val	Gly	Glu						

- 126 -

420							425					430				
Gly	Phe	Leu	Thr	Ser	Ser	His	Gly	Val	Asn	Thr	Ile	Tyr	Lys	Gly	Tyr	
		435					440					445				
Phe	Leu	Ile	Gly	Asp	Leu	Lys	Ala	Ala	Cys	Leu	Leu	Glu	Thr	Gly	Asp	
	450					455					460					
Glu	Lys	Thr	Gln	Val	Lys	Met	His	Asn	Val	Val	Arg	Ser	Phe	Ala	Leu	
465					470					475					480	
Trp	Met	Ala	Ser	Glu	Gln	Gly	Thr	Tyr	Lys	Glu	Leu	Ile	Leu	Val	Glu	
				485					490					495		
Pro	Ser	Met	Gly	His	Thr	Glu	Ala	Pro	Lys	Ala	Glu	Asn	Trp	Arg	Gln	
			500				505					510				
Ala	Leu	Val	Ile	Ser	Leu	Leu	Asp	Asn	Arg	Ile	Gln	Thr	Leu	Pro	Glu	
		515					520					525				
Lys	Leu	Ile	Cys	Pro	Lys	Leu	Thr	Thr	Leu	Met	Leu	Gln	Gln	Asn	Ser	
	530					535					540					
Ser	Leu	Lys	Lys	Ile	Pro	Thr	Gly	Phe	Phe	Met	His	Met	Pro	Val	Leu	
545					550					555					560	
Arg	Val	Leu	Asp	Leu	Ser	Phe	Thr	Ser	Ile	Thr	Glu	Ile	Pro	Leu	Ser	
				565					570					575		
Ile	Lys	Tyr	Leu	Val	Glu	Leu	Tyr	His	Leu	Ser	Met	Ser	Gly	Thr	Lys	
			580					585					590			
Ile	Ser	Val	Leu	Pro	Gln	Glu	Leu	Gly	Asn	Leu	Arg	Lys	Leu	Lys	His	
		595					600					605				
Leu	Asp	Leu	Gln	Arg	Thr	Gln	Phe	Leu	Gln	Thr	Ile	Pro	Arg	Asp	Ala	
	610					615					620					
Ile	Cys	Trp	Leu	Ser	Lys	Leu	Glu	Val	Leu	Asn	Leu	Tyr	Tyr	Ser	Tyr	
625					630					635					640	
Ala	Gly	Trp	Glu	Leu	Gln	Ser	Phe	Gly	Glu	Asp	Glu	Ala	Glu	Glu	Leu	
				645					650					655		
Gly	Phe	Ala	Asp	Leu	Glu	Tyr	Leu	Glu	Asn	Leu	Thr	Thr	Leu	Gly	Ile	
			660					665					670			
Thr	Val	Leu	Ser	Leu	Glu	Thr	Leu	Lys	Thr	Leu	Phe	Glu	Phe	Gly	Ala	
		675					680					685				
Leu	His	Lys	His	Ile	Gln	His	Leu	His	Val	Glu	Glu	Cys	Asn	Glu	Leu	
	690					695					700					
Leu	Tyr	Phe	Asn	Leu	Pro	Ser	Leu	Thr	Asn	His	Gly	Arg	Asn	Leu	Arg	
705					710					715					720	
Arg	Leu	Ser	Ile	Lys	Ser	Cys	His	Asp	Leu	Glu	Tyr	Leu	Val	Thr	Pro	
				725					730					735		
Ala	Asp	Phe	Glu	Asn	Asp	Trp	Leu	Pro	Ser	Leu	Glu	Val	Leu	Thr	Leu	
			740					745					750			
His	Ser	Leu	His	Asn	Leu	Thr	Arg	Val	Trp	Gly	Asn	Ser	Val	Ser	Gln	

- 127 -

(2) INFORMATION FOR SEQ ID NO:143:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 22 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:143:

Pro Lys Ala Glu Asn Trp Arg Gln Ala Leu Val Ile Ser Leu Leu Asp
1 5 10 15
Asn Arg Ile Gln Thr Leu
20

(2) INFORMATION FOR SEQ ID NO:144:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 23 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

- 128 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:144:

Pro Glu Lys Leu Ile Cys Pro Lys Leu Thr Thr Leu Met Leu Gln Gln
1 5 10 15
Asn Ser Ser Leu Lys Lys Ile
20

(2) INFORMATION FOR SEQ ID NO:145:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 24 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:145:

Pro Thr Gly Phe Phe Met His Met Pro Val Leu Arg Val Leu Asp Leu
1 5 10 15
Ser Phe Thr Ser Ile Thr Glu Ile
20

(2) INFORMATION FOR SEQ ID NO:146:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 23 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:146:

Pro Leu Ser Ile Lys Tyr Leu Val Glu Leu Tyr His Leu Ser Met Ser
1 5 10 15
Gly Thr Lys Ile Ser Val Leu
20

(2) INFORMATION FOR SEQ ID NO:147:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 24 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:147:

- 129 -

Pro Gln Glu Leu Gly Asn Leu Arg Lys Leu Lys His Leu Asp Leu Gln
 1 5 10 15

Arg Thr Gln Phe Leu Gln Thr Ile
 20

(2) INFORMATION FOR SEQ ID NO:148:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 37 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:148:

Pro Arg Asp Ala Ile Cys Trp Leu Ser Lys Leu Glu Val Leu Asn Leu
 1 5 10 15

Tyr Tyr Ser Tyr Ala Gly Trp Glu Leu Gln Ser Phe Gly Glu Asp Glu
 20 25 30

Ala Glu Glu Leu Gly
 35

(2) INFORMATION FOR SEQ ID NO:149:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 25 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:149:

Phe Ala Asp Leu Glu Tyr Leu Glu Asn Leu Thr Thr Leu Gly Ile Thr
 1 5 10 15

Val Leu Ser Leu Glu Thr Leu Lys Thr
 20 25

(2) INFORMATION FOR SEQ ID NO:150:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 27 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

- 130 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:150:

Leu Phe Glu Phe Gly Ala Leu His Lys His Ile Gln His Leu His Val
1 5 10 15
Glu Glu Cys Asn Glu Leu Leu Tyr Phe Asn Leu
20 25

(2) INFORMATION FOR SEQ ID NO:151:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 26 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:151:

Pro Ser Leu Thr Asn His Gly Arg Asn Leu Arg Arg Leu Ser Ile Lys
1 5 10 15
Ser Cys His Asp Leu Glu Tyr Leu Val Thr
20 25

(2) INFORMATION FOR SEQ ID NO:152:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 29 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:152:

Pro Ala Asp Phe Glu Asn Asp Trp Leu Pro Ser Leu Glu Val Leu Thr
1 5 10 15
Leu His Ser Leu His Asn Leu Thr Arg Val Trp Gly Asn
20 25

(2) INFORMATION FOR SEQ ID NO:153:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 30 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: protein

- 131 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:153:

Ser Val Ser Gln Asp Cys Leu Arg Asn Ile Arg Cys Ile Asn Ile Ser
1 5 10 15
His Cys Asn Lys Leu Lys Asn Val Ser Trp Val Gln Lys Leu
20 25 30

(2) INFORMATION FOR SEQ ID NO:154:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 28 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:154:

Pro Lys Leu Glu Val Ile Glu Leu Phe Asp Cys Arg Glu Ile Glu Glu
1 5 10 15
Leu Ile Ser Glu His Glu Ser Pro Ser Val Glu Asp
20 25

(2) INFORMATION FOR SEQ ID NO:155:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 22 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:155:

Pro Thr Leu Phe Pro Ser Leu Lys Thr Leu Arg Thr Arg Asp Leu Pro
1 5 10 15
Glu Leu Asn Ser Ile Leu
20

(2) INFORMATION FOR SEQ ID NO:156:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 23 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:156:

- 132 -

Pro Ser Arg Phe Ser Phe Gln Lys Val Glu Thr Leu Val Ile Thr Asn
 1 5 10 15

Cys Pro Arg Val Lys Lys Leu
 20

(2) INFORMATION FOR SEQ ID NO:157:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5134 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:157:

AAGCTTTACA GATTGGATGA TCTCTTAATG CATGCTGAAG TGACTGCAAA AAGGTTAGCA	60
ATATTCAGTG GTTCTCGTTA TGAATATTTT ATGAACGGAA GCAGCACTGA GAAAATGAGG	120
CCCTTGTTAT CTGATTTTCT GCAAGAGATT GAGTCTGTCA AGGTAGAGTT CAGAAATGTT	180
TGCTTGCAAG TTCTGGATAT ATCACCTTTT TCCCTGACAG ATGGAGAAGG CCTTGTTAAT	240
TTCTTATTAA AAAACCAGGC CAAGGTGCCG AATGATGATG CTGTTTCTTC TGATGGAAGT	300
TTAGAGGATG CAAGCAGCAC TGAGAAAATG GGA CTTCAT CTGATTTTCT CCGAGAGATT	360
GAGTCTGTTG AGATAAAGGA GGCCAGAAAA TTATATGATC AAGTTTTGGA TGCAACACAT	420
TGTGAGACGA GTAAGCACGA TGGAAAAAGC TTTATCAACA TTATGTTAAC CCAACAGGAC	480
AAGGTGCTGG ACTATGATGC TGGTTCAGTG TCTTATCTTC TTAACCAAAT CTCAGTAGTT	540
AAAGACAAAA TATTGCACAT TGGCTCTTTA CTTGTAGATA TTGTACAGTA CCGGAATATG	600
CATATAGAAC TTACAGATCT CGCTGAACGT GTTCAAGATA AAAACTACAT TCGTTTCTTC	660
TCTGTCAAGG GTTATATTCC TGCTTGGTAT TACACACTAT ATCTCTCTGA TGTCAAGCAA	720
TTGCTTAAGT TTGTTGAGGC AGAGGTAAAG ATTATTTGTC TGAAAGTACC AGATTCTTCA	780
AGTTATAGCT TCCCTAAGAC AAATGGATTA GGATATCTCA ATTGCTTTTT AGGCAAATTG	840
GAGGAGCTTT TACGTTCTAA GCTCGATTG ATAATCGACT TAAAACATCA GATTGAATCA	900
GTCAAGGAGG GCTTATTGTG CCTAAGATCA TTCATTGATC ATTTTTCAGA AAGCTATGTT	960
GAGCATGATG AAGCTTGTGG TCTTATAGCA AGAGTTTCTG TAATGGCATA CAAGGCTGAG	1020
TATGTCATTG ACTCATGCTT GGCCTATTCT CATCCACTCT GGTACAAAGT TCTTTGGATT	1080
TCTGAAGTTC TTGAGAATAT TAAGCTTGTA AATAAAGTTG TTGGGGAGAC ATGTGAAAGA	1140
AGGAACACTG AAGTTACTGT GCATGAAGTT GCAAAGACTA CCACTAATGT AGCACCATCT	1200
TTTTTCAGCTT ATACTCAAAG AGCAAACGAA GAAATGGAGG GTTTTCAGGA TACAATAGAT	1260

- 133 -

GAATTAAAGG ATAAACTACT TGGAGGATCA CCTGAGCTTG ATGTCATCTC AATCGTTGGC	1320
ATGCCAGGAT TGGGCAAGAC TACACTAGCA AAGAAGATTT ACAATGATCC AGAAGTCACC	1380
TCTCGCTTCG ATGTCCATGC TCAATGTGTT GTGACTCAAT TATATTCATG GAGAGAGTTG	1440
TTGCTCACCA TTTTGAATGA TGTGCTTGAG CCTTCTGATC GCAATGAAAA AGAAGATGGA	1500
GAAATAGCTG ATGATCTACG CCGATTTTTG TTGACCAAGA GATTCTTGAT TCTCATTGAT	1560
GATGTGTGGG ACTATAAAGT GTGGGACAAT CTATGTATGT GCTTCAGTGA TGTTTCAAAT	1620
AGGAGTAGAA TTATCCTAAC AACCCGCTTG AATGATGTCG CCGAATATGT CAAATGTGAA	1680
AGTGATCCCC ATCATCTTCG TTTATTCAGA GATGACGAGA GTTGGACATT ATTACAGAAA	1740
GAAGTCTTTC AAGGAGAGAG CTGTCCACCT GAACTTGAAG ATGTGGGATT TGAAATATCA	1800
AAAAGTTGTA GAGGGTTGCC TCTCTCAGTT GTGTTAGTAG CTGGTGTTCT GAAACAGAAA	1860
AAGAAGACAC TAGATTCATG GAAAGTAGTA GAACAAAGTC TAAGTTCCCA GAGGATTGGC	1920
AGCTTGGAAG AGAGCATATC TATAATTGGA TTCAGTTACA AGAATTTACC AACTATCTT	1980
AAGCCTTGTT TTCTCTATTT TGGAGGATTT TTGCAGGGAA AGGATATTCA TGA CTCAAAA	2040
ATGACCAAGT TGTGGGTAGC TGAAGAGTTT GTACAAGCAA ACAACGAAAA AGGACAAGAA	2100
GATACCCGCA CAAGGTTTCT TGGACGATCT TATTGGTAGG AATCTGGTGA TGGCCATGGA	2160
GAAGAGACCT AATGCCAAGG TGAAAACGTG CCGCATT CAT GATTTGTTGC ATAAATTCTG	2220
CATGGAAAAG GCCAAACAAG AGGATTTCTT TCTCCAGATC AATAGGTAAA AAAA ACTGTA	2280
TTAATTTTAC ATTACAAAA AAAAGAACTG TATTAATTTT ACTGTATTAT GTTTATGCCA	2340
ACTCTCATTT CCATGTGTTT TCTTTTATTC AATTCAGTGG AGAAGGTGTA TTTCTGAAC	2400
GATTGGAAGA ATACCGATTG TTCGTTTATT CTTACCAAGA TGAAATTGAT CTGTGGCGCC	2460
CATCTCGCTC TAATGTCCGC TCTTTACTAT TCAATGCAAT TGATCCAGAT AACTTGTTAT	2520
GGCCGCGTGA TATCTCCTTC ATTTTTGAGA GCTTCAAGCT TGTTAAAGTG TTGGATTG	2580
AATCATTCAA CATTGGTGGT ACTTTTCCCA TTGAAACACA ATATCTAATT CAGATGAAGT	2640
ACTTTGCGGC CCAA ACTGAT GCAAATTCAA TTCCTTCATC TATAGCTAAG CTTGAAAATC	2700
TTGAGACTTT TGTGTAAGA GGATTGGGAG GAGAGATGAT ATTACCTTGT TCACTTCTGA	2760
AGATGGTGAA ATTGAGGCAT ATACATGTAA ATGATCGGGT TTCTTTTGGT TTGCGTGAGA	2820
ACATGGATGT TTAACTGGT AACTCACAAT AACCTAATTT GGAAACCTTT TCTACTCCGC	2880
GTCTCTTTTA TGGTAAAGAC GCAGAGAAGA TTTTGAGGAA GATGCCAAAA TTGAGAAAAT	2940
TGAGTTGCAT ATTTTCAGGG ACATTTGGTT ATTCAAGGAA ATTGAAGGGT AGGTGTGTTT	3000
GTTTTCCCAG ATTAGATTTT CTAAGTCACC TTGAGTCCCT CAAGCTGGTT TCGAACAGCT	3060
ATCCAGCCAA ACTTCCTCAC AAGTTCAATT TCCCCTCGCA ACTAAGGGAA CTGACTTTAT	3120
CAAAGTTCCG TCTACCTTGG ACCCAAATTT CGATCATTGC AGAACTGCCC AACTTGGTGA	3180

- 134 -

TTCTTAAGTT	ATTGCTCAGA	GCCTTTGAAG	GGGATCACTG	GGAAGTGAAA	GATTCAGAGT	3240
TCCTAGAACT	CAAATACTTA	AAACTGGACA	ACCTCAAAGT	TGTACAATGG	TCCATCTCTG	3300
ATGATGCTTT	TCCTAAGCTT	GAACATTTGG	TTTTAACGAA	ATGTAAGCAT	CTTGAGAAAA	3360
TCCCTTCTCG	TTTTGAAGAT	GCTGTTTGTC	TAAATAGAGT	TGAGGTGAAC	TGGTGCAACT	3420
GGAATGTTGC	CAATTCAGCC	CAAGATATTC	AAACTATGCA	ACATGAAGTT	ATAGCAAATG	3480
ATTCATTAC	AGTTACTATA	CAGCCTCCAG	ATTGGTCTAA	AGAACAGCCC	CTTGACTCTT	3540
AGCAAAGGTT	TGTTCTTGCT	GTGTTCATCC	AAGTGCATTT	AACATTTATT	CATTTTGTTT	3600
TACACCAGAA	CATGTTTATT	TTGCTAGTAT	TACTTGATAC	ATTAAAAGAA	ATCGAACTCA	3660
TATTTCTGCT	ACAGTCTTAA	CTTTTCTTGG	GCTTACTTGA	GGTCTAGATT	AGATCAATGG	3720
TTCATGTAAT	TTTTAATTCA	CTGTTTCATT	CAACTGTCTT	ATGATAGTTG	TGAAATGACA	3780
ATATTGTTAT	CCCTAGCCAA	ATTTATTATG	TTCAAATGAA	AACTGATGTC	ACAACTACTT	3840
TTTTGTGAAA	TGTTTTTGAA	TTTTTTGCTA	TAAAATTGAC	GAATTGACAG	CTTCTATATT	3900
TGTCAGCTAA	ACTCTTTGTC	ACCAGAAGTG	TATTTAGAAT	TACTGTGGTT	TTATGAAAGA	3960
GTTCTGTAGA	ATTTTATGCT	TTTGCAGAAT	ATAGTTTAAA	ACAACAACAC	TTCTCTGTTT	4020
CAGAGATAGC	AGAAGCTAAA	GTTCAAGGCA	TTTTGTTTAT	TTCTAGAACA	AGTGGAGTTC	4080
TTATGTTGAA	TTCTTGAAAA	GAAGAAGAAT	CAGGAGCAGG	TAAAGTTATC	TCTTTTTTATG	4140
TTTTTCTTCT	TTTAGATGTT	ATTTCTTCAT	CTTGAACGTG	AACACCGCTG	AAAGCATTTT	4200
AATAAAACCG	GAGAGAAAAA	TAAGATCTTT	TTATATAAAG	CATTATCATG	TAAATATGCC	4260
TAAATCCATA	TGGTACAACT	GTTTGACAAA	ATGATAGAGA	GGGGAGTTTT	ATAGTATAAG	4320
TAAAACAGGA	TTGAGAAAAA	AATCCTTGCA	CGATTTTCAA	TTTCTGGCCA	CATCACAATG	4380
TGTGTCAAAG	TTCCCCTCTT	TAAGTGGAAC	AAGCAATCAG	AAAAGCTCAT	TCTTATCGGT	4440
GACATACCAA	TACCAGCTGA	CTGTCTCATC	TTGGTTAACT	TAGCCTTGCT	TACTTAGACT	4500
ATTAGATTAG	TTACTAATGA	ACTGGTAAAT	TGGAACCAAA	TGTAGTTAGC	TTGATGAGCT	4560
GGTAGACATG	TATATATGAA	GATACACGCG	TAACTTTAGT	CGATGGTTAA	TTTTTCATTT	4620
TTGATTTTTT	TTCTTCACAG	AGTATATATG	AACTTGGCCT	AAAAGTTTTG	CTTCACTAAT	4680
TTAACTATTA	CCGTGGATGA	AACAAGCATG	GCAACATTTT	CAACAACATAT	CACTCAAGCA	4740
ATGTAAAAAA	TGGAGGTTCT	ACGAGCGGTA	CATGTAAGAG	TTTTGTGCAC	ACAAGAGGTT	4800
CTGAGACTTG	AACCATCCAT	GTCCAAGGCA	GTTGAGATGC	TAGTAAAGAA	AGAAGAAGAT	4860
GAGCCTGCAC	TAATTAATCT	CCCTGTATGA	ATGAGAGAAT	GAGAAAAAGA	TGGAGCTTCA	4920
TGAACCAAAA	GTTACCTTTT	TTTTTTCTTC	TTAATGGCAT	TACTTTGAAG	CACATGTTTG	4980
TTAGTTGTAA	ATTGTAATGG	TGAAGTGTTT	GTAAATATAG	GGAGTGATAT	TTGAAAGAAT	5040
GTTGTGTGTTA	TCTTTACAAA	CCGGAATCAT	TTCTGTATAA	TTTTCTTCTG	TAATTTTTTG	5100

- 135 -

TTTCGGTTTA TTCATTACTC ATTTTCAGTAA GCTT

5134

(2) INFORMATION FOR SEQ ID NO:158:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 26 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:158:

GGNATGGGNG GNNTNGGNAA RACNAC

26

(2) INFORMATION FOR SEQ ID NO:159:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:159:

NCGNGWNGTN AKDAWNCGNA

20

(2) INFORMATION FOR SEQ ID NO:160:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 17 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:160:

GGWNTBGGWA ARACHAC

17

(2) INFORMATION FOR SEQ ID NO:161:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 33 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- 136 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:161:

NRYNRDNGTN GTYTTNCCNA NNCCNNSNRK NCC

33

(2) INFORMATION FOR SEQ ID NO:162:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 26 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:162:

GGNMYNSSNG GNNTNGGNAA RACNAC

26

(2) INFORMATION FOR SEQ ID NO:163:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 13 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:163:

TYGAYGAYRT BRA

13

(2) INFORMATION FOR SEQ ID NO:164:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 16 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:164:

TYCCAVAYRT CRTCA

16

(2) INFORMATION FOR SEQ ID NO:165:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 26 base pairs
 - (B) TYPE: nucleic acid

- 137 -

- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:165:

VYMNAYRTCR TCNADNAVNA NNARNA

26

(2) INFORMATION FOR SEQ ID NO:166:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 26 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:166:

WWNMRRDTNY TNNTNBTNHT NNARNA

26

(2) INFORMATION FOR SEQ ID NO:167:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:167:

NCGNGWNGTN AKDWNCNGNA

20

(2) INFORMATION FOR SEQ ID NO:168:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:168:

NCKNSWNGTN ADDATDAATN G

21

- 138 -

(2) INFORMATION FOR SEQ ID NO:169:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:169:

NARNGGNARN CC

12

(2) INFORMATION FOR SEQ ID NO:170:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 17 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:170:

GGWYTBCCWY TBGCHYT

17

(2) INFORMATION FOR SEQ ID NO:171:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 17 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:171:

ARDGCVARWG GVARNCC

17

(2) INFORMATION FOR SEQ ID NO:172:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 24 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- 139 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:172:

NRNNWYNAVN SHNARNGGNA RNCC

24

(2) INFORMATION FOR SEQ ID NO:173:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:173:

GGNYTNCCNY TNDSNBT

17

(2) INFORMATION FOR SEQ ID NO:174:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:174:

ARRTTRTCRT ADSWRAWYTT

20

(2) INFORMATION FOR SEQ ID NO:175:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:175:

ARNYYNTYRT ANSRNANNYY

20

(2) INFORMATION FOR SEQ ID NO:176:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

- 140 -

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:176:

RRNWTWHSNT AYRANRVNY

19

(2) INFORMATION FOR SEQ ID NO:177:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:177:

GTNTTYTWN SNTTYMGRGG

20

(2) INFORMATION FOR SEQ ID NO:178:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 23 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:178:

CCNATHTTYT AYRWBGTNGA YCC

23

(2) INFORMATION FOR SEQ ID NO:179:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 17 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:179:

GTNGGNATHG AYRMNCA

17

(2) INFORMATION FOR SEQ ID NO:180:

- (i) SEQUENCE CHARACTERISTICS:

- 141 -

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:180:

RAARCANGCD ATRTCNNARRA A

21

(2) INFORMATION FOR SEQ ID NO:181:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:181:

TTYTNGAYA THGCNTGYTT

20

(2) INFORMATION FOR SEQ ID NO:182:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:182:

CCCATCYK NADNWRRTCR TGCAT

25

(2) INFORMATION FOR SEQ ID NO:183:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:183:

- 142 -

ATGCAYGAYY WNHTNMRRGA YATGGG

26

(2) INFORMATION FOR SEQ ID NO:184:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 15 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:184:

NARNSWYTYN ARYTT

15

(2) INFORMATION FOR SEQ ID NO:185:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 17 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:185:

WSNAARYTNR ARWSNYT

17

(2) INFORMATION FOR SEQ ID NO:186:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:186:

DWWYTCNARN SWNYKNARN C

21

(2) INFORMATION FOR SEQ ID NO:187:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 17 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- 143 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:187:

GGNYTNMRNW NNYTNGA

17

(2) INFORMATION FOR SEQ ID NO:188

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 13 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:188:

Leu Lys Phe Ser Tyr Asp Asn Leu Glu Ser Asp Leu Leu
1 5 10

(2) INFORMATION FOR SEQ ID NO:189:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 16 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:189:

Gly Val Tyr Gly Pro Gly Gly Val Gly Lys Thr Thr Leu Met Gln Ser
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:190:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 14 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:190:

Gly Gly Leu Pro Leu Ala Leu Ile Thr Leu Gly Gly Ala Met

(2) INFORMATION FOR SEQ ID NO:191:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 11 amino acids
 - (B) TYPE: amino acid

- 144 -

- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 2
- (D) OTHER INFORMATION: /note= "Xaa is Met or Pro"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 3
- (D) OTHER INFORMATION: /note= "Xaa is Gly or Pro"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 5
- (D) OTHER INFORMATION: /note= "Xaa is Ile, Leu or Val"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 10
- (D) OTHER INFORMATION: /note= "Xaa is Ile, Leu or Thr"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 11
- (D) OTHER INFORMATION: /note= "Xaa is Ala or Met"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:191:

Gly	Xaa	Xaa	Gly	Xaa	Gly	Lys	Thr	Thr	Xaa	Xaa
1			5						10	

(2) INFORMATION FOR SEQ ID NO:192:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 11 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 1
- (D) OTHER INFORMATION: /note= "Xaa is Phe or Lys"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 2
- (D) OTHER INFORMATION: /note= "Xaa is Arg or Lys"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 3
- (D) OTHER INFORMATION: /note= "Xaa is Ile, Val or Phe"

- 145 -

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 5
- (D) OTHER INFORMATION: /note= "Xaa is Ile, Leu or Val"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 6
- (D) OTHER INFORMATION: /note= "Xaa is Ile or Leu"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 7
- (D) OTHER INFORMATION: /note= "Xaa is Ile or Val"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 10
- (D) OTHER INFORMATION: /note= "Xaa is Ile, Leu or Val"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 11
- (D) OTHER INFORMATION: /note= "Xaa is Asp or Trp"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:192:

Xaa	Xaa	Xaa	Leu	Xaa	Xaa	Xaa	Asp	Asp	Xaa	Xaa
1			5						10	

(2) INFORMATION FOR SEQ ID NO:193:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 8 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 1
- (D) OTHER INFORMATION: /note= "Xaa is Ser or Cys"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 2
- (D) OTHER INFORMATION: /note= "Xaa is Arg or Lys"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 3
- (D) OTHER INFORMATION: /note= "Xaa is Phe, Ile or Val"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 4
- (D) OTHER INFORMATION: /note= "Xaa is Ile or Met"

(ix) FEATURE:

- 146 -

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 5
- (D) OTHER INFORMATION: /note= "Xaa is Ile, Leu or Phe"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 7
- (D) OTHER INFORMATION: /note= "Xaa is Ser, Cys or Thr"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:193:

Xaa Xaa Xaa Xaa Xaa Thr Xaa Arg
1 5

(2) INFORMATION FOR SEQ ID NO:194:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 8 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 5
- (D) OTHER INFORMATION: /note= "Xaa is Thr, Ala or Thr"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 6
- (D) OTHER INFORMATION: /note= "Xaa is Leu or Val"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 7
- (D) OTHER INFORMATION: /note= "Xaa is Ile, Val or Lys"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 8
- (D) OTHER INFORMATION: /note= "Xaa is Val or Thr"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:194:

Gly Leu Pro Leu Xaa Xaa Xaa Xaa
1 5

(2) INFORMATION FOR SEQ ID NO:195:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

- 147 -

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 1
- (D) OTHER INFORMATION: /note= "Xaa is Lys or Gly"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 2
- (D) OTHER INFORMATION: /note= "Xaa is Ile or Phe"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 5
- (D) OTHER INFORMATION: /note= "Xaa is Asp or Lys"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 6
- (D) OTHER INFORMATION: /note= "Xaa is Ala, Gly or Asn"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:195:

Xaa Xaa Ser Tyr Xaa Xaa Leu
1 5

(2) INFORMATION FOR SEQ ID NO:196:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 4 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:196:

Asn Ser His Arg
1

(2) INFORMATION FOR SEQ ID NO:197:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:197:

Arg Asp Arg Arg Arg Val Asp Pro Cys
1 5

(2) INFORMATION FOR SEQ ID NO:198:

- 148 -

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 4 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:198:

Thr Gly Asp Leu

1

(2) INFORMATION FOR SEQ ID NO:199:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 4 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:199:

His Gly Thr Tyr

1

(2) INFORMATION FOR SEQ ID NO:200:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 11 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:200:

Arg Met Ser His Gly Phe Arg Asn Ser Gln Ser
1 5 10

(2) INFORMATION FOR SEQ ID NO:201:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 27 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

- 149 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:201:

Gly	Glu	Met	Val	Glu	Ser	Thr	Gly	Lys	Arg	Ser	Thr	Lys	Arg	Arg	Ala
1				5				10					15		
Leu	Leu	Phe	Thr	Ala	Leu	Cys	Ser	Lys	Leu	Ile					
				20				25							

- 150 -

Claims

1. A substantially pure oligonucleotide comprising the sequence:
5' GGNATGGGNGGNNTNGGNAARACNAC 3', [SEQ. ID NO:158]
5 wherein N is A, T, G, or C; and R is A or G.
2. A substantially pure oligonucleotide comprising the sequence:
5' NARNGGNARNCC 3', [SEQ. ID NO: 169] wherein N is A, T, G or C; and R is A or G.
- 10 3. A substantially pure oligonucleotide comprising the sequence:
5' NCGNGWNGTNAKDAWNCNGA 3', [SEQ. ID NO: 159]
wherein N is A, T, G or C; W is A or T; D is A, G, or T; and K is G or T.
- 15 4. A substantially pure oligonucleotide comprising the sequence:
5' GGWNTBGGWAARACHAC 3', [SEQ ID NO: 160] wherein N is A, T, G or C; R is G or A; B is C, G, or T; H is A, C, or T; and W is A or T.
- 20 5. A substantially pure oligonucleotide comprising the sequence:
5' TYGAYGAYRTBKRBRA 3', [SEQ. ID NO: 163] wherein R is G or A; B is C, G, or T; D is A, G, or T; Y is T or C; and K is G or T.
- 25 6. A substantially pure oligonucleotide comprising the sequence:
5' TYCCA VAYRTCRTCNA 3', [SEQ ID NO: 164] wherein N is A, T, G or C; R is G or A; V is G or C or A; and Y is T or C.

- 151 -

7. A substantially pure oligonucleotide comprising the sequence:

5' GGWYTBCCWYTBGCHYT 3', [SEQ ID NO.: 170] wherein B is C, G, or T; H is A, C, or T; W is A or T; and Y is T or C.

8. A substantially pure oligonucleotide comprising the sequence:

5' ARDGCVARWGGVARNCC 3', [SEQ ID NO: 171] wherein N is A, T, G or C; R is G or A; W is A or T; D is A, G, or T; and V is G, C, or A.

9. A substantially pure oligonucleotide comprising the sequence:

5' ARRTTRTCRTADSWRAWYTT 3', [SEQ ID NO: 174] wherein R is G or A; W is A or T; D is A, G, or T; S is G or C; and Y is C or T.

10. A recombinant plant gene comprising the DNA sequence:

5' GGNATGGGNGGNNTNGGNAARACNAC 3', [SEQ ID NO: 158] wherein N is A, T, G or C; and R is A or G.

20 11. The gene of claim 10, further comprising the sequence:

5' NARNGGNARNCC 3', [SEQ ID NO: 169] wherein N is A, T, G or C; and R is A or G.

25 12. The gene of claim 11, further comprising the sequence:

5' NCGNGWNGTNAKDAWNCNGNA 3', [SEQ ID NO: 167] wherein N is A, T, G or C; W is A or T; D is A, G or T; and K is G or T.

- 152 -

13. A recombinant plant gene comprising a combination of any two or more sequences of claims 10, 11 and 12.

14. A substantially pure plant polypeptide
5 comprising the amino acid sequence:

Gly Xaa₁ Xaa₂ Gly Xaa₃ Gly Lys Thr Thr Xaa₄ Xaa₅,
[SEQ ID NO: 191], wherein Xaa₁ is Met or Pro; Xaa₂ is Gly
or Pro; Xaa₃ is Ile, Leu, or Val; Xaa₄ is Ile, Leu, or
Thr; and Xaa₅ is Ala or Met.

15. A substantially pure plant polypeptide
10 comprising the amino acid sequence:

Xaa₁ Xaa₂ Xaa₃ Leu Xaa₄ Xaa₅ Xaa₆ Asp Asp Xaa₇
Xaa₈, [SEQ ID NO: 192],
wherein Xaa₁ is Phe or Lys; Xaa₂ is Arg or Lys; Xaa₃ is
15 Ile, Val, or Phe; Xaa₄ is Ile, Leu, or Val; Xaa₅ is Ile or
Leu; Xaa₆ is Ile or Val; Xaa₇ is Ile, Leu, or Val; and
Xaa₈ is Asp or Trp.

16. A substantially pure plant polypeptide
comprising the amino acid sequence:

20 Xaa₁ Xaa₂ Xaa₃ Xaa₄ Xaa₅ Thr Xaa₆ Arg, [SEQ ID NO:
193]

wherein Xaa₁ is Ser or Cys; Xaa₂ is Arg or Lys; Xaa₃ is
Phe, Ile, or Val; Xaa₄ is Ile, or Met; Xaa₅ is Ile, Leu,
or Phe; Xaa₆ is Ser, Cys, or Thr.

25 17. A substantially pure plant polypeptide
comprising the amino acid sequence:

Gly Leu Pro Leu Xaa₁ Xaa₂ Xaa₃ Xaa₄, [SEQ ID NO:
194],

wherein Xaa₁ is Thr, Ala, or Ser; Xaa₂ is Leu or Val; Xaa₃
30 is Ile, Val, or Lys; and Xaa₄ is Val or Thr.

- 153 -

18. A substantially pure plant polypeptide comprising the amino acid sequence:

Xaa₁ Xaa₂ Ser Tyr Xaa₃ Xaa₄ Leu, {SEQ ID NO: 195},
wherein Xaa₁ is Lys or Gly; Xaa₂ is Ile or Phe; Xaa₃ is
5 Asp or Lys; and Xaa₄ is Ala, Gly, or Asn.

19. A method of isolating a disease-resistance gene or fragment thereof from a plant cell, comprising:

- (a) providing a sample of plant cell DNA;
- (b) providing a pair of oligonucleotides
10 having sequence homology to a conserved region of an RPS disease-resistance gene;
- (c) combining said pair of oligonucleotides with said plant cell DNA sample under conditions suitable for polymerase chain reaction-mediated DNA amplification;
15 and
- (d) isolating said amplified disease-resistance gene or fragment thereof.

20. The method of claim 19, wherein said amplification is carried out using a reverse-
20 transcription polymerase chain reaction.

21. The method of claim 19, wherein said reverse-transcription polymerase chain reaction is RACE.

22. A method of identifying a plant disease-resistance gene in a plant cell, comprising:

- 25 (a) providing a preparation of plant cell DNA;
- (b) providing a detectably-labelled DNA sequence having homology to a conserved region of an RPS gene;
- (c) contacting said preparation of plant cell DNA with said detectably-labelled DNA sequence under
30 hybridization conditions providing detection of genes having 50% or greater sequence identity; and

- 154 -

(d) identifying a disease-resistance gene by its association with said detectable label.

23. The method of claim 22, wherein said DNA sequence is produced according to the method of claim 19.

5 24. The method of claim 22, wherein said preparation of plant cell DNA is isolated from a plant genome.

 25. A method of isolating a disease-resistance gene
10 from a recombinant plant cell library, comprising:
 (a) providing a recombinant plant cell library;
 (b) contacting said recombinant plant cell library with a detectably-labelled gene fragment produced according to the method of claim 19 under hybridization
15 conditions providing detection of genes having 50% or greater sequence identity; and
 (c) isolating a member of a disease-resistance gene by its association with said detectable label.

 26. A method of isolating a disease-resistance
20 gene from a recombinant plant cell library, comprising:
 (a) providing a recombinant plant cell library;
 (b) contacting said recombinant plant cell library with a detectably-labelled oligonucleotide of any of claims 1-9 under hybridization conditions providing
25 detection of genes having 50% or greater sequence identity; and
 (c) isolating a disease-resistance gene by its association with said detectable label.

- 155 -

27. A recombinant plant polypeptide capable of conferring disease-resistance wherein said plant polypeptide comprises a P-loop domain or nucleotide binding site domain.

5 28. The recombinant plant polypeptide of claim 27, wherein said polypeptide further comprises a leucine-rich repeating domain.

29. A recombinant plant polypeptide capable of conferring disease-resistance wherein said plant
10 polypeptide contains a leucine-rich repeating domain.

30. A plant disease-resistance gene isolated according to the method comprising:

- (a) providing a sample of plant cell DNA;
- (b) providing a pair of oligonucleotides having
15 sequence homology to a conserved region of an RPS disease-resistance gene;
- (c) combining said pair of oligonucleotides with said plant cell DNA sample under conditions suitable for polymerase chain reaction-mediated DNA amplification; and
20 (d) isolating said amplified disease-resistance gene or fragment thereof.

31. A plant disease-resistance gene isolated according to the method comprising:

- (a) providing a preparation of plant cell DNA;
- 25 (b) providing a detectably-labelled DNA sequence having homology to a conserved region of an RPS gene;
- (c) contacting said preparation of plant cell DNA with said detectably-labelled DNA sequence under hybridization conditions providing detection of genes
30 having 50% or greater sequence identity; and

- 156 -

(d) identifying a disease-resistance gene by its association with said detectable label.

32. A plant disease-resistance gene according to the method comprising:

- 5 (a) providing a recombinant plant cell library;
 (b) contacting said recombinant plant cell library with a detectably-labelled gene fragment produced according to the method of claims 1-4 under hybridization conditions providing detection of genes having 50% or
10 greater sequence identity; and
 (c) isolating a disease-resistance gene by its association with said detectable label.

33. A method of identifying a plant disease-resistance gene comprising:

- 15 (a) providing a plant tissue sample;
 (b) introducing by biolistic transformation into said plant tissue sample a candidate plant disease-resistance gene;
 (c) expressing said candidate plant disease-
20 resistance gene within said plant tissue sample; and
 (d) determining whether said plant tissue sample exhibits a disease-resistance response, whereby a response identifies a plant disease-resistance gene.

34. The method of claim 33, wherein said plant
25 tissue sample comprises leaf, root, flower, fruit, or stem tissue.

35. The method of claim 33, wherein said candidate plant disease-resistance gene is obtained from a cDNA expression library.

- 157 -

36. The method of claim 33, wherein said disease-resistance response is the hypersensitive response.


37. A plant disease-resistance gene isolated according to the method comprising:

- 5 (a) providing a plant tissue sample;
- (b) introducing by biolistic transformation into said plant tissue sample a candidate plant disease-resistance gene;
- (c) expressing said candidate plant disease-
10 resistance gene within said plant tissue sample; and
- (d) determining whether said plant tissue sample exhibits a disease-resistance response, whereby a response identifies a plant disease-resistance gene.

38. A purified antibody which binds specifically
15 to an rps family protein.

39. A DNA sequence substantially identical to the DNA sequence shown in Figure 12.

40. A substantially pure polypeptide having a sequence substantially identical to a Prf amino acid
20 sequence shown in Figure 5 (A or B).



1
AAGTAAAGAAAGAGCGAGAAATCATCGAAATCGATTTTCATCTCTCTTATCGCTGGC
-----+ 60
TTCAATTTCTTTCTCGCTCTTTTAGTAGCTTTACCTAAAGTAGAGTAGAGAAATAGCAACCG
a
b K * K K E R E I I E (M) D F I S S L I V G -
c S K R K S E K S S K W I S S H L L S L A -
V K E R A R N H R N G F H L I S Y R W L -
61
TGTGCTCAGGTGTTGTGTGCAATCTATGAAATATGGCGGAGAGAAGAGGACATAAGACTGAT
-----+ 120
ACACGAGTCCACAAACACACTTAGATACTTATACCGGCTCTCTTCTCCTGTATTCTGACTA
a
b C A Q V L C E S M N M A E R R G H K T D -
c V L R C C V N L * I W R R E E D I R L I -
C S G V V * I Y E Y G G E K R T * D * S -
121
CTTAGACAAGCCATCACTGATCTTCAAAACAGCCATCGGCTGACTTGAAGGCCATACCTGAT
-----+ 180
GAATCTGTTTCGGTAGTGACTAGAACTTTGTTCGGTAGCCACTGAACCTTCGGGTATGCCACTA
a
b L R Q A I T D L E T A I G D L K A I R D -
c L D K P S L I L K Q P S V T * R P Y V M -
* T S H H * S * N S H R * L E C H T * * -
181
GACCTGACTTTACGGATCCAAACAAGACGGTCTAGAGGGACGAAGCTGCTCAAAATCGTGGC
-----+ 240
CTGGACTGAAAATGCCTAGGTTGTTCTGCCAGATCTCCCTGCTTCGACCACTTACGACCG
a
b D L T L R I Q Q D G L E G R S C S N R A -
c T * L Y G S N K T V * R D E A A Q I V P -
P D F T D P T R R S R G T K L L K S C Q -
241
AGAGAGTGGCTTAGTCCGGTGCAACTAACGGAGACTAAAACAGCCCTACTTTTACTGAGG
-----+ 300
TCTCTCAGCGAATCAGGCCACGTTTCATTGCTCTGATTTTGTGCGGAAGAAAATCACTCC
a
b R E W L S A V Q V T E T K T A L L L V R -
c E S G L V R C K * R R L K Q P Y F * * G -
R V A * C C A S N G D * N S P T F S E V -
301
TTTAGGGCTGGGGAACAGAGGACGGGAATGAGGAGGAGATACCTCAGTTGTTTCGGTTGT
-----+ 360
AAATCCGGAGCGCTTGTCTCGCTGGGCTTACTGCTGCTCTATGGAGTCACAAAGCCCAACA
a
b F R R R E Q R T R M R R R Y L S C F G C -
c L G V G N R G R E * G G D T S V V S V V -
* A S G T E D A N E E E I P Q D F R L C -
GCGGACTACAAACTGTGCAAGAAGGTTTCTGCCATATTGAAGAGCATTCGTGAGGTGAG

FIGURE 2

```

361 ----- 420
CGGCTCATGTTTACACGTTTCTTCCAAAGACGGTATAACTTTCTGTAACCACTCGACTCT
a   A D Y K L C K K V S A I L K S I G E L R -
b   P T T N C A R R F L P Y * R A L V S * E -
c   R L Q T V Q E G F C H I E E H W * A E R -

GAACGCTCTCAAGCTATCAAAACAGATGCGCGGCTCAATTCAAGTAACTTCTAGAGAGATA
421 ----- 480
CTTCCGAGACTTCCATAGTTTTCTGTACCGCGGCACTTAAGTTCAATTGAACATCTCTCTAT
a   E R S E A I K T D G G S I Q V T C R E I -
b   N A L K L S K Q H A G Q F K * L V E R Y -
c   T L * S Y Q N R W R V N S S N L * R D T -

CCCATCAAGTCCGTTCTCGGAAATACCAAGATCATCGAACAGGTTTTCGAAATTTCTCAGT
481 ----- 540
CGGTAGTTACGCAACAGCGCTTTATGGTGTCTACTACGTTCTCGAAACGTTAAAGACTCA
a   P I K S V V G N T T M H Z Q V L E P L S -
b   P S S P L S E I P R * W N R F W N P S V -
c   H Q V R C R K Y H D D G T G F G : S Q * -

GAAGAAGAAGAAAGAGGAATCATTTGGTGTCTTATGGACCTCTCGGGTTCCGAAGACAACG
541 ----- 600
CTTCTTCTCTTTCTCTCTTAGTAACCAAAATACCTCGACCAAGCCCAAGCCTTCTCTTGC
a   E E E E R G I I G V Y G P G G V G K T T -
b   K K K K E E S L V F M D L V G L G R Q R -
c   R R R K R N H W C L W T W W C W E D N V -

TTAATGCAGAGCATTAAACAACCAAGCTCATCAAAAAGGACATCATATCATGTACTGATT
601 ----- 660
AATTACGCTCTGTAATTCTTGTCTGCTAGTAGTGTCTTCTGTAGTCATCTACATCACTAA
a   L M Q S I N N E L I T K G H Q Y D V L I -
b   * C R A L T T S * S Q K D I S M H Y * F -
c   N A E H * Q R A D H K R T S V * C T D L -

TGGGTTCAAAATCTCCAGAGAATTCCGCGAGTGTACAATTCAAGCAAGCGGTTCCAGCAACG
661 ----- 720
ACCCAAGTTTACAGGCTCTCTTAAGCCGCTCACATGTTAAGTCTTCCGCAACCTCTGCCC
a   W V Q M S R E F G E C T I Q Q A V C A R -
b   G F K C P E N S A S V Q F S K P L E H G -
c   G S N V Q R I R R V Y N S A S R W S T V -

TTGGGTTTATCTTGGGACGAGAGGAGACCGGCGGCAAAACAGAGCTTTTAAAGATATACAGA
721 ----- 780
AACCCAAATAGAACCCCTGCTCTTCTCTTGGCGCGCTTTTCTCTCGAAACCTCTATATGCT
a   L G L S W D E K E T G E N P A L K I Y R -
b   W V Y L G T R R R P A K T E L * R Y T E -
c   G F I L G R E G D R R K Q S P E D I Q S -

GCTTTGAGACAGAAACGTTTCTTGTGTTGCTAGATCATCTCTCGGAAGAGATAGACTTC
781 ----- 840
CGAAACTCTCTCTTTCCAAAGAACAAACAGATCTACTACAGACGCTTCTCTATCTGAAC
a   A L R Q K R F L L L L D D V W E E I D L -
b   L * D R N V S C C C * H H S G K R * T W -
c   F E T E T F L V V A R * C L G R D R L C -

```

FIG. 2 CONTINUED

841 GAGAAAACCTGCACTTCTCTGACCTGACAGGGAAAACAAATGCAAGGTGATGTTTCAGGACA
----- 900
CTCTTTTGACCTCAAGGAGCTGCACTGTGTCCTTTTGTGTTTACGTTCCACTACAAGTGCTGT

a E K T G V P R P D R E N K C K V H F T T -
b R K L E F L D L T G K T N A R * C S R H -
c E N W S S S T * Q G K Q M Q G D V H D T -

901 CCGTCTATACCATTTATGCAACAATATGCGTCCCGAATACAAGTTGAGAGTGGAGTTTCTC
----- 960
GCCAGATATCGTAATACGTTGTTTATACCCACCGCTTATGTTCAACTCTACCTCAAAGAC

a R S I A L C N N M G A E Y K L R V E F L -
b G L * H Y A T I W V R N T S * E W S P W -
c V Y S I M Q Q Y G C G I Q V E S C V S C -

961 GAGAAGAAAACACCGGTGGGAGCTGTTCTCTAGTAAGGTATGAGAGAAAACATCTTTTACAG
----- 1020
CTCTTCTTTGTCGGCACCCTCGACAAGACATCATTCATACCTCTTTTCTAGAAAATCTC

a E K K H A W E L F C S K V W R K D L L E -
b R R N T R G S C S V V R Y C E K I F * S -
c E E T R V G A V L * * C M E K R S F R V -

1021 TCATCATCAATTGCGCGGCTGCGCGAGATTATAGTGAGTAAATGTGGAGCATGCGCACTA
----- 1080
AGTAGTAGTTAAGCGCGCGAGCGGCTCTAATATCACTCATTTAGACCTCCTAACGGTCTAT

a S S S I R R L A E I I V S K C G G L P L -
b H H Q P A G S R R L * * V N V E D C R * -
c I I N S P A R G D Y S E * M W R I A T S -

1081 CCGTTGATCACTTTAGGAGCAGCCATGCGTCTATAGACACACAGAAGAGAGTGGATCCAT
----- 1140
CGCAACTAGTCAATCTCTCTGCGTACCGAGTATCTCTCTGTTCTCTCTACCTAGGTA

a A L I T L G G A M A H R E T E E E W I H -
b R * S L * E E P W L I E R Q K K S G S H -
c V D H F R R S H G S * R D R R E V D P C -

1141 CCTAGTGAACTTCTCACTAGATTTCCAGCAGAGATCAAGGGTATCAACTATGTATTTGCC
----- 1200
CGATCACTTCAAGACTGATCTAAAGCTGCTCTCTACTTCCCATACTTCATACATAAACCG

a A S E V L T R F P A E M K G M N Y V F A -
b L V K P * L D P Q Q R * R V * T M Y L P -
c * * S S D * I S S R D E G Y E L C I C P -

1201 CTTTTGAAATTCAGCTACGACAACCTCGAGAGTCACTCTGCTTCCGTCTGTTTCTGTAC
----- 1260
GAAAACCTTAAAGTCGATGCTGTTTGGAGCTCTCACTAGACGAAGCCAGAACAAAGAACATG

a L L K F S Y D N L E S D L L R S C P L Y -
b F * N S A T T T S R V I C F G L V S C T -
c F E I Q L R Q P R E * S A S V L F L V L -

1261 TCGCGCTTTATTCGAGAGAAGCAATTCTATAGAGATCGAGCAGCTTGTGAGTACTGGGTC
----- 1320
ACGCGAAATAAGGGTCTTTTCTTAAGATATCTCTAGCTCTGCAACAACCTCATGACCCAC

a C A L F P E E H S I E I E Q L V E Y W V -

FIG. 2 CONTINUED

b A L Y S Q K N I L * R S S S L L S T G S -
 c R P I P R R T F Y R D R A A C * V L G R -
 1321 ----- 1380
 GCGGAAGCGTTTCTCACCAGCTCCCATGCCCTTAACACCATTTACAAAGCGATATTTTCTC
 CCGCTTCCCAAGAGTGGTCGAGGGTACCGCAATTGTGGTAAATGTTCCCTATAAAAGAG
 a G E G F L T S S H G V N T I Y K G Y F L -
 b A K G F S P A P M A L T P F T R D I F S -
 c R R V S H Q L P W R * H H L Q G I F S H -
 1381 ----- 1440
 ATTGGGGATCTGAAAGCGGCATGTTTCTTCCAAACCGGAGATGAGAAAACACAGCTCAAG
 TAACCCCTAGACTTTCCCGGTACAAACAACCTTTGGCCTCTACTCTTTGTGTCCACTTC
 a I C D L K A A C L L E T G D E K T Q V K -
 b L G I * K R H V C W K P E M R K H R * R -
 c W G S E S G M F V G N R R * E N T G E D -
 1441 ----- 1500
 ATGCATAATGTGGTCAGAAGCTTTGCCATTGTGGATGGCATCTGAACAGGGGACTTATAAG
 TACGTATTACACCAGCTCTTCCGAAACGTAACACCTACCGTAGACTTGTCCCTCAATATTC
 a M H N V V R S F A L W M A S E Q G T Y K -
 b C I M W S E A L H C G W H L N R G L I R -
 c A * C G Q K L C I V D G I * T G D L * G -
 1501 ----- 1560
 GAGCTGATCCTAGTTGAGCCTAGCATGGGACATACTGAAGCTCTAAAGCAGAAAACCTGG
 CTGCACTAGCATCAACTCCGATCGTACCCTGTATGACTTCCAGGATTTCTGTCTTTTGACC
 a E L I L V E P S M G H T E A P K A E N W -
 b S * S * L S L A W D I L K L L K Q K T C -
 c A D P S * A * H G T Y * S S * S R K L A -
 1561 ----- 1620
 CGACAAGCGTTGGTGATCTCATTTGTTAGATAACAGAAATCCAGACCTTGGCTGAAAAACTC
 GCTGTTCCCAACCACTAGAGTAACAATCTATTGTCTTAGGTCTGGAACCGACTTTTTCAG
 a R Q A L V I S L L D N R I Q T L P E K L -
 b D K R W * S H C * I T E S R P C L R N S -
 c T S V G D L I V R * Q N P D L A * R T H -
 1621 ----- 1680
 ATATCCCCGAAACTGACAACACTGATGCTCCAACAGAACAGCTCTTTGAAGAAGATTCCA
 TATACCGGCTTTGACTGTTGTGACTACGAGCTTGTCTTGTCCGAGAAACTTCTTCTAAGCT
 a I C P K L T T L M L Q Q N S S L R K X I P -
 b Y A R N * Q H * C S N R T A L * R R F Q -
 c M P E T D N T D A P T E Q L F E E D S N -
 1681 ----- 1740
 ACAGCGTTTTTCATGCATATGCCCTGTTCTCAGAGTCTTTGAGCTTGTGTTTCAAGTATC
 TGTCCCCAAAAAGTACGTATACGGACAAGAGTCTCAGAACCTGAACAGCAAGTGTTCATAG
 a T G F F M H M P V L R V L D L S F T S I -
 b Q G F S C I C L F S E S W T C R S Q V S -
 c R V F H A Y A C S Q S L G L V V H R Y R -
 1741 ----- 1800
 ACTGAGATTCCGTTGTCTATCAAGTATTTGGTGGAGTTGTATCATCTGTCTATCTCAGGA
 TGACTCTAAGGCAACAGATAGTTTCATAAACCACTCAACATAGTAGACAGATACAGTCT

FIG. 2 CONTINUED

```

a   T E I P L S I K Y L V E L Y H L S M S G -
b   L R P R C L S S I N W S C I I C L C Q E -
c   * D S V V Y Q V P G G V V S S V Y V R N -

ACAAAGATAAGTGTATTGCGACAGGAGCTTGGGAATCTTAGAAAAGTGAAGCATCTGGAC
1801 ----- 1860
TGTTCCTATTGACATAACGGTGTCTGCGAAGCGTTAGAACTCTTTGAGTTCTAGACCTG

a   T K I S V L P Q E L G N L R K L K H L D -
b   Q R * V Y C H R S L G I L E N * S I W T -
c   K D R C I A T G A W E S * K T E A S G P -

CTACAAAGAACTCAGTTTCTTCAGAGCGATGCCAGGAGATGCCATATGTTGCGCTAGCAAG
1861 ----- 1920
GATGTTTCTTGAGTCAAAGAAGTCTGCTAGGGTCTCTACCGTATACAACCGAGCTCTTC

a   L Q R T Q F L Q T I P R D A I C W L S K -
b   Y K E L S F F R R S H E M P Y V G * A S -
c   T K N S V S S D D P T R C H M L A E Q A -

CTCGAGGTTCTGAAGTTGTACTACAGTTACCGCGGTTGGGAAGTGCAGAGCTTTGGAGAA
1921 ----- 1980
GAGCTCGAGAGCTTGAACATGATGTCAATGCCGCGCAACCGTTGAGCTCTCGAAACCTCTT

a   L E V L N L Y Y S Y A G W E L Q S F G E -
b   S R P * T C T T V T P V G N C R A L E X -
c   R G S E L V L Q L R R L G T A E L W R R -

GATGAAGCAGAAAGAACTCGGATTGCGCTGACTTGGAACTCTCGAAAAAGCTAACCACACTC
1981 ----- 2040
CTACTTCTCTCTTCTTGAGCGCTAAGCGACTGAACCTTATGAACCTTTTGGATTCTCTGAG

a   D E A E E L G F A D L E Y L E N L T T L -
b   M K Q K N S D S L T W N T W K T * P H S -
c   * S R R T R I R * L G I L G K F N H T R -

CGTATCACTGTTCTCTCATTCGAGAGCGCTAAAAAGTCTCTTTCGAGTTTGGTCTTTCCAT
2041 ----- 2100
CCATAGTCACAAGAGAGTAACCTCTCGGATTTTTCAGAGAAGCTCAAGCCACGAAACGTA

a   G I T V L S L E T L R T L F E F G A L H -
b   V S L F S H W R P * K L S S S S V L C I -
c   Y H C S L I G D P K N S L R V R C F A * -

AAACATATACAGCATCTCCAGCTTGAAGAGTCCAATGAAGTCTCTTACTTCAATCTCCCA
2101 ----- 2160
TTTGTATATCTCTGAGAGCTGCAACTTCTCAGCTTACTTGAGGAGATGAAGTTAGAGCGT

a   K H I Q H L H V E E C N E L L Y F N L P -
b   N I Y S I S T L K S A M N S S T S I S H -
c   T Y T A S P R * R V Q * T P L L Q S P I -

TGACTCACTAACCATGGCAGCAACCTGAGAAGACTTAGCATTAAGATTTGCCATGACTTC
2161 ----- 2220
AGTGAGTGATTCGTACCGTCTCTTGGAGCTCTTCTGAATCCTAATTTTCAACGGTACTGAA

a   S L T N H G R N L R R L S I K S C H D L -
b   H S L T M A G T * E D L A L K V A M T W -
c   T H * P W Q E P E R T * H * K L P * L C -

GAGTACCTGCTGACACCGCGAGATTTTCAAAAATGATTCGGCTTCCGAGCTAGAGCTTCT

```

FIG. 2 CONTINUED

2221 ----- 2280
CTCATGGACCACTGTGGGGCTCTAAAACCTTTTACTAACCGAAGGCTCAGATCTCCAAAGAC
a E Y L V T P A D F E N D W L P S L E V L -
b S T W S H P Q I L K M I G F R V * R P * -
c V P G H T R R F * K * L A S E S R G S D -
ACGTTACACAGCCTTCACAACCTTAACCAGAGTGTGGGAAATTCTCTAACCGAAGATTCT
2281 ----- 2340
TCCAATCTCTCGGAAGTCTTGAATTCTCTACACCCCTTTAAGACATTCTGGTTCTAACA
a T L H S L H N L T R V W G N S V S Q D C -
b R Y T A F T T * P E C G E I L * A K I V -
c V T Q P S Q L N Q S V G K F C X P R L S -
CTGCGGAATATCCGTTGCATAAACATTTACACTGCCAACAGCTGAAGAATCTCTCATCG
2341 ----- 2400
GACCGCTTATAGGCAACGTATTTGTAAAGTGTGACGTTGTTCTGACTTCTTACAGAGTACC
a L R N I R C I N I S H C N K L K N V S W -
b C G I S V A * T F H T A T S * R M S H G -
c A E Y P L H K H F T L Q Q A E E C L M G -
GTTCAGAAACTCCCAAAGCTAGAGCTGATTTGAACCTTTCTGACTCCAGAGAGATAGAGGAA
2401 ----- 2460
CAAGTCTTTTACGGCTTTTCATCTCCACTAAGTTGACAAGCTGACCTCTCTCTATCTCTCT
a V Q K L P K L E V I E L F D C R E I E E -
b F R N S Q S * R * L N C S T A E R * R N -
c S E T P K A R G D * T V R L Q R D R G I -
TTGATAAGCGAACACGAGAGTCCATCTCTGGAAGATCCAACATTGTTTCCCAAGCCTGAAAG
2461 ----- 2520
AACTATTCTGCTTGTCTCTCAGGTAGGCAGCTTCTAGGTTGTAAACAAGGGTTCCGACTTC
a L I S E H E S P S V E D P T L P P S L K -
b * * A N T R V H P S K I Q H C S Q A * R -
c D K R T R E S I R R R S N I V F K P E D -
ACCTTGAGAAGCTAGGATCTTCCCAAGCTAAACAGGATCTCTCCATCTCCATTTTCAATTC
2521 ----- 2580
TGGAACTCTTCTATCTCTAGACCTTTTCAATTTCTCTAGGAGGCTAGAGCTAAAAGTAAAG
a T L R T R D L P E L N S I L P S R F S F -
b P * E L G I C Q N * T A S S H L D F H S -
c L E N * G S A R T K Q H P P I S I F I P -
CAAAAAGTTGAAACATTAGTCATCACAATTTGCCCCAGAGTTAAGAAAGTCCCGTTTCAG
2581 ----- 2640
GTTTTTCAACTTTGTAAATCAGTAGTGTTTAACGGGGTCTCAATTCTTTGACCGCAAGTC
a Q K V E T L V I T N C P R V K K L P P Q -
b K K L K H * S S Q I A P E L R N C R F R -
c K S * N I S H H K L P Q S * E T A V S G -
GAGAGGAGGAGCCGAGATGAACCTTGGCAACAGTTTATTGTGAGGAGAAATGGTGGAAAGCA
2641 ----- 2700
CTCTCTCTCTGGGTCTACTTGAACGGTTGTCAAAATAACACTCTCTTTTACCACCTTTCTCT
a E R R T Q M N L P T V Y C E E K W W K A -
b R G G P R * T C Q Q F I V R R N G G K H -
c E E D P D E L A N S L L * G E M V E S T -

```

2701 CTCGAAAAGATCACCACGAGAGCTTTGTTATTTACGGCGCTTTGTTCCAAATTCA
-----
GACCTTTTCTAGTTGCTTCTCTCGAAACATAAATGGCGCGAAACAAGGTTTAACT 2760
a   L E K D Q P N E E L C Y L P R F V P N . -
b   W K K I N Q T K S P V I Y R A L F Q I D -
c   G K R S T K R R A L L F T A L C S K L I -

2761 TATAAGAGCTAAGAGCACTCTCTACAAATATGTCGATTGATAAGTAGGAGGAGCCAGGA
-----
ATATTCTCGATTCTGCTGAGACATGTTTATACAGGTAAGTATTGATCCTCTTCCGCTCCT 2820
a   Y K S . E H S V Q I C P F I S S R K P G -
b   I R A K S T L Y K Y V H S . V A G S Q E -
c   . E L R A L C T N M S I H K . Q E A R K -

2821 AGGTTGTTCCAGTGAAGTCATCAACTTTCCACATAGCCACAAAAGTACAGATTATGTAAT
-----
TCCAACAAGGTCAGTTCTAGTAGTTGAAAGGTGTATCGGTTGTTTGATCTCTAATACATTA 2880
a   R L F Q . S H Q L S T . F Q N . R L C N -
b   C C S S E V I N F P H S H K T R D Y V I -
c   V V P V K S S T F H I A T K L E I M . S -

2881 CATAAAAACCAAACTATCCGCGA
-----
GTATTTTGGTTTCTATAGGCGCT 2903
a   H K N Q T I R -
b   I K T K L S A -
c   . K P N Y P R -

```

Enzymes that do cut:

NONE

Enzymes that do not cut:

KpnI

FIG. 2 CONTINUED

-146
ATCCATTGATCTCTGGCTTCACTCCCACTAGTCCATTCTAGACCACTGCTACCCCCCTCT -26
GCGCATCTATCGAGCTATTTTGGAAATTTTCCGACGGTTATGCAATTCCTAGTCCGAACCCATT -26
CATTCCTTCCAAACCACCAACCGACCAATTAACAAGCTTCTCTAGCTCATCATCAAAATT
MetLysIle 35
GCTCCAGTTCCGATAAATCAGACGCTCTCAGCAGCGAGCTCTCTTCACACGCGCGACCC 95
AlaProValAlaIleAsnHisSerProLeuSerArgGluValProSerHisAlaAlaPro
ACTCAGCCAAAGCAACCAACCTTCAATCTCAAGCTGGCGATTAGATCCAAAGAAAAGT 155
ThrGlnAlaLysGlnThrAsnLeuGlnSerGluAlaGlyAspLeuAspAlaArgLysSer
AGCCCTTCAAGCCCCGAAACCCGCTATTACTCTCTTCTAAGACAGTACTCCGACACAC 215
SerAlaSerSerProGluThrArgAlaLeuLeuAlaThrLysThrValLeuGlyArgHis
AAGATAGAGCTTCTCCGCTTTTCCAGCGTGGTTCAAAAAGAAATCATTTAGCAGCAGACG 275
LysIleGluValProAlaPheGlyGlyTrpPheLysLysLysSerSerLysHisGluThr
GCGCTTCAAGTCCCAACCGACATATTTGAGCGTGGCTTCCGATTTACCGAAAAAGCT 335
GlyGlySerSerAlaAsnAlaAspSerSerSerValAlaSerAspSerThrGluLysPro
TTGTTCCCTCTACCGACGTTCTTACGATATCCCAAGCTAATGACCGAATGGGATCTTCC 395
LeuPheArgLeuThrHisValProTyrValSerGlnGlyAsnGluArgMetGlyCysTrp
TATGCTCTGCGCAAGAAATGCTTGGGCAATCTCTGCAAGCTTGGGCTTGGCTTGGCTTGGG 455
TyrAlaCysAlaArgMetValGlyHisSerValGluAlaGlyProArgLeuGlyLeuPro
GAGCTTATGACGCAAGCGACGCGCGCGAGCTGGGCTACAAATTTTACAGATGTACAAAGC 515
GluLeuTyrGluGlyArgGluAlaProAlaGlyLeuGlnAspPheSerAspValGluArg
TTTATTGACAATCAAGCATTAAGTCTGGGTAGACCTTCCAGACATGACAGATTTACACAC 575
PheIleHisAsnGluGlyLeuThrArgValAspLeuProAspAsnGluArgPheThrHis

Figure 3.

GAAGAGTTCGGTCCACTGTTGTATTAAGCACCGCCGATTATATTTCGGTCCAAAACCTCCG 615
GluGluLeuGlyAlaLeuLeuTyrLysHisGlyProIleIlePheGlyTyrLysThrPro

AATGACAGCTCCACATGTTGGTCTCTACTGGTGTGATTAAGAGAGCTGGTCCATTACT 635
AsnAspSerTyrHisMetSerValLeuThrGlyValAspLysGluThrSerSerIleThr

TTCACGATCCCGACAGCGCCCGACCTAGCAATCCCGTGGATTACTTTAATACGGCA 755
PheHisAspProArgGlnGlyProAspLeuAlaMetProLeuAspTyrPheAsnGlnArg

TTGGCATGCCAGCTTCCACAGCCAAATGCTGTACCGCTAAGTACGAGGCTATCTTACCGT 815
LeuAlaTyrGlnValProHisAlaMetLeuTyrArgEnd

CGCCCATCATCACAAGCCCATCATGCCCGCCAGCAGCTACCTGAATCCCGTTCGGTTCCT 875
← →

CGTCCGATTTGCTGATCCCGAAGATGACCTTAAGAAATTCGGCAAGAGCTTCCTTCCT 935

CGACTCTCTACGCTTCGGATGATGATCAGCTCGTTTCCGAGCGCCGCTTCCTCATGAGCAT 995

CTGCCACAGCTGCTGCTGATGTTGTCTCTACGCTAAGCGGATTTTCACTGACAACCATGCC 1055

CAACTGCCCGCTTCGGATACGCTGATCTCTCAAGCCCGCTTGTGATGGCAGCCCGCAAGAA 1115

AAAGACATAATTCGGCGCTGTGAGCTTGTAGCTGTGTGCGCGCGCGAGCTGTGCGCGAT 1175

AAACACCGCTTCAGTCCCGATCTTCTGTGAAAGCATCAATCCCGCTTCCTGCTTCTTCGG 1235

CGAGTCACTCCCGACCAACGTCACCGACCCGACCGCAAGCTTCAGCGAGTCTCTCCCGAA 1295

CGTCCCGACCGATTCCTGATCTCTGAGAGAGGATACCTTCTCTCTGAC 1346

Figure 3 continued




Figure 4.

12 / 25

FIGURE 5A

1 50
 L6pro MSYLREVATA VALLPFIIL NGGWPHSKO STVWEDDST SEVDALDST
 NprotM
 PstP
 rps2

6 100
 L6pro NPSGSPBVE YVTLSPROP STREQTFPL YQSLRAYKIM TTADDELLK
 Nprot ABBBBBBSWS YVTLSPROE STRTTTTSHL YEVLDKRGIK TTQDDKRLY
 PstP LRSKLDLID LKHQIESVGE
 rps2 MDPISSEIVE CAGVLCZMN MAERRGHKTQ LRQAITSLET

101 150
 L6pro GZIGFNLLR AIDQSKIYF IISGTYADSK NCLMELAEIV RQZEDPARI
 Nprot CATPGEELK AIZESQTAIV VTSINYATSR NCLMELVKIM ECK.TRFKQT
 PstP GLICLRSTPD NPSZSYVHD EA CELIAKVEVM AYKAE.....
 rps2 AIGDLKQIRD DLTRIQQDG LIGRSCSNRA RFWLSAVQVT EYKTA.....

7 200
 L6pro IIPFYKVPF SDVRHQTCY KXAFKXANK Y..DGQTIQN WKDALKVGD
 Nprot VIFPITVDVF SHVTHQKEST AKAFEEHSTK YKDDVEGIOR WRIALNEAAN
 PstPYVIDS CLAYSEPLNY KVLW..... IS..... EVLENIKLV
 rps2LLIVR FRREQRIRM RRY..... LSCFCAD YKCKKVSAT

8 250
 L6pro LKQWHIGQD XQQAIAKVS ADINSHISKE NLILZ...TD EIVGIDDHET
 Nprot LKQSCDNRK TDADCIHQIV QOISSKLCXI SLST....LQ NIVGIDHLE
 PstP NGVVGETCER RNTVTVREV AKITTVAPS FBAYTCRANE EMEQFQTED
 rps2 LKSIGELNER SZAKTDEGS IOVTCREIPI KSVVG..... NTRM

① - P-loop 300
 L6pro AVLEKLSLS ENVTWGLYE MGGGKTTA KAVYNKI... SSC.FDCC
 Nprot KESALLTGI NGVRIMING MGGVQKTTA RAIFDTLLR MOSSYQFDGA
 PstP KLEKLLGGS FELDVISIV MGLGKTTA KRINDPEVT... SRFDVHAQ
 rps2 EQVLEPLSEE EERGIIQVQ MGGVQKTTA QSIMBELTK...G.....MOY

Fig. 5A (cont)

301 350
 L6pro CTIDNIRETO ENGGVVVLQK KLVSEILRID ..SGSVGTNN EIGGKRTIKE
 Nproct CFLKDIKE.. NRGGRSLON ALLSELR.. ..KXNNYNN EEDOKHOMAS
 Prip CWVTLYSWR EL..LITLND VLEP...S... ..DNWKEO GE..IADELRR
 rps2 DVLIWVOMR EF..GCTTIOQ AVGA...RLG ..LSWDEKET GENRALKIYR

351 400
 L6pro XVARFKYLVV LDDVLEKFXE EOMLGSPKDF ISO..RPFYT SPMRVLGTL
 Nproct RLRSKVLIV LDDIDAGBY LYLACGLDW FGNGRIIIT CPOKHLI...
 Prip FLLIDRLLIL IDDVVYKVM ONLCHTFSD. VNRRIIIT..FADVAEIV
 rps2 ALRCKRLLIL LDDVLEZIDL EXTGVPRPD. RKNCKVMIL..FADVAEIV

401 450
 L6pro MEN..OCHLYE VGSMKPRSL ELFSKAPFK NT...PFSY YETLANDVVD
 Nproct ..EK..NDITTYE VTALPOHESI OLFKORAFGE EV...ENEN FEXLSLEWVW
 Prip ..KC..ZSDPHH LRLPRODESW TLCKREVTGO E...SCPPZ LDDVGFZSK
 rps2 ..GA..ZYK..LR VEFLEKXWAM ELFGSKVWRK OLLSSSIRL LAET...EVS

451 500
 L6pro FEGDPLTAK VIGSLLPKOE IAV..WEDTL ZOL...ART LNLDEVDRL
 Nproct YAKGFLYALK VIGSLLHNLK LIZ..HKSAL EHM...KNN SYSGIIDNV
 Prip SCRLPLFVW LVAGVLKQK KILDSKVVZ QSL..SQR GSLEESTISII
 rps2 KCGDPLTAK VIGSLLHNLK LIZ..HKSAL EHM...KNN SYSGIIDNV

501 550
 L6pro KISYDAIPE AKZIFLDIA CFFICQ..NK EEPYMTDC NFYPASHIIF
 Nproct KISYDAIPE AKZIFLDIA CFFICQ..NK EEPYMTDC NFYPASHIIF
 Prip GFSYN..PH YLKGCFLYFG GFLGCKDIHD SKMTLWVAZ EFVQANN...
 rps2 KISYDNISSD LLRSCFLYCA LTFEHSIET ZGLVETWVGE GFLTSSHGVN

551 600
 L6pro LIQRCHIQVGDD DEFMHDLA DMREIVRE DVLEWKRRI
 Nproct LIDKSLVFIEEY NGVCHDLQ DMKXIVNQ KD..FGERSRL
 PripEK GEDTATRE..LRSY.....
 rps2 TTYKGYFLIC DLKACILIT ODERTQVWM NVVRSPALWM ASQGTYKEL

601 650
 L6pro WSAZEGIDLL LNRKGSSEKVK AISI..PKOVK YEFK..SECFI NLSZLYLHA
 Nproct WLAKEVLEVM SNNTGTNAME AIWVSSYST LRFS..NQAVK NQKRLRVEM
 Prip
 rps2 ILVERPMGHT EAPKAENWRO ALVISLENA IOTL..PEKLI CPKLTPLC

651 700
 L6pro REAMLTODFN NLLPNLWLE LFFYKHGEDD SBLNMYTWN LII..VILENS
 Nproct GRESTHYAID YLPNNLRCTV CTRYPH...E SFFSTPELW LVM..LOLHM.
 Prip
 rps2 QNSBLKRIPT OTTAMPVLR VLDSF... TSITTEPLSC MYL..VELVHL

Fig 5A(cont)

L6pro HITADDWGCN RUDSGMAERL KVVRLASNYE LYGARVR...
 NprotNSL RUCNTITKHL PSL.....RAID...
 Pref
 rps2 SMSGTXISVL PQELCNLAL KHLDLORTQY LQTIPROAIC WLSKLEVLNL

781 800
 L6pro .LSD.CHRFP KBIEVLSMCA IEMDEVQICE LKKLKTIVLK PCPIQKISGT
 Nprot .LSW.SKRUT RIFDFTQMFN LEY..VNLVQ CSNLEEVHMS LQCCSKVIGL
 Pref
 rps2 YYSY.AGWEL QSTGEDEAKZ LCFADLYLZ NLTTIGITVL SLETKLTFZ

801 850
 L6pro TPGCKKOLRE L.CIEFWGT NLRVVDIG QLSKLVKXV CGAKEVEINE
 Nprot YLNDCKSLKX P.....PCVNVESLE
 Pref
 rps2 PQALEHGHQH L.HVEICHNL LYFNLFSLTX HGNLRALST XSCMELEYLV

851 900
 L6pro PFLOLK....ELSTSR IPNLSQLDL ZVLKVYCKXO CFOMPPASP8
 Nprot Y.LGLR....SCHSLX LPFIYORCKP EI.....QIMXGSGIR
 Pref
 rps2 TPADFENDWL PSLEVLTLRS LKHLTRVWGN SVSQDCLRNI RCINISHCNK

901 950
 L6pro ZDESSVWVKV SKLKSLOLEK TRINVVVDV ASSGGHLPRY LLFTSLTYLK
 Nprot ELSSIFQYK THVTKLL...MGGQNLVAL PSSICRL...KSLVSL
 Pref
 rps2 LQVSWVQXL PKLEVLELD CKEIZELISE HESPSVZDPT LFP.SLTLR

951 1000
 L6pro IYQCTEPTWL P.GIENLENL TSELVNDIFQ TLGGDLGLO GLKSLILRI
 Nprot VSGCKLESLE PEEIGDLNL KVFASDTL.....ILRP
 Pref
 rps2 TKDLPELNSI LPSRFSQKV ETLVITNCPR VQULFQERR TQNLPTVYC

1001 1050
 L6pro RKVNGLARIX GLKDLCSST CLKKTYITE CPDLZELLPC ELGVQTVVVP
 Nprot P.....SSI TRINKLILM PRGFKDGVHF EFPFVAEGLH
 Pref
 rps2 EEKQNKALEK CGFMEEZCYL PRFVFN.....

1051 1100
 L6pro SMAELTEROC PRELVGMIR SLKFFPMKK LCLAVANIX EEDLDAIGSL
 Nprot SLEYLNL.SY CNLIDGGLPY EIGSLSSLKK LCLERNNE..ENLPSSLAQL
 Pref
 rps2

1101 1150
 L6pro ELVLSLELEL DOTSSEIERS VSSSKLYLT TLVWKVPSLR EIGLEELKS

Fig. 5A(cont)

Nprot	GALQSLDLK	DCORLTCLF	ELFFELNELH	VDCHMALNF
PrEP
eps2

	1151			1200
L6pro	LODLYLEGCT	SLQRLPLEKL	KE.....LD	IGGC7DLTEL
Nprot	IHDLVTKRK	KLHRVNLQCA	HNQTYNLF	YTFQNISSH
PrEP
eps2

	1201			1250
L6pro	RLTIRDCPR	LEVQPMIOSL	PKFFMLNELT	LSMVTITKCO
Nprot	SLTY.....	FTGQYPPEKI	PSWPHHQGD	SBVBVNLPE
PrEP
eps2

	1251			1300
L6pro	LD.SLELTLD	STESSIERIS	FLSKLQNTT	LIVEVPSLE
Nprot	FAVCYSRSLI	STTAHLIPVC	DDRMARMTC	KLALSECCTE
PrEP
eps2

	1301			1350
L6pro	RILYL.....	EGCTSLRL	WPDQQLGSL
Nprot	HEFFVPPAQL	WDTKANOKT	PNDYGIIRLS	FSGLZKATOL
PrEP
eps2

	1351			1387
L6pro	CKSLBVDHLS	ALKTTLEPPA	RITWPDQSYR
Nprot	MALLORENS	NEPTEHSTGI	RTQYDRTG	EYELING
PrEP
eps2

FIGURE 5B

691 SLYGRRVRLSDCWRFPKSIIEVLSMTAIE MDEV DIGELKGLKTLVLKFCPI 740
 666 SKVIGLYLNDCKSLKRFPCVNVESLEYLGLRSCDSLEKLPETYGKMP... 713
 741 QKISGGTFGMLKGLREL.....CLEFNWCTNLREVVAIDIGQLSSLK 761
 714EIQIHMOGSGIRELP.SSIFQYKTHVTKLLWNM....KNLV 750
 782 VLKMTGAKEVEINEFPLGLKELSTSSRIIPNLSQLLDLEVLKVYDCKDGF 831
 751 ALPSSICRLKSLVSLVSGOSKLESIPPEIGDLDNLRVFDASDTLILRP. 799
 832 MPPASPEDESSVWVKV...SKLKSLEKTRINNVVDDASSGGHLPY 876
 800PSSIIRLNKLIILMFRGFKDGVHFEFPPVAE 830
 879 LLPTSLTYLKIYQCTEPTWLPFIENLENLTSLEVNDIFQTLGGDLGL.Q 927
 831 GLHSLEYINLSYCNLID..GGLPEET.GSLSSLKKLDL..SRNNFEHLPS 875
 928 GLRSLEILRIKRVNGLARIKGLKDLLCSSTCKLPKFYITECPDLIELLFC 977
 876 SIA....QLGALQSLDLKDCORLTQLPELPPELNELHVCHMALKFIHYL 921
 978 ELGVQTVVVP SMAELTIRDCPRL.EVGFMIRSLPKFPM.....LKKLDLA 1021
 922 VTKRKGLHRVKLDDAHNDTMYNLFAYTMFQNISSMRHDISASDSLSLTVF 971
 1022 VANITKEEDLDAIGSLEELV..SLELEDDTSSGIERTVSSSKLOKLTTL 1069
 972 TGOPYPEKIPSWFHHQGWDSVSVN.....LPENWYIPDKFLGFAVCY 1014
 1070 VV.....KVPSLREIEGLEELKSLQDLYLEGCTSLGRLPLEKLELDIGG 1114
 1015 SRSLIDTTAHLIPVCDDK.....MSRMTQKLA....LSECTES 1049
 1115 CPDLTELVTQTVAVPSLRGLTIRDCPRLEVGPMIQSLPKFPMLNELTSLM 1164
 1050 SNYSEWDIHFFVFPFAGLWDTSKANGKTPNDYGIIRLSFSGEEKMYGLRL 1099
 1165 VNITKEDELEVLGSLEELDSLELTLDTCSSIERISF.LSKLOKLTTLIV 1213
 1100 LYKEGPEVNALLQMRENSNEPTEHSTGIRRTQYNNRTSFYELIN 1143
 1214 EVPSLREIEGLAELKSLRIILEGCTSLER.LWPDQQQLGSLKN 1256

FIGURE 5B (CON'T)

FIGURE 6


consensus PXXaXX LXXLXXLXaXXXX aXXa

505	PKAENW RQALVISLLD NR IQTL	
527	PEKLIK PK LTTLMLQQNSSLKKI	
550	PTGFFMHMPVLRVLDLSFTS ITEI	
574	PLSIKY LVELYHLSMSGTK ISVL	
597	PQELGN LRKLKHLDLQRTQFLQTI	
621	PRDAICWLSKLEVLNLYYSYAGWEL	QSFGZDEAEELG
658	FADLEY LENLTTLGITVLS LETL	KT
683	LFEFGALHKHIQHLHVEECNELLYF	NL
710	P SLTNHGRNLRRLSIKSDHLEYL	VT
736	PADFENDWLPSEVLTLHSLHNLTRV	WGN
765	SVSQDC LRNIRCINISHCNKLKNV	SWVQKL
795	PK LEV IELFDCREIEELISEHES	PSVED
823	PT LFPSLKTLLRTRDLPELNSI L	
845	PSRFS FQKVETLVITNCPRVKKL	

FIGURE 7

RPS2 Transient Expression Assay

Principle of the assay



Actual procedure




FIGURE 9




FIGURE 10




FIGURE 11

24 / 25

1	aagctttaca	gattgagatga	tctcttcaatg	40	catgctgaaag	tgactgcaaa	aaggttaqca	60
61	atattcagtg	gttcttcgtta	tgaatatttc		atgaacggaa	gcagcactga	gaaaatqagg	120
121	cccttgctat	ctgattttct	gcaagagatt		gaqtctgtca	aggttagagt	cagaaatgct	180
181	tgcttgcaag	ttcttgatct	atcacccttt		tccctgacag	atggagaaag	ccttgctaat	240
241	ttcttattaa	aaaaccaggc	caaggtgccc		aattgactgat	ctgtttcttc	tgatggaagt	300
301	ttagaggatg	caagcagcac	tgagaaatg		ggacttccat	ctgattttct	ccgagagatt	360
361	gagtcctgtg	agataaaaga	ggccagaaaa		ttatatgata	aagtctttgga	tgcaacacat	420
421	tgtagagcga	gttaagacaga	tggaaaaaagc		tttatcaaca	ttatgtttaa	ccaacaggac	480
481	aaggtgcttg	actatgatgc	tggttcagtc		tcttatcttc	tttaaccaaa	ctcagtagtt	540
541	aaagacaaac	tattgcacat	tggtctctta		cttgtagata	tgtagagta	ccggaatatg	600
601	catatagaac	ttacagatct	cgctgaacgt		gttcaagata	aaaactacat	ctgtttcttc	660
661	tctgtcaagg	gttatattcc	tgcttggtat		tacacactat	atctctctga	tgtagagcaa	720
721	tcgtctaaat	ttgttgaggc	agaggttaag		attatctgtc	tgaaagtacc	agattcttca	780
781	agttatagct	tccctaaagc	aaatggattg		ggatattctc	attgcttttt	aggcaaatgc	840
841	gagtagcttt	tacgttctaa	gctcgatttg		ataatcgact	taaaacatca	gattgaaatca	900
901	gtcaaggagg	gcttattgtg	cttaagatca		ttcattgata	atttttcaqa	aagctatgct	960
961	gagcatgatg	aagcttgctg	tcttatagca		agagcttctg	taattggcata	caaggtctag	1020
1021	tatgtcattg	actcatgctt	ggcttatttc		catccactct	ggtagaaaat	tcttggtgac	1080
1081	tctgaagctt	ttgagaaat	taagcttgta		aataaaatctg	ttggggagac	atgtgaaaga	1140
1141	aggaacactg	aagttactgt	gcattgaagt		gtcaagacta	ccactaatgt	agcaccatct	1200
1201	ttttcagctt	atactcaaa	agcaaacgaa		gaaatggagg	gttttcaqga	tacaatagat	1260
1261	gaattaaagg	ataaactact	tgaggtatca		cttgagcttg	atgtcatctc	aatccttggt	1320
1321	atgccaggat	tgggcaagac	tacactagca		aagaaatctt	acaatgaccc	agaagctcac	1380
1381	tctctgcttg	atgtccatgc	tcaatgtgtt		gtgactcaat	tatatctatg	gagagagctg	1440
1441	ttgctcacca	ttctgaatga	tgtgtctgag		cttctctgac	gcaatgajaa	agaagatcga	1500
1501	gaaatagctg	atgagctacg	ccgattcttg		ttgaccaaga	gattcttgat	tcttattgat	1560
1561	gatgtgtggg	actatamaag	gtgggacaa		ctatgtatgt	gcttcaqctg	tgcttcaaat	1620
1621	agtagtagaa	ttatcttaac	aaacctcttg		aattgatgtc	ccgaaatagt	caaatgtgaa	1680
1681	agtagtcccc	atcatctctg	tttattcaga		gatgacaga	gttggaatct	attacagaaa	1740
1741	gaagtccttc	aaggagagag	ctgtccacct		gaacttgaa	atgtgggatt	tgaaatatca	1800
1801	aaaagttgta	gaggttgctc	tctctcaagt		gtgttagtag	ctgtgttctt	gaaacagaaa	1860
1861	aagaaagcac	tagattcaatg	gaaagtagta		gaacaaagtc	taagttccca	gaggtatttg	1920
1921	agcttggaag	agagcatatc	tataattgga		tccagttaca	agaaattacc	acactatctt	1980
1981	aagcttctgt	ttctctatct	tgagagattt		ttgcagggaa	aggaatttca	tgactcaaaa	2040
2041	atgaccaagt	tgtgggtagc	tgaaagagtt		gtacaaagca	acaaagaaaa	aggaacaaaga	2100
2101	gatccccgca	caaggtttct	tgagcagatc		tattggtagg	aatctgggtg	tggtccatgga	2160
2161	gaagagacct	aatgccaagg	tgaaaacgtg		ccgcattcat	gattttgttc	ataaattctg	2220
2221	catggaaaa	gccaaacaa	aggatttctt		tctccagatc	aatagggtaa	aaaaactgta	2280
2281	tcaattttac	attacaaaa	aaaagaactg		tattaaattt	actgtattat	gtttatgccca	2340
2341	actctccatt	ccatgtgttc	tcttttattc		aatccagctg	agaaggtgta	tttcttgaa	2400
2401	gattggaaqa	ataccgattg	ttctgtcatt		cttaccgaqa	tgaaattgat	ctgtggcgcc	2460
2461	catctcgttc	taattgtccg	tctttactat		tcaatgcaat	tgatccagat	aatctgttat	2520
2521	ggcgcgctga	tatctctctc	atctttcaga		gcttcaagct	tgcttaaaag	ttggattctg	2580
2581	aatcattcaa	cattgtgtgt	acrtttccca		ttgaaataca	atattctaat	cagatgaaat	2640
2641	actttgcggc	ccaaactgat	gcaaatccaa		ttctttctac	tatagtctaa	cttgaaaaat	2700
2701	ttgagacttc	tgtcgttaaga	ggattgggag		gagagagatg	attaccttgt	tcaactctga	2760
2761	agatgggtgaa	attgagggcat	atacatgtaa		atgactcgggt	ttcttttggt	ttgcttgaga	2820
2821	acatggatgt	tttaactggg	aaactcaaat		tacctaatct	ggaaaccttt	tctactccgc	2880
2881	gtctctttta	tggttaagac	gcagagaaqa		ttttgagga	gatgccaaaa	ttgagaaaa	2940
2941	tgagttgcat	attttcaggg	acatttggtt		attcaagga	attgaagggt	aggtgtgttc	3000
3001	gttttccca	atttagattt	ctaaagccac		ttgagctcc	caagctgggt	tcgaacagct	3060
3061	atccagccaa	acttctctac	aagttcaatt		tccctctgca	actaaaggga	ctgactttat	3120
3121	caaaagttcc	tctaccttg	acccaaattt		cgatcattgc	agaactgccc	aaacttggtg	3180
3181	ttcttgaagt	ttgtctcaga	gcttttgaag		gggtcactg	ggaaagtga	gattcagagt	3240
3241	tcttgaact	caaatactta	aaactggaca		acctcaaaat	tgtagaatg	tccatctctg	3300
3301	atgagctctt	tcttaagctt	gaacacttg		ttttaaagaa	atgttaagcat	cttgagaaaa	3360
3361	tccctctctg	ttttgaagat	gctgtttgtc		taaaatagat	tgaggtgaa	tggtgcaact	3420
3421	ggaaatgttc	caattcagcc	caagataatt		aaactatgca	acatgaagtt	atagcaaatg	3480
3481	attcattcac	agttactata	caagctccag		attggtctaa	agaaacagccc	cttgactctt	3540
3541	agcaaaaggt	tggtcttggt	gtgttctatc		aagtcgactt	aaactttatc	catcttgctt	3600
3601	tacaccaaga	catgtttatt	ttgttagtat		tacttgatag	attaaaagaa	atctgaactca	3660
3661	tattctctgt	acagctctaa	ttttctcttg		gcttactctg	gggttagatt	agatcaaatg	3720
3721	ttcatgtaat	ttttaatca	ctgttctatt		caactgtctt	atgtagcttc	tgaaatgaca	3780
3781	atattgttat	ccccagccaa	atttattatg		ttcaaatgaa	aactgtagtc	acaactacct	3840

FIGURE 12

25 / 25

```

3841 ttttctgaaa tgtttttt tttt .gcta taaaaattgac gaattgacag ctt 32 300
3901 tgtcagctaa actcttttgc accagaagtg tttttagaat tactgtggtt ttatgaaaga 3960
3961 gttctctaga attttatgct ttgcagaat atagttttaa acaacaacac ttctctgttt 4020
4021 cagagatagc aqaagctaaa gttcaaggca ttttctttat ttctagaaca agtggagttc 4080
4081 ttatgttgaa ttcttgaaaa gaagaagaat caggagcagg taagcttacc tctttttttg 4140
4141 tttttcttct tttagatgtt atttcttcat cttagaacgtg aacaccgttg aaagcatttt 4200
4201 aataaaaaccg gagagaaaaa taagatcttt ttatataaag cattatcatg taatatgtcc 4260
4261 taaatccata tggtagcaact gtttgacaaa atgatataga ggggagtttt atagtataag 4320
4321 taaaacagga ttgagaaaaa aatccttgca cgtttttcaa ttcttggtcca catcacatg 4380
4381 tgtgtcaaaq tttctcttct taagtggaaac aagcaatcag aaaaqctcat tcttatcggc 4440
4441 gacataccaa taccagctga ctgtctcttc ttgggttaact tagccttgct tacttagact 4500
4501 attagattag ttactaatga actggttaact tggaaaccaa tgtagcttag ttgatgagct 4560
4561 ggtagacatg tatatatgaa gatcacgcgt taactttagt cgtaggctta tttttcattt 4620
4621 tcgatttttt ttcttcacag agtatatatg acttgacctt aaagtttttg ctccactaat 4680
4681 ttaactatta ccgtggatga aacaagcatg gcaacatttt caacaactat cactcaagca 4740
4741 atgtaaaaaa tggaggttct acgagcggtt catgtaaag ttttgtgcac acaagaggtt 4800
4801 ctgagacttg aaccatccat gtccaaggca gttgagatgc taqtaaaaga agaagaaagt 4860
4861 gagcttgca taattaatct ccttgtatga atgagagaaat gagaaaaaga tggagcttca 4920
4921 tgaacaaaaa gttacctttt tttttcttct ttaattggcat tactttgaaq ccatgctttg 4980
4981 ttagtgttaa attgtaatgg tgaagtgttt gtaaatataq gtagtgatgc ttgaaagaac 5040
5041 ggttctgtta tctttacaaa ccggaatcat ttctgtataa ttttcttctg taattttttg 5100
5101 tttcggttta ttcattactc atttcagtaa gctt 5134

```

1 10 20 30 40 50 60

FIGURE 12 (CON'T)

INTERNATIONAL SEARCH REPORT

national application No.

PCT/US95/04589

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C07K 14/415; C12N 15/29; C12Q 1/68

US CL : 536/23.1, 23.6, 24.3; 435/6; 530/370

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/23.1, 23.6, 24.3; 435/6; 530/370

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

MPSRCH, SDC, CAS ONLINE, IG

Search terms: SEQ ID NOS: 1-5, 105, 158, AND 191

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X ----- P, Y	Cell, Volume 78, issued 23 September 1994, S. Whitham et al., "The product of the tobacco mosaic virus resistance gene <i>N</i> : Similarity to Toll and the interleukin-1 receptor", pages 1101-1115, see sequences.	1, 10-13 ----- 14, 19-32
P, X ----- P, Y	Science, Volume 265, issued 23 September 1994, A.F. Bent et al., " <i>RPS2</i> of <i>Arabidopsis thaliana</i> : a leucine-rich repeat class of plant disease resistance genes", pages 1856-1860, see the entire document.	10-14, 19-32 ----- 1
P, X ----- P, Y	Cell, Volume 78, issued 23 September 1994, M. Mindrinos et al., "The <i>A. thaliana</i> disease resistance gene <i>RPS2</i> encodes a protein containing a nucleotide-binding site and leucine-rich repeats", pages 1089-1099, see the entire document.	10-14, 19-32 ----- 1

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

13 JULY 1995

Date of mailing of the international search report

04 AUG 1995

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

CHE SWYDEN CHERESKIN

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/04589

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Proceedings National Academy of Sciences, USA, Volume 90, issued December 1993, F. Bunz et al., "cDNAs encoding the large subunit of human replication factor C", pages 11014-11018, see peptide sequences.	14
X	Proceedings National Academy of Sciences, USA, Volume 90, issued December 1993, P.D. Burbelo, et al., "Cloning of the large subunit of activator 1 (replication factor C) reveals homology with bacterial DNA ligases", pages 11543-11547, see peptide sequences.	14
X	Biochem. Biophys. Res. Commun., Volume 193, Number 2, issued 15 June 1993, Y. Lu, et al., "Cloning and expression of a novel human DNA binding protein, PO-GA", pages 779-786, see peptide sequences.	14
Y	The Plant Cell, Volume 5, issued August 1993, B.N. Kunkel et al., "RPS2, an Arabidopsis disease resistance locus specifying recognition of <i>Pseudomonas syringae</i> strains expressing the avirulence gene <i>avrRpt2</i> ", pages 865-875, see the entire document.	1, 10-14, and 19-32
Y	Phil. Trans. R. Soc. Lond. B, Volume 342, issued 29 November 1993, C. Dean, "Advantages of Arabidopsis for cloning plant genes", pages 189-195, see especially Table 1.	1, 10-14, 19-32

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/04589

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1, 10-14, 19-32

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

Group I, claims 1-32, drawn to RPS oligos and polypeptides.

This application contains claims directed to more than one species of the generic invention. These species are deemed to lack Unity of Invention because they are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for more than one species to be examined, the appropriate additional examination fees must be paid. The species are as follows:

1. The oligos of claim 1
2. The oligos of claim 2.
3. The oligos of claim 3.
4. The oligos of claim 4.
5. The oligos of claim 5.
6. The oligos of claim 6.
7. The oligos of claim 7.
8. The oligos of claim 8.
9. The oligos of claim 9.

and

1. The peptides of claim 14.
2. The peptides of claim 15.
3. The peptides of claim 16.
4. The peptides of claim 17.
5. The peptides of claim 18.

The species listed above do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, the species lack the same or corresponding special technical features for the following reasons: each group of oligos comprise a separate and distinct chemical entity which does not share a special technical feature with any other species. Likewise, each group of peptides comprise a separate and distinct chemical entity which does not share a special technical feature with any other species of peptide.

Group II, claims 33-37, drawn to identification of plant disease resistance using biolistics.

Group III, claim 38, drawn to an antibody.

Group IV, claim 39-40, drawn to the Prf amino acid sequence.

The inventions listed as Groups I and II do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: Group I is drawn to specific products and the use of those products to isolate disease resistance genes while Group II is not drawn to the use of any specific product to isolate genes but rather to the use of biolistics to isolate genes. The use of biolistics is a different inventive concept than the use of the specific products of Group I. Therefore, Groups I and II do not share a special technical feature.

The inventions listed as Groups I and III do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: Group III is drawn to an antibody which is a product that is chemically distinct from the products of Group I which are specific oligos and peptides. Therefore, Groups I and III do not share a special technical feature.

The inventions listed as Groups I and IV do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: Group I is drawn to specific nucleotides and peptides related to RPS while Group IV relates to a Prf amino acid sequence. These appear to be separate and distinct chemical entities. Therefore, Groups I and IV do not share a special technical feature.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/04589

The inventions listed as Groups II and III do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: the antibodies of Group III could not be used in the method of Group II. Therefore, Groups II and III do not share a special technical feature.

The inventions listed as Groups II and IV do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: the Prf amino acid sequence of Group IV could not be used in the method of Group II. Therefore, Groups II and IV do not share a special technical feature.

The inventions listed as Groups III and IV do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: the antibody of Group III and the Prf amino acid sequence of Group IV appear to be separate, distinct and unrelated chemical entities. Therefore, Groups III and IV do not share a special technical feature.

Accordingly, the claims are not so linked by a special technical feature within the meaning of PCT Rule 13.2 so as to form a single inventive concept.